



RNF43 Is Frequently Mutated in Colorectal and Endometrial Cancers

Citation

Hodis, Eran. 2020. RNF43 Is Frequently Mutated in Colorectal and Endometrial Cancers. Doctoral dissertation, Harvard Medical School.

Permanent link

<https://nrs.harvard.edu/URN-3:HUL.INSTREPOS:37365198>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

RNF43 is frequently mutated
in colorectal and endometrial cancers

A THESIS PRESENTED

BY

ERAN HODIS, PH.D.

SUBMITTED IN PARTIAL FULFILLMENT OF THE REQUIREMENTS FOR THE
M.D. DEGREE WITH HONORS IN A SPECIAL FIELD AT
HARVARD MEDICAL SCHOOL

MAY 2020

RNF43 is frequently mutated in colorectal and endometrial cancers

ABSTRACT

We report somatic mutations of *RNF43* in over 18% of colorectal adenocarcinomas and endometrial carcinomas. *RNF43* encodes an E3 ubiquitin ligase that negatively regulates Wnt signaling. Truncating mutations of *RNF43* are more prevalent in microsatellite-unstable tumors and show mutual exclusivity with inactivating *APC* mutations in colorectal adenocarcinomas. These results indicate that *RNF43* is one of the most commonly mutated genes in colorectal and endometrial cancers.*

*The material in this thesis is reproduced, in accordance with *Springer Nature's* author reuse guidelines, from Giannakis, M.[#], Hodis E.[#] *et al. Nature Genetics* 2014.¹ (# denotes equal contribution).

Contents

1	GLOSSARY	1
2	INTRODUCTION	3
3	RESULTS	5
4	DISCUSSION AND SUMMARY	12
5	METHODS	14
6	FIGURES	18
7	SUPPLEMENTARY FIGURES AND TABLES	21
	REFERENCES	27

Acknowledgments

The work reproduced in this thesis has been previously published, with the following full citation:

Giannakis M*, Hodis E*, Mu XJ, Yamauchi M, Rosenbluh J, Cibulskis K, Saksena G, Lawrence MS, Qian ZR, Nishihara R, Van Allen EM, Hahn WC, Gabriel SB, Lander ES, Getz G, Ogino S, Fuchs CS, Garraway LA. RNF43 is frequently mutated in colorectal and endometrial cancers. *Nat Genet.* 2014 Dec;46(12):1264-6. *These authors contributed equally to this work.

The above publication is reproduced herein, in accordance with with *Springer Nature's* author reuse guidelines.

AUTHOR CONTRIBUTIONS: M.G., E.H., W.C.H., G.G., S.B.G., E.S.L., S.O., C.S.F. and L.A.G. designed research. M.G., E.H., X.J.M., M.Y., J.R., K.C., G.S., M.S.L., Z.R.Q., R.N. and E.M.V.A. performed research. G.G., S.B.G.,

E.S.L., S.O. and C.S.F. contributed new reagents and analytic tools. M.G., E.H., X.J.M., E.M.V.A. and L.A.G. analyzed data. M.G., E.H. and L.A.G. wrote the manuscript.

We thank S. Bahl, E. Nickerson and S. Chauvin for project management and A. Regev for help with computing resources, as well as A. Sivachenko, F. Huang and P. Tamayo for helpful discussions. This work was supported by the Dana-Farber Cancer Institute Leadership Council, the 2014 Colon Cancer Alliance-American Association for Cancer Research Fellowship for Biomarker Research, grant 14-40-40-GIAN, the Perry S. Levy Endowed Fellowship (M.G.), award T32GM007753 from the National Institute of General Medical Sciences, the Herchel Smith Fellowship (E.H.) and the Agrusa Fund for Colorectal Cancer Research (C.S.F.), as well as National Human Genome Research Institute grant U54HG003067 (E.S.L. and S.B.G.) and National Cancer Institute grants, K07CA190673 (R.N.), P01CA87969, UM1CA167552 and R01CA151993 (S.O.), R01CA118553 and R01CA168141 (C.S.F.) and P50CA127003 (M.G., W.C.H., L.A.G. and C.S.F.).

1

Glossary

- *APC*: adenomatous polyposis coli
- C•G: cytosine-guanine base pair
- CGHub: Cancer Genomics Hub
- dbGaP: database of Genotypes and Phenotypes
- dsDNA: double-stranded DNA
- FFPE: formalin-fixed paraffin-embedded

- HPFS: Health Professionals Follow-Up Study
- MSI: microsatellite instability
- MSI-H: microsatellite instability high
- MSI-L: microsatellite instability low
- MSI-NA: microsatellite instability status not available
- MSS: microsatellite stable
- NHS: Nurses' Health Study
- PCR: polymerase chain reaction
- *PTPRK*: protein tyrosine phosphatase receptor type K
- RNA-seq: bulk RNA sequencing
- *RNF43*: ring finger protein 43
- RSPO: R-spondin
- *RSPO3*: R-spondin 3
- TCGA: The Cancer Genome Atlas

2

Introduction

The E3 ubiquitin-protein ligase RNF43 negatively regulates the Wnt signaling pathway². Somatic mutations in *RNF43* have been associated with heightened sensitivity to compounds that target the Wnt-specific acyltransferase porcupine (PORCN) in preclinical models³. In parallel, clinical trials of small-molecule porcupine inhibitors (for example, LGK974) are ongoing in Wnt ligand-dependent

malignancies (melanoma, pancreatic, breast, head and neck, and colorectal cancers). *RNF43* is frequently mutated in pancreatic cystic neoplasms⁴ and in <5% of pancreatic carcinomas with acinar differentiation⁵; however, *RNF43* mutations have not been reported in melanoma⁶, breast cancer⁷, or head and neck malignancies⁸. In colorectal cancer, Wnt signaling is more commonly dysregulated through loss-of-function *APC* mutations⁹, whereas *RNF43* has not been reported to be significantly mutated in previous sequencing studies^{10,11}.

3

Results

Unexpectedly, our whole-exome sequencing of colorectal cancers identified a large number of non-silent somatic mutations in *RNF43*. DNA isolated from a discovery set of 185 formalin-fixed, paraffin-embedded colorectal tumor and matched normal samples collected from participants in 2 prospective cohort studies, the Nurses' Health Study (NHS) and the Health Professionals Follow-Up Study (HPFS)¹²,

showed *RNF43* mutations in 35 (18.9%) cases (median allelic fraction of 0.23, range of 0.01-0.68) (Supplementary Table 7.1). Frameshift mutations encoding p.Gly659fs and p.Arg117fs constituting insertions or deletions of 1 bp in homopolymeric tracts (microsatellite instability (MSI) loci) of seven and six C•G pairs, respectively, accounted for 41.7% (p.Gly659fs) and 8.3% (p.Arg117fs) of the *RNF43* mutations identified (Figure 6.1A). To exclude the possibility that these mutations represented technical artifacts, we validated 31 of the *RNF43* mutations (97% of the 32 reactions that had leftover DNA available and achieved coverage of >50× in resequencing or successfully underwent Sanger sequencing) in the mutant tumors and their matched normal tissue (Supplementary Figure 7.1 and Supplementary Table 7.1).

The unexpectedly high frequency of truncating *RNF43* mutations in our colorectal tumor cohort contrasted with the paucity of *RNF43* mutations reported by previous studies of a comparable scale, including a TCGA (The Cancer Genome Atlas) study of 224 colorectal tumor-normal tissue pairs¹⁰. We hypothesized that previous studies might have inadvertently filtered out many bona fide *RNF43* frameshift events owing to their similarity to the polymerase slip errors that may arise during the massively parallel sequencing process. Therefore, we reanalyzed 222 TCGA colorectal tumor-normal exomes (representing all TCGA colorectal ex-

omes available on our local servers in September 2013). Of these, 49 cases (22%) were described in the published TCGA study¹⁰. We discovered *RNF43* mutations with high allelic fraction at a frequency of 17.6% (median allelic fraction of 0.38, range of 0.04-0.77; 48.0% encoding p.Gly659fs and 12% encoding p.Arg117fs). We then orthogonally validated these mutations by examining matched RNA sequencing (RNA-seq) data, additionally confirming mRNA expression of the mutant *RNF43* alleles (100% validation rate, 44 of 44 mutations in cases with at least 10-fold coverage at the relevant base pair; Supplementary Table 7.2). These results confirmed that *RNF43* is mutated at a high frequency in colorectal tumors (Figure 6.1B and Supplementary Table 7.2).

In light of this discovery, we reasoned that inactivating mutations in *RNF43* might also have been overlooked in previous whole-exome sequencing studies of endometrial cancer, another Wnt-dependent tumor type in which MSI is common. A reanalysis of all 248 endometrial tumor-normal exome pairs from the published TCGA study¹³ identified the presence of non-silent *RNF43* mutations in 18.1% of cases (median allelic fraction of 0.31, range of 0.04-0.87), with the p.Gly659fs variant accounting for 47.3% and the p.Arg117fs variant accounting for 3.6% of the alterations (Figure 6.1C and Supplementary Table 7.3). Matched RNA-seq data orthogonally validated these events (91% validation rate, 20 of 22 mutations

in cases with at least 10-fold coverage of the relevant basepair; Supplementary Table 7.3).

The high frequency of truncating mutations at this locus (together with a low frequency of synonymous mutations) strongly suggested that *RNF43* mutations had undergone positive selection during colorectal and endometrial tumor evolution. To investigate this possibility, we analyzed *RNF43* in each tumor exome cohort using InVEx, an algorithm we previously developed to infer the presence of positive selection in tumors with high background mutation rates⁶. InVEx uses the exon/intron mutational distribution at a given locus to determine whether a gene has more mutations of predicted functional consequence than expected under a model of no selection. *RNF43* emerged as significantly mutated in all three sample sets (NHS/HPFS colorectal cancer, $P = 1.4 \times 10^{-7}$; TCGA colorectal cancer, $P = 4.1 \times 10^{-10}$; TCGA endometrial cancer, $P = 2.0 \times 10^{-7}$). Furthermore, given the empirical somatic indel rate at exonic homopolymeric tracts of seven C•G pairs in tumors with MSI, *RNF43* indels causing p.Gly659fs occurred significantly more often than expected by chance ($P < 2.2 \times 10^{-16}$, binomial test; 0.08-0.14% mutation frequency across all exonic 7-bp C•G tracts and 9.2-10.8% mutation frequency for the 7-bp C•G tract in *RNF43* causing p.Gly659fs, calculated in tumors with MSI). *RNF43* indels causing the p.Gly659fs alteration represented the most

recurrent indels at exonic C•G homopolymeric tracts of any length in all three cohorts. Thus, *RNF43* mutations apparently conferred a fitness advantage to the colorectal and endometrial cancer cells in which they occurred.

As noted earlier, most *RNF43* mutations (73-75% of all non-silent variants) were truncating events (defined as frameshift indels, nonsense mutations and splice-site mutations), consistent with the previous characterization of *RNF43* as a tumor-suppressor gene². Additionally, 16 of 74 (21.6%) *RNF43*-mutant colorectal tumors (Supplementary Table 7.1 and Supplementary Table 7.2) and 8 of 45 (17.8%) *RNF43*-mutant endometrial tumors (Supplementary Table 7.3) harbored at least 2 non-silent *RNF43* mutations, in line with the Knudson 2-hit hypothesis for tumor-suppressor genes. We cannot exclude the possibility that the mutations affected the same allele in some two-hit cases, but this seems less likely given the overall loss-of-function pattern.

Given that the *RNF43* mutations were predominated by small insertions or deletions in homopolymeric tracts in tumor types with a high prevalence of MSI, we hypothesized that these mutations might be enriched in MSI tumors. Indeed, truncated mutations of *RNF43* were statistically linked to the microsatellite instability high (MSI-H) phenotype in both tumor types ($P < 2.2 \times 10^{-16}$ for colorectal cancer and $P < 9.0 \times 10^{-16}$ for endometrial cancer, Fisher's exact test;

Figure 6.2A,B). Overall, *RNF43* was mutated in 79.7% of MSI-H colorectal tumors and 50.7% of MSI-H endometrial tumors (Supplementary Table 7.4). This association suggests that mismatch repair deficiency creates a permissive environment for the predominant *RNF43* frameshift alterations affecting codons 659 and 117.

Inactivating *RNF43* mutations should result in augmented Wnt signaling, owing to increased cell surface abundance of the Wnt receptor Frizzled^{2,3}. We thus asked whether any of the significantly mutated Wnt pathway genes displayed a mutually exclusive pattern of mutation with *RNF43* in colorectal or endometrial cancer. Indeed, truncating mutations in *APC*, the most frequently mutated gene in colorectal cancer, exhibited profound mutual exclusivity with truncating mutations in *RNF43* (6 of 74 *RNF43*-mutant samples had co-occurring truncating mutations in *RNF43* and *APC*, $P < 2.2 \times 10^{-16}$, Fisher's exact test; 4 of 55 mutant samples had equivalent co-occurring mutations when analysis was restricted to MSI-H tumors, $P = 3.6 \times 10^{-5}$; Figure 6.2A). Moreover, we detected *RSPO3-PTPRK* fusions (which activate Wnt signaling¹¹) in two tumors, neither of which harbored *RNF43* or *APC* mutations. The mutually exclusive relationship between mutations in *RNF43* and *APC* provides additional genetic evidence in support of an activating role for truncating mutations of *RNF43* in the Wnt pathway in

colorectal cancer.

4

Discussion and Summary

Our study demonstrates that *RNF43* is a frequently mutated gene in colorectal and endometrial cancers, enriched in the setting of MSI. These findings further predict that stomach cancer, a third tumor type characterized by Wnt pathway dysregulation and frequent MSI¹⁴, should also harbor frequent mutations in *RNF43*. Indeed, we found that 3 of 37 (8.1%) published stomach cancer exomes^{14,15} had

non-silent *RNF43* mutations, including a mutation encoding p.Gly659fs. Moreover, while this manuscript was under review, two studies reported *RNF43* mutations in gastric cancer^{16,17}. These studies also demonstrated a higher percentage of *RNF43* mutations in microsatellite-unstable samples in comparison to microsatellite-stable ones (54.6% versus 4.8% and 33% versus 3% in refs.¹⁶ and¹⁷, respectively). The discovery of multiple tumor types with frequent mutations in *RNF43* may have near-term clinical relevance in light of preclinical studies showing that *RNF43* mutations render Wnt-driven cancer cells susceptible to pharmacological porcupine inhibition³. More generally, these findings suggest that large-scale genomic analyses remain a fruitful means to discover biologically and clinically relevant changes in human cancer.

5

Methods

5.1 TUMOR SPECIMENS

We collected 185 colorectal tumor-normal pairs from the participants of NHS and HPFS who were diagnosed with colorectal carcinomas (Supplementary Table 7.5). All of the colorectal tumor and adjacent normal colorectal tissue samples were collected according to Partners Human Research Committee institutional re-

view board (IRB)-approved protocols, and informed consent was obtained from all subjects. Hematoxylin and eosin slides prepared from formalin-fixed paraffin-embedded blocks were examined for cancer extent, tumor grade and other tumor characteristics through pathology review. MSI status for each colorectal cancer sample was determined using standard protocols¹². We extracted genomic DNA from formalin-fixed, paraffin-embedded samples using the Qiagen QIAamp DNA FFPE Tissue kit. DNA quality was evaluated by quantification using the Quant-iT PicoGreen dsDNA Assay kit (Invitrogen) according to the manufacturer’s protocol.

5.2 DNA SEQUENCING

TCGA exome sequence data were downloaded from CGHub (see 5.5). The generation of sequencing data from the NHS and HPFS cohorts was performed using a protocol for formalin-fixed, paraffin-embedded samples that has been detailed previously¹⁸. Briefly, whole-exome capture libraries were constructed from tumor and normal DNA after sample shearing, end repair, phosphorylation and ligation to barcoded sequencing adaptors. DNA was then subjected to solution-phase hybrid capture with SureSelect v2 Exome bait (Agilent Technologies), and we performed multiplexing of the samples and sequencing on Illumina HiSeq 2000 instruments.

Average mean target coverage was $96\times$ (Supplementary Table 7.6). For validation, we used the Fluidigm Access Array microfluidic device. PCR products were barcoded, pooled and subjected to Illumina sequencing on a MiSeq instrument to a mean coverage of $11,000\times$. For six samples, validation of the c.1976delC (p.Gly659fs) mutation was performed by amplifying and Sanger Sequencing (Genewiz) a 211-bp product containing the change of interest (Supplementary Figure 7.1 and Supplementary Table 7.7).

5.3 EXOME SEQUENCING DATA ANALYSIS

Sequence analysis was performed as previously described^{18,19} to identify somatic single-nucleotide variants and short insertion or deletion events (Supplementary Table 7.8, Supplementary Table 7.9 and Supplementary Table 7.10). Somatic mutations were called with the MuTect algorithm¹⁹, and somatic indels were called with the Indelocator algorithm (see 5.5). Manual review of all *RNF43* indels in the NHS and HPFS cohorts was conducted using the Integrated Genomics Viewer²⁰, leading to deletions at base pair 56,435,160 being considered sequencing artifacts and excluded from further analysis. During manual review, it was discovered that several indels in *RNF43* resulting in p.Gly659fs were discarded as 'germline' indels by Indelocator owing to the presence of a small number of reads supporting the

variant in the normal exome. Indels likely to be somatic but originally filtered out as germline by Indelocator were rescued across the exome if they passed the following conservative criteria: coverage of at least 50 reads in both tumor and normal samples, >20% of reads supporting the variant in tumor samples and <5% of reads supporting the variant in normal samples.

5.4 *RSPO* FUSION DETERMINATION

TCGA RNA-seq data corresponding to the 222 samples in the analyzed colorectal cancer exome TCGA set were downloaded from CGHub (219 of 222 samples had available RNA-seq data). Gene fusions were called with TopHat-Fusion²¹.

5.5 URLs

CGHub, <https://cghub.ucsc.edu/>; Indelocator, <http://www.broadinstitute.org/cancer/cga/indelocator>.

5.6 ACCESSION CODES

All primary sequence files have been deposited in the database of Genotypes and Phenotypes (dbGAP) under accession phs000722.v1.p1.

6

6.1 FIGURES




Figure 6.1: *RNF43* mutations in colorectal and endometrial cancers. (a-c) Distribution and type of *RNF43* mutations in colorectal cancer, NHS, and HPFS set (a); colorectal cancer, TCGA set (b); and endometrial cancer, TCGA set (c). The domains of *RNF43* are depicted schematically at the bottom.




Figure 6.2: Association of *RNF43* mutations with MSI status and *APC* mutations. (a) Matrix displaying MSI status, *APC* truncating events and *RNF43* truncating events for each of 407 colorectal tumor samples (NHS and HPFS set and TCGA set). Each vertical line represents one colorectal cancer. Top, MSI status. MSI-H, microsatellite instability high; MSI-L, microsatellite instability low; MSI-NA, MSI status not available; MSS, microsatellite stable. Middle, a filled box shows that an *APC* truncating mutation is present in that tumor. Bottom, a filled box shows that an *RNF43* truncating mutation is present in that tumor. (b) Matrix displaying MSI status and *RNF43* truncation status for each of 248 endometrial tumor samples (TCGA set). Each vertical line represents one endometrial cancer. Top, MSI status. Bottom, a filled box shows that an *RNF43* truncating mutation is present in that tumor. Mutations in *APC* and *RNF43* are colored by type. In cases where multiple mutations were present in the same cancer sample, the presence of a frameshift mutation took priority in determining the displayed color, followed by the presence of a nonsense mutation.

7

Supplementary Figures and Tables

7.1 SUPPLEMENTARY FIGURES




Figure 7.1: Manual review of *RNF43* p.Gly659fs mutations using Integrated Genomics Viewer and validation by Sanger sequencing. Representative Integrated Genomics Viewer screenshots and Sanger sequencing chromatograms for *RNF43* p.Gly659fs-mutated tumor (a,c) and matched normal (b,d) tissue.

7.2 SUPPLEMENTARY TABLES

Tables 7.1-10 can be downloaded as an Excel file from
<https://www.nature.com/articles/ng.3127>.

Table 7.1: *RNF43* mutations in colorectal cancer, NHS/HPFS set ($n=185$).

Table 7.2: *RNF43* mutations in colorectal cancer, TCGA set ($n=222$).

Table 7.3: *RNF43* mutations in endometrial cancer, TCGA set ($n=248$).

Table 7.4: *RNF43* mutations and MSI status in colorectal and endometrial cancer samples.

Table 7.5: Clinico-pathological features of the NHS and HPFS patients and tumors.

Table 7.6: Exome coverage statistics for NHS and HPFS FFPE samples.

Table 7.7: PCR/Sanger primer sequences.

Table 7.8: Somatic coding mutations in colorectal cancer, NHS/HPFS set ($n=185$)

Table 7.9: Somatic coding mutations in colorectal cancer, TCGA set ($n=222$).

Table 7.10: Somatic coding mutations in endometrial cancer, TCGA set ($n=248$).

References

1. Giannakis M, Hodis E, Jasmine Mu X, Yamauchi M, Rosenbluh J, Cibulskis K, et al. RNF43 is frequently mutated in colorectal and endometrial cancers. *Nat Genet.* 2014 Dec;46(12):1264–6.
2. Koo BK, Spit M, Jordens I, Low TY, Stange DE, van de Wetering M, et al. Tumour suppressor RNF43 is a stem-cell E3 ligase that induces endocytosis of Wnt receptors. *Nature.* 2012 Aug;488(7413):665–9.
3. Jiang X, Hao HX, Growney JD, Woolfenden S, Bottiglio C, Ng N, et al. Inactivating mutations of RNF43 confer Wnt dependency in pancreatic ductal adenocarcinoma. *Proc Natl Acad Sci USA.* 2013 Jul;110(31):12649–54.
4. Wu J, Jiao Y, Molin M, Maitra A, de Wilde RF, Wood LD, et al. Whole-exome sequencing of neoplastic cysts of the pancreas reveals recurrent mutations in components of ubiquitin-dependent pathways. *Proc Natl Acad Sci USA.* 2011 Dec;108(52):21188–21193.
5. Jiao Y, Yonescu R, Offerhaus G, Klimstra D, Maitra A, Eshleman J, et al. Whole-exome sequencing of pancreatic neoplasms with acinar differentiation. *J Pathol.* 2014 Mar;232(4):428–35.
6. Hodis E, Watson I, Kryukov G, Arold S, Imielinski M, Theurillat J, et al. A landscape of driver mutations in melanoma. *Cell.* 2012 Jul;150(2):251–63.
7. TCGA. Comprehensive molecular portraits of human breast tumours. *Nature.* 2012 Oct;490(7418):61–70.
8. Stransky N, Egloff AM, Tward AD, Kostic AD, Cibulskis K, Sivachenko A, et al. The mutational landscape of head and neck squamous cell carcinoma. *Science.* 2011 Aug;333(6046):1157–60.

9. Clevers H, Nusse R. Wnt/ β -catenin signaling and disease. *Cell*. 2012 Jun;149(6):1192–205.
10. TCGA. Comprehensive molecular characterization of human colon and rectal cancer. *Nature*. 2012 Jul;487(7407):330–7.
11. Seshagiri S, Stawiski EW, Durinck S, Modrusan Z, Storm EE, Conboy CB, et al. Recurrent R-spondin fusions in colon cancer. *Nature*. 2012 Aug;488(7413):660–4.
12. Liao X, Lochhead P, Nishihara R, Morikawa T, Kuchiba A, Yamauchi M, et al. Aspirin Use, Tumor PIK3CA Mutation, and Colorectal-Cancer Survival. *N Engl J Med*. 2012 Oct;367(17):1596–606.
13. TCGA. Integrated genomic characterization of endometrial carcinoma. *Nature*. 2013 May;497(7447):67–73.
14. Wang K, Kan J, Yuen S, Shi S, Chu K, Law S, et al. Exome sequencing identifies frequent mutation of ARID1A in molecular subtypes of gastric cancer. *Nat Genet*. 2011 Oct;43(12):1219–23.
15. Zang Z, Cutcutache I, Poon S, Zhang S, McPherson J, Tao J, et al. Exome sequencing of gastric adenocarcinoma identifies recurrent somatic mutations in cell adhesion and chromatin remodeling genes. *Nat Genet*. 2012 May;44(5):570–4.
16. Wang K, Yuen S, Xu J, Lee S, Yan HH, Shi ST, et al. Whole-genome sequencing and comprehensive molecular profiling identify new driver mutations in gastric cancer. *Nat Genet*. 2014 Jun;46(6):573–582.
17. TCGA. Comprehensive molecular characterization of gastric adenocarcinoma. *Nature*. 2014 Sep;513(7517):202–9.
18. Van Allen E, Wagle N, Stojanov P, Perrin D, Cibulskis K, Marlow S, et al. Whole-exome sequencing and clinical interpretation of formalin-fixed,

- paraffin-embedded tumor samples to guide precision cancer medicine. *Nat Med*. 2014 Jun;20(6):682–8.
19. Cibulskis K, Lawrence M, Carter S, Sivachenko A, Jaffe D, Sougnez C, et al. Sensitive detection of somatic point mutations in impure and heterogeneous cancer samples. *Nat Biotechnol*. 2013 Mar;31(3):213–9.
 20. Robinson J, Thorvaldsdóttir H, Winckler W, Guttman M, Lander E, Getz G, et al. Integrative genomics viewer. *Nat Biotechnol*. 2011 Jan;29(1):24–6.
 21. Kim D, Salzberg S. TopHat-Fusion: an algorithm for discovery of novel fusion transcripts. *Genome Biol*. 2011 Aug;12(8):R72.