



(P0-176) Suicidal Ideation and Behavior During the COVID-19 Pandemic: A Retrospective, Single-Center Case Series

Citation

Sorg, Emily Emily, Mladen Mladen Nisavic, Scott Scott Beach, Rachel Rachel MacLean, Nicholas Nicholas Kontos, and Felicia Felicia Smith. 2022. "(P0-176) Suicidal Ideation and Behavior During the COVID-19 Pandemic: A Retrospective, Single-Center Case Series." *Journal of the Academy of Consultation-Liaison Psychiatry* 63: S81–S81.

Published Version

<https://doi.org/10.1016/j.jaclp.2022.03.168>

Permanent link

<https://nrs.harvard.edu/URN-3:HUL.INSTREPOS:37374410>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

Share Your Story

The Harvard community has made this article openly available. Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

Table 1: Demographic Characteristics of Patients Presenting with Suicidality Related to COVID-19 Pandemic

<i>Category</i>	<i>Demographics</i>	<i>Number (Percentage)</i>	
Patient Age	18-40 years	15 (65.2%)	
	41-60 years	6 (26%)	
	61 years and above	2 (8.8%)	
Gender ^a	Female	7 (30.4%)	
	Male	16 (69.6%)	
Housing Status	Housed	11 (47.8%)	
	Undomiciled	11 (47.8%)	
	Institutionalized (e.g. skilled nursing facility)	1 (4.4%)	
Suicidality	Suicidal ideation only	16 (69.6%)	
	Status post suicide attempt	7 (30.4%)	
Prior Psychiatric and/or Neuropsychiatric Diagnoses	No	6 (26%)	
	Yes	17 (74%)	
		Major Depressive Disorder	11
		Post-Traumatic Stress Disorder (PTSD)	5
	Bipolar Disorder	3	

		Schizophrenia and Schizoaffective Disorder	3
		Anxiety Spectrum Illness	2
		Attention Deficit Hyperactivity Disorder (ADHD)	2
		Antisocial Personality Disorder	1
		Borderline Personality Disorder	1
		Frontal Lobe Injury	1
Substance Use Disorder ^b	No		10 (43.5%)
	Yes		13 (56.5%)
		Alcohol	10
		Sedative/Hypnotic	2
		Cocaine	5
		Amphetamine	2
Prior Suicidality ^c	No		10 (43.5%)
	Yes		13 (56.5%)
COVID-19 status	Positive		8 (34.7%)
	Negative		15 (65.3%)

Footnotes:

a. All participants cisgender per information available in chart

b. All diagnostic information taken directly from chart; formal disorder titles per DSM-5 intentionally not used as unsure if symptoms fully met/meets DSM-5 criteria

c. Defined for these purposes as suicidal ideation or attempt(s)