



Religious service attendance and lower depression among women - a prospective cohort study

Citation

Li, Shanshan, Olivia I Okereke, Shun-Chiao Chang, Ichiro Kawachi, and Tyler J. VanderWeele. 2016. "Religious Service Attendance and Lower Depression Among Women—a Prospective Cohort Study." *Annals of Behavioral Medicine* 50 (6): 876–84. <https://doi.org/10.1007/s12160-016-9813-9>.

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:41275544>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Open Access Policy Articles, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#OAP>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)



Published in final edited form as:

Ann Behav Med. 2016 December ; 50(6): 876–884. doi:10.1007/s12160-016-9813-9.

Religious service attendance and lower depression among women - a prospective cohort study

Shanshan Li, ScD., Olivia I Okereke, MD., Shun-Chiao Chang, ScD., Ichiro Kawachi, Ph.D., and Tyler J. VanderWeele, Ph.D.

Departments of Nutrition (S.L.), Department of Epidemiology (O.I.O, T.J.V), Department of Biostatistics (T.J.V), Department of Social and Behavioral Sciences (I.K), Harvard T.H. Chan School of Public Health, 655 Huntington Avenue, Boston, MA 02115, United States; Channing Division of Network Medicine, Department of Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, 02115 (O.I.O, S-C.C.); Department of Psychiatry, Harvard Medical School and Brigham and Women's Hospital, Boston MA, 02115 (O.I.O)

Abstract

Objective—Previous studies on the association between religious service attendance and depression have been mostly cross-sectional, subject to reverse causation, and did not account for the potential feedback between religious service attendance and depression. We prospectively evaluated whether religious service attendance decreased risk of subsequent risk of depression and whether depression increased subsequent cessation of service attendance, while explicitly accounting for feedback with potential effects in both directions.

Method—We included a total of 48,984 U.S. nurses who were participants of the Nurses' Health Study with mean age 58 years and who were followed up from 1996 to 2008. Religious services attendance was self-reported in 1992, 1996, 2000, and 2004. Depression was defined as self-reported physician-diagnosed clinical depression, regular anti-depressant use, or severe depressive symptoms. Multivariate logistic regression and marginal structural models were used to estimate the odds ratio of developing incident depression, adjusted for baseline religious service attendance, baseline depression and time-varying covariates.

Results—Compared with women who never attended services, women who had most frequent and recent religious service attendance had the lowest risk of developing depression (odds ratio [OR] =0.71, 95% confidence interval [CI]:0.62–0.82). Compared with women who were not depressed, women with depression were less likely to subsequently attend religious services once or more per week (OR=0.74, 95% CI: 0.68–0.80).

Address reprint requests to: Dr. Tyler J VanderWeele, 677 Huntington Avenue, Kresge Building Boston, Massachusetts 02115, tvanderw@hsph.harvard.edu, Tel:617-432-7855, Fax: 617-432-1884.

Previous presentation:

/This work has been previously presented by Dr. VanderWeele at American Public Health Association Annual Meeting, New Orleans, LA, November 15–19, 2014 and the Annual Conference on Religion and Medicine, Cambridge, MA, March 6–8, 2015.

Location of work:

Department of Epidemiology, Harvard T.H. Chan School of Public Health, 655 Huntington Avenue, Boston, MA 02115

Disclosures: The authors have no conflicts of interest to declare. The authors report no competing interests.

Conclusions—In this study of U.S. women, there is evidence that higher frequency of religious service attendance decreased the risk of incident depression and women with depression were less likely to subsequently attend services.

Keywords

Religious service attendance; depression; causal models

INTRODUCTION

Depression is the leading cause of disability worldwide, affecting 350 million people of all ages (1). Approximately 6.7% of Americans are currently suffering from some form of depression (1). Depression is also a major risk factor for suicide (1). Compared with men, women are 70% more likely to experience depression over the course of a lifetime. The mechanisms underlying depression are poorly understood and the standard medications and treatments reduce depressive symptoms by only about 50% (2). Biological factors (e.g., genetic risk, medical illness) as well as psychological and/or spiritual factors (e.g., poor coping or resilience, low levels of mindfulness) may contribute to the development of depression (3). For religious patients, religious practice may constitute a complementary therapy as it can provide social support and may promote a sense of peace, hope, and meaning in life (4).

Previous studies have suggested a modest inverse association between religious service attendance and depression (5–9). However, the majority of these studies have had serious methodological limitations. Most research has been based on cross-sectional study designs (10–15), which generally cannot provide evidence of causality. Limited longitudinal research has been carried out (16–19). Complicating matters further, a recent study by Maselko *et al.* suggested that onset of major depression in early life (before 18 years of age) is associated with subsequent discontinuation of religious service attendance (20). Maselko *et al.* pointed out that if this is so, then cross-sectional studies could find a seemingly protective association between service attendance and depression even if there is no actual effect, simply because those who are depressed stop attending (20). VanderWeele noted that even longitudinal studies of service attendance and depression suggested a protective effect but no study had yet adequately addressed the methodological challenge of feedback with potential effects in both directions (21). Moreover, previous studies have only used one measurement of service attendance which captured “prevalent exposure” instead of “incident exposure”. Previous studies were thus still subject to reverse causation (21). To the best of our knowledge, no previous study has used repeated measurements of both depression and religious service attendance, and has been able to evaluate the potential feedback between religious service attendance and depression.

Using the Nurses’ Health Study (NHS) with repeated measures of service attendance and depression for up to 12 years of follow-up, we investigated the evidence for whether and to what extent: 1) religious service attendance is associated with the risk of subsequent depression; and 2) depression is associated with subsequent religious service attendance.

METHODS

Study design

The Nurses' Health Study (NHS) included 121,701 nurses aged 30–55 years at baseline, from across the United States, and who were followed up from 1976 to the present (22). Information on lifestyle and medical history was collected biennially using a self-administered questionnaire. Participants who had depression or religious service attendance information missing in 1992 or 1996 ($n=49,037$), or died before 2000 ($n=3,877$) were excluded. Compared with the general population, patients with cancer or cardiovascular disease have higher depression risk and might alter their religious service attendance. We, therefore, considered analyses both with and without participants who had a diagnosis of cardiovascular disease or cancer at baseline ($n=19,803$). In our primary analysis, we included a final sample size of 48,984 NHS participants with mean age of 58 years in 1996 (baseline) and followed up to 2008. The study protocol for the Nurses' Health Study was approved by the institutional review boards of Brigham and Women's Hospital and Harvard School of Public Health.

Assessment of religious service attendance

Religious service attendance in 1996 and 2000 and risk of depression in 2004

—Information on frequency of religious services attendance was self-reported in the 1992, 1996, 2000, 2004, and 2008 questionnaires, with the question “How often in the past year do you attend religious services?” Responses were “More than once per week, once per week, 1–3 times per month, less than once per month, never or almost never”. To ensure reasonably large categories, we modeled religious service attendance as four categories: “More than once per week, once per week, less than once per week, and never or almost never (reference group).”

Depression in 1996 and 2000 and subsequent religious service attendance in 2004

—We also examined associations between depression and subsequent religious service attendance. For the service attendance outcome in these analyses, we used the information reported in 2004 and 2008 questionnaires and modeled it as a binary variable (≥ 1 /week, <1 /week (reference group)).

Assessment of depression

Religious service attendance in 1996 and 2000 and risk of depression in 2004

—In the 2000, 2004, and 2008 questionnaires, study participants self-reported whether they had physician-diagnosed depression. Beginning in 1996 they reported biennially whether they had regular use of antidepressants. Depression measures in 2004 were taken as the primary outcomes. In 2004, participants were given the 10-item version of the Center for Epidemiologic Studies Depression scale (CESD-10). CESD-10 was designed for brief screening and identification of depressive symptoms in the general population (total scores range from 0 to 30), and a cutoff point of 10 is recommended for clinically significant depressive symptoms (23). As a secondary analysis, depression in 2008 was measured using the Geriatric Depression scale (GDS-15) with 15 items (24, 25); prior data indicate that a cutoff point of 10 yields a sensitivity of 100% and a specificity of 84% (24, 25); Cronbach

alpha coefficients were 0.75, with modest construct validity (25). Frequently recommended cutoff points for GDS-15 are currently 5 and 6 (24, 26). We performed the analysis using a cutoff point of 6 points, as unpublished data from an ongoing depression validation study suggested that 6 is the optimal cutoff point when compared to the gold-standard of DSM-IV (Diagnostic and Statistical Manual of Mental Disorders-IV) depression by semi-structured diagnostic interview.

Depression in 1996 and 2000 and subsequent religious service attendance in 2004—For the association between depression and subsequent religious service attendance, the depression exposure was defined as self-reported physician or clinician-diagnosed depression or Mental Health Index ≥ 5 , which was available in the Nurses' Health Study in 1996 and 2000 (yes or no).

Covariates

We considered a wide range of covariates in our analysis based on evidence from the literature and *prior* subject knowledge. We included socio-demographic, social, lifestyle factors, smoking status, alcohol consumption, post-menopausal hormone use, and health/medical conditions. Physical/functional limitations or disability was defined as having limitations activities requiring moderate exertion (e.g., climbing a flight of stairs) or having more than moderate limitations in demanding activities (e.g., lifting heavy objects). Covariate information was updated during the follow up and, for each exposure period, covariates were adjusted for from the questionnaire prior to the exposure assessment. For example, we adjusted covariates in 1992 for exposure in 1996, and covariates in 1996 for exposure in 2000.

Statistical methods

Religious service attendance in 1996 and 2000 and risk of depression in 2004

—Multivariate logistic regression and marginal structural models were used to calculate odds ratios between religious service attendance and risk of depression. The marginal structural models were our primary analysis and explicitly account for feedback between service attendance and depression, and allow for the estimation of the effects over time even in the presence of such feedback (27). They control for time-varying confounding by weighting techniques (27), described below, and were also used to assess the sensitivity of our conclusions to missing covariates and censoring. Specifically, we generated inverse probability weights for the service attendance exposure in 1996 and 2000, trimmed using a 5% rule, and applied stabilized weights to the marginal structural model (27). We also fit multivariate logistic regression models for comparison with other literature.

For associations between religious service attendance and risk of depression in 2004, information on service attendance in 1992 was adjusted as a baseline covariate, and service attendance in 1996 and in 2000 were modeled as categorical exposures with never attending as the reference group. To examine the robustness of the association between religious service attendance and depression, we used 3 definitions for the outcome of depression in 2004: 1) depression (yes or no) defined as self-reported physician or clinician-diagnosed depression, regular antidepressant medication use, or using CES-D-10 ≥ 10 points; 2)

depression (yes or no) defined using only the CES-D-10 cutoff of 10; 3) depressive symptoms (continuous) measured by CES-D-10 (Table 2). We adjusted for demographic, lifestyle, medical conditions, social support and physical function limitation covariates measured in 1992 for the 1996 exposure and measured in 1996 for exposure in 2000. We also adjusted for past religious service attendance frequency and depression status. The marginal structural models adjusted for feedback and time-varying confounding by weighting (27). We also evaluated the joint effects of service attendance in 1996 and 2000 in relation to depression risk. As secondary analyses, we further excluded participants who were depressed at baseline; we also examined the association among all participants including those with cardiovascular disease and cancer, and also among Catholics only, among Protestants only, and using depression risk in 2008.

To test the robustness of our results, we compared the following: a complete case approach, replacing missing data by a carry-forward from the previous questionnaire, and using an inverse probability weighting method with and without censoring weights to examine the influence of censoring during 1996–2000. We used sensitivity analysis for unmeasured confounding to assess how substantial residual unmeasured confounding would need to be to explain away the observed associations (28).

Depression in 1996 and 2000 and subsequent religious service attendance in 2004—Multivariate logistic regression and marginal structural models were used to estimate odds ratios between depression in 1996 and 2000 and subsequent religious service attendance. We performed subgroup analysis among participants with no baseline depression in 1992. Further, to assess robustness of results to censoring, we used several methods including a complete case approach; replacing missing data by a carry forward from previous questionnaire; and an inverse probability weighting method with and without censoring weights.

RESULTS

Religious service attendance and risk of depression

Among 48,984 women at 1996 baseline, 37,114 women reported religious service attendance and 9,244 of them attended more than once per week (Table 1). Participants who attended religious services more than once per week drank less, were less likely to be depressed, and were less likely to be a current smoker. In 1996, 6,973 (10.5%) women were depressed and 42,011 (89.5%) women were depression free (Supplementary Table 1). From 1996 to 2004, 6,432 women developed incident depression. Women who attended religious services often had higher mental health index scores (i.e., fewer depressive symptoms) and lower prevalence of self-reported physician or clinician-diagnosed depression (Table 1). In our study, 41% of women were Catholic and 53% were Protestant.

Using marginal structural models, compared with women who never or almost never attended religious services in 2000, adjusted odds ratios for depression in 2004 were 0.87 (95% CI: 0.79, 0.97) for women who attended less than once per week, 0.75 (95% CI: 0.67, 0.84) for women who attended weekly, and 0.71 (95% CI: 0.62, 0.82) for women who attended more than once/week (Table 2). There appeared to be a gradient in the estimates of

2000 religious service attendance levels with depression risk in 2004. Service attendance in 1996 did not appear to reduce depression risk beyond the associations with attendance in 2000. Results were similar with different analytic approaches or methods for handling censoring and missing data, or different definitions for depression; results were similar from both the logistic regression model and the marginal structural model (Data not shown). The results were consistent across 3 definitions for the outcome depression in 2004 (Table 2). The results did not substantially change when stratified by Catholic vs. Protestant (Supplementary Table 2). The results were also similar and significant, although slightly attenuated, using 2008 depression outcomes (Data not shown).

For an unmeasured confounder to fully explain away the odd ratio estimate of 0.71, the unmeasured confounder would have to both increase the likelihood of service attendance and decrease the likelihood of depression by 2.1-fold, above and beyond the measured confounders. This would constitute relatively substantial confounding; weaker confounding would not be sufficient to explain away the association. For an unmeasured confounder to bring the upper confidence limit of 0.82 for this estimate above 1, the unmeasured confounder would still have to both increase the likelihood of service attendance and decrease the likelihood of depression by 1.7-fold. Results were similar after further excluding participants with physician or clinician-diagnosed depression at baseline and only restricted to incident depression cases (Data not shown). Models that included an interaction between religious service attendance in 1996 and 2000 again indicated that more recent service attendance in 2000 was the stronger predictor, but a change in service attendance (either increase or decrease) was associated with somewhat higher risk of depression than continuous attendance (Table 3). Women who attended services frequently in both 1996 and 2000 had the lowest risk of subsequent depression; while women who had frequent attendance in 1996 but never or less than once per week in 2000 had elevated risk for depression (Table 3). Regardless of analytic methods used, more recent frequent religious service attendance was consistently associated with significantly lower risk of depression. The results were similar after including participants who had cardiovascular diseases and cancer and also when restricting the analysis to that population, with slightly larger estimates for that population (Supplementary Table 4). The effect size for service attendance is comparable to or larger than that of most, but not all, of many other risk factors for depression (Supplementary Table 3).

Depression and subsequent religious service attendance

Among 48,984 women at baseline, 6,973 had Mental Health Index ≥ 52 , which was the cutoff point used to define presence of clinical depression. Compared with women who were not depressed, women with depression had lower religious service attendance in 1992 and 1996 (Supplementary Table 1). They were more likely to feel stressed either at work or home, less likely to be currently married, and less likely to be unemployed in the past two years (Supplementary Table 1).

Using marginal structural models, compared to women without depression, women who were depressed in 2000 were less likely to attend frequent religious services (once per week or more) in 2004, with adjusted OR=0.74, (95% CI: 0.68, 0.80) (Table 4). Results were

similar across religious service attendance outcomes in 2004 and 2008; results were similar in logistic regression models, but the magnitude of the odds ratios was smaller when accounting for missing data and censoring (Data not shown). For depression there was also some evidence for an effect of depression in 1996 lowering the likelihood of subsequent service attendance (OR=0.91; 95% CI: 0.83, 1.00) beyond the effects of depression in 2000.

DISCUSSION

In this prospective cohort study, with repeated measurements of religious service attendance and depression, women with more frequent service attendance had lower risk of incident depression. Women with depression were subsequently less likely to attend religious services compared with those who were not depressed. Results were robust across different years of services attendance, different analytic methods, accounting for feedback, missing data and censoring. Our results provide evidence for potential feedback in both directions with effect sizes of roughly equal magnitude. Our conclusions are consistent with previous results, and moreover provide evidence of effects in both directions.

Religiosity is multidimensional, and different aspects of religion and spirituality may therefore be associated with depression differently. In our study, we only have one domain measure of religiosity or spirituality, namely, service attendance. Our study did not measure spirituality and other dimensions of a person's religiosity. Further, frequency of religious service attendance is also a form of social participation and social engagement, and closely related to social support. In our study, we attempted to capture religious service attendance in everyday life in our primary analysis by excluding women with chronic diseases or comorbidities at baseline since these conditions may alter patterns of religious service attendances. However, results were similar also when they were included.

Measurement error is also inevitable in self-reported religious service attendance. However, previous studies suggest that service attendance is often over-reported (29), which might lead to an underestimation of effects. Over-reporting might also still preserve relative ordering. Depression was measured using different validated questionnaires for 1996, 2000, 2004, and 2008. In our study, prevalence of depression may be misclassified due to the potential of under-detection/under-treatment for depression and under-diagnosis of depression. However, regardless of the different measures for depression, our results were robust and consistent. To separate out new cases with persistent or unremitted depression, we further excluded depression cases at baseline and only evaluated incident depression. Although we adjusted for major confounders for the association between religious service attendance and depression, our study is an observational study and may still be subject to unmeasured confounders and residual confounding. However, for an unmeasured confounder to fully explain away the association of service attendance with subsequent depression it would have to both increase the likelihood of service attendance and decrease the likelihood of depression by 2.1-fold, above and beyond the measured covariates, which may not be likely. In our study, we did not inquire about potential mediating mechanisms. All covariates we adjusted for in our analyses were measured in the period prior to the exposure and thus would not be mediators. Future studies are needed to investigate potential underlying mediating mechanisms between religious service attendance and depression, and

also to better assess the effects of other measures of religiosity than frequency of religious service attendance.

Religiosity and spirituality vary greatly in style and content across different cultures and race/ethnicity groups. The strengths of the association between religiosity and health may differ across gender, race/ethnicity, and different forms of religiosity. Our study mainly consisted of white Christians, and consisted entirely of U.S. women who were nurses. In this study, about 90% reported being Protestant or Catholic. The health-related behaviors of nurses may differ from the general population. Our results might not be generalizable to other race/ethnicity group and demographic groups with other religious beliefs. Individuals suffering from depression tend to withdraw from all types of social activities and patterns may differ with other chronic diseases. However, the extensive confounding adjustment and robustness of our results to different analytic approaches help ensure the internal validity of our study.

Our study findings do not imply that physicians should prescribe religious service recommendations for patients. The ethical questions are of course complex. However for patients who are already religious, service attendance might be encouraged as a form of meaningful social participation to alleviate depression. Somewhat paradoxically, however, persons with depression are more likely to cease attending. Further study of this phenomenon merits attention. The results may also be of interest to religious communities in that those who become depressed tend to cease attendance, which may exacerbate depression further. Further support could be extended to such persons before they leave. It may moreover be important for psychiatrists, physicians, and other health care providers to be aware of and take into account the potential role of faith and belief in depression prevention and treatment.

The strengths of our study included the large sample size, repeated measures on service attendance and depression, and analytic methods accounting for feedback. To the best of our knowledge, no previous study had examined incident rather than prevalent religious service attendance, the latter of which could still be subject to reverse causation, as depression onset during adulthood may lead to a lower level of religious service attendance in later life. By using marginal structural models, we have been able to tease out the feedback between religious service attendance and depression, and appropriately account for reverse causation and time dependent confounding. We also have clear temporality of covariates, exposure, and outcome. Using validated questionnaires, we adjusted for lifestyle, medical history, and a wide range of potential confounders, along with baseline exposure, baseline outcome, and time-varying covariates.

In conclusion, based on a large longitudinal cohort study with repeated measures of religious service attendance and depression, we found evidence that women with a higher level of religious service attendance were less likely to develop depression, and women who were depressed were less likely to attend religious services frequently.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Funding: This work was support by the Templeton Foundation and National Institutes of Health grant R01 ES017876. The funding agencies played no role in the design and conduct of this study; collection, management, analysis, and interpretation of the data; and preparation, review, or approval of the manuscript.

SL and TJV contributed to the study design, data analysis, interpretation, and manuscript preparations. OIO, S-C C, and IK contributed to the study design and critical review of the manuscript. SL and TJV had full access to all the data in the study and take responsibility for the integrity of the data, and the accuracy of the data analysis.

References

1. Brundtland GH. From the World Health Organization. Mental health: new understanding, new hope. *JAMA*. 2001; 286(19):2391. [PubMed: 11712923]
2. Blackburn–Munro G. Hypothalamo–pituitary–adrenal axis dysfunction as a contributory factor to chronic pain and depression. *Curr Pain Headache Rep*. 2004; 8(2):116–24. [PubMed: 14980146]
3. Koenig HG. Religion, spirituality, and health: a review and update. *Adv Mind Body Med*. 2015; 29(3):19–26. [PubMed: 26026153]
4. Mytko JJ, Knight SJ. Body, mind and spirit: towards the integration of religiosity and spirituality in cancer quality of life research. *Psycho-oncology*. 1999; 8(5):439–50. [PubMed: 10559803]
5. Koenig HG. Depression in chronic illness: does religion help? *Journal of Christian nursing: a quarterly publication of Nurses Christian Fellowship*. 2014; 31(1):40–6. [PubMed: 24592603]
6. McCullough ME, Larson DB. Religion and depression: a review of the literature. *Twin Res*. 1999; 2(2):126–36. [PubMed: 10480747]
7. Hayward RD, Owen AD, Koenig HG, Steffens DC, Payne ME. Longitudinal relationships of religion with posttreatment depression severity in older psychiatric patients: evidence of direct and indirect effects. *Depression res and Treat*. 2012:745970.
8. Koenig HG. Religion and depression in older medical inpatients. *Am J Geriatr Psychiatry*. 2007; 15(4):282–91. [PubMed: 17384313]
9. Koenig HG, Hays JC, George LK, Blazer DG, Larson DB, Landerman LR. Modeling the cross–sectional relationships between religion, physical health, social support, and depressive symptoms. *Am J Geriatr Psychiatry*. 1997; 5(2):131–44. [PubMed: 9106377]
10. Rasic D, Kisely S, Langille DB. Protective associations of importance of religion and frequency of service attendance with depression risk, suicidal behaviours and substance use in adolescents in Nova Scotia, Canada. *J Affect Disord*. 2011; 132(3):389–95. [PubMed: 21458077]
11. King DA, Lyness JM, Duberstein PR, He H, Tu XM, Seaburn DB. Religious involvement and depressive symptoms in primary care elders. *Psychol Med*. 2007; 37(12):1807–15. [PubMed: 17498321]
12. Chatters LM, Bullard KM, Taylor RJ, Woodward AT, Neighbors HW, Jackson JS. Religious participation and DSM–IV disorders among older African Americans: findings from the National Survey of American Life. *Am J Geriatr Psychiatry*. 2008; 16(12):957–65. [PubMed: 19038894]
13. Agorastos A, Demiralay C, Huber CG. Influence of religious aspects and personal beliefs on psychological behavior: focus on anxiety disorders. *Psychol Res Behav Manag*. 2014; 7:93–101. [PubMed: 24648780]
14. Maselko J, Gilman SE, Buka S. Religious service attendance and spiritual well–being are differentially associated with risk of major depression. *Psychol Med*. 2009; 39(6):1009–17. [PubMed: 18834554]
15. Koenig, HG.; McCullough, ME.; Larson, DB. *Handbook of Religion and Health*. Oxford ; New York: Oxford University Press; 2001.

16. Koenig HG, George LK, Peterson BL. Religiosity and remission of depression in medically ill older patients. *The American journal of psychiatry*. 1998; 155(4):536–42. [PubMed: 9546001]
17. Barton YA, Miller L, Wickramaratne P, Gameroff MJ, Weissman MM. Religious attendance and social adjustment as protective against depression: a 10-year prospective study. *J Affect Disord*. 2013; 146(1):53–7. [PubMed: 22959684]
18. Barton YA, Miller L, Wickramaratne P, Gameroff MJ, Weissman MM. Religiosity and major depression in adults at high risk: a ten year prospective study. *The American journal of psychiatry*. 2012; 169(1):89–94. [PubMed: 21865527]
19. Rasic D, Asbridge M, Kisely S, Langille D. Longitudinal associations of importance of religion and frequency of service attendance with depression risk among adolescents in Nova Scotia. *Can J Psychiatry*. 2013; 58(5):291–9. [PubMed: 23756289]
20. Maselko J, Hayward RD, Hanlon A, Buka S, Meador K. Religious service attendance and major depression: a case of reverse causality? *Am J Epidemiol*. 2012; 175(6):576–83. [PubMed: 22350581]
21. Vanderweele TJ. Re: "Religious service attendance and major depression: a case of reverse causality?". *Am J Epidemiol*. 2013; 177(3):275–6. [PubMed: 23292958]
22. Belanger CF, Hennekens CH, Rosner B, Speizer FE. The nurses' health study. *The American journal of nursing*. 1978; 78(6):1039–40. [PubMed: 248266]
23. Andresen EM, Malmgren JA, Carter WB, Patrick DL. Screening for depression in well older adults: evaluation of a short form of the CES–D (Center for Epidemiologic Studies Depression Scale). *American journal of preventive medicine*. 1994; 10(2):77–84. [PubMed: 8037935]
24. Lyness JM, Noel TK, Cox C, King DA, Conwell Y, Caine ED. Screening for depression in elderly primary care patients. A comparison of the Center for Epidemiologic Studies–Depression Scale and the Geriatric Depression Scale. *Archives of internal medicine*. 1997; 157(4):449–54. [PubMed: 9046897]
25. Friedman B, Heisel MJ, Delavan RL. Psychometric properties of the 15-item geriatric depression scale in functionally impaired, cognitively intact, community-dwelling elderly primary care patients. *Journal of the American Geriatrics Society*. 2005; 53(9):1570–6. [PubMed: 16137289]
26. Blank K, Gruman C, Robison JT. Case –Finding for Depression in Elderly People: Balancing Ease of Administration With Validity in Varied Treatment Settings. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*. 2004; 59(4):M378–M84.
27. Robins JM, Hernan MA, Brumback B. Marginal structural models and causal inference in epidemiology. *Epidemiology (Cambridge, Mass)*. 2000; 11(5):550–560.
28. Vanderweele TJ, Arah OA. Bias formulas for sensitivity analysis of unmeasured confounding for general outcomes, treatments, and confounders. *Epidemiology (Cambridge, Mass)*. 2011; 22(1): 42–52.
29. Hadaway CK, Marler PL, Chaves M. What the polls don't show: a closer look at U.S. Church attendance. *American Sociological Review*. 1993; 58:741–52.

Table 1

Baseline Characteristics of 48,984 Nurses' Health Study Participants by Frequencies of Religious Services Attending in 1996

	Frequency of religious service attendance			
	Never or almost never (n=11,870)	Less than once/wk. (n=7,899)	Once/wk. (n=19,974)	More than once/wk. (n=9,244)
Age, years	61.5(7.0)	61.5(7.1)	61.5(7.0)	61.5(7.0)
Race/ethnicity, %				
Caucasian	97	97	97	98
African American	1	1	1	1
Other	2	2	2	1
Currently married, %	73	74	78	77
Nurses' education level, %				
High school	68	67	71	70
Bachelor	21	22	19	21
Master or above	11	11	9	8
Religious group, %				
Catholic	29	28	53	40
Protestant	62	63	43	53
Live alone, %	15	15	13	14
Not Employed in Last 2 Years, %	37	34	37	40
Number of close friends, continuous	3.2(1.0)	3.4(1.0)	3.4(1.0)	3.5(1.1)
Have someone close to talk to (yes or no), %	92	94	94	95
Self-reported clinician-diagnosed depression in 1996, binary, % ^b	13	11	10	9
Mental Health Index-5 score in 1996, continuous ^a	78.4(14.7)	78.9(13.8)	79.9(13.2)	81.5(12.1)
Physical activity, 10 MET-hrs. /wk.	2.04(2.4)	2.0(2.4)	1.9(2.2)	1.9(2.2)
Weight change since age 18, kg	13.2(26.1)	13.3(25.9)	13.3(25.3)	13.6(25.4)
Alternative Healthy Eating Index 2010, continuous ^c	47.9(10.5)	47.9(10.3)	46.9(10.1)	47.7(10.1)
Alcohol intake (g/day), %				
0	59	58	55	43
< 5	28	32	31	26
5.1– 15	19	18	17	12
15.4–30	6	5	4	3
>30	6	3	3	2
Smoking status, %				
Past smoker	45	44	41	35
Current smoker	20	15	11	6
Never smoker	65	59	52	41
Body Mass Index (BMI) , %				
<18 kg/m ²	17	16	17	17
18.1–21 kg/m ²	32	34	32	32
21.1–23 kg/m ²	22	24	24	24

	Frequency of religious service attendance			
	Never or almost never (n=11,870)	Less than once/wk. (n=7,899)	Once/wk. (n=19,974)	More than once/wk. (n=9,244)
23.1–25 kg/m ²	20	19	19	19
25.1–27 kg/m ²	4	4	4	5
27.1–30 kg/m ²	3	2	3	2
30+ kg/m ²	2	1	1	1
Hypertension, %	27	27	26	26
High cholesterol, %	41	42	42	42
Diabetes, %	3	3	3	3
Have physical/functional limitations, %	48	48	46	47

^a Measured by Mental Health Index. Higher score represent less depressive symptoms. Depression was defined as MHI \geq 52.

^b Self-reported depression diagnosis or anti-depressant use

^c Higher Alternative Healthy Eating Index 2010 (AHEI2010) represents better diet quality

Multivariate Associations Between Religious Services Attendance in 1996 and 2000 with Subsequent Onset of Depression in 2004

Table 2

Marginal Structural Model						
Depression ^a	Presence of severe depressive symptoms ^b	Continuous level of depressive symptoms ^c	Estimate	Std Err	P value	
Odds ratios	95% CI	Odds ratios	95% CI	Estimate	Std Err	P value
Religious service attendance in 1996						
Never	Reference	Reference	Reference	Reference	Reference	Reference
< 1/wk.	0.89, 1.12	1.04	0.91, 1.19	0.03	0.09	0.70
1/wk.	0.95, 1.24	1.12	0.95, 1.30	0.11	0.10	0.27
> 1/wk.	0.98, 1.36	1.18	0.98, 1.43	0.12	0.12	0.29
Religious service attendance in 2000						
Never	Reference	Reference	Reference	Reference	Reference	Reference
< 1/wk.	0.79, 0.97	0.87	0.77, 0.98	-0.20	0.08	0.01
1/wk.	0.67, 0.84	0.72	0.63, 0.82	-0.48	0.09	<.0001
> 1/wk.	0.62, 0.82	0.65	0.55, 0.76	-0.53	0.10	<.0001

CI: confidence interval

Multivariate model adjusted for age (years), baseline depressive symptom in 1992(continuous), baseline religious service attendance in 1992 (never, less than once/week, more than once/week), live alone (yes, no), not employed in the past 2 years (yes, no), marital status (currently married, yes, no), nurses' education level (high school, bachelor, master), husband education level (under high school, high school graduate, college or above), postmenopausal hormone use (never, current, past), race/ethnic groups(Caucasian, African American, Asian, Hispanic, other), geographic region (middle, north, south), census tract level median family income (dollars per year, continuous), height (continuous), family history of Myocardial Infarction(yes, no), family history of cancer(yes, no), family history of diabetes(yes, no), body mass index (BMI,<18, 18-1-21, 21.1-23, 23.1-25, 25.1-27, 27.1-30, >30+ kg/m2), weight change since age 18 (kg), physical activity (MET hrs/wk), hypertension(yes, no), hypercholesterolemia (yes, no), diabetes(yes, no), diet quality (Alternative Healthy Eating Index 2010, continuous), alcohol consumption (0, <5, 5.1-15, 15.4-30, >30 g/day), number of close friends (continuous), have someone close to talk to(yes, no), smoking status (never, current, past), physical/functional limitations (yes, no)

Marginal structural model with weights accounted for censoring and missing data

^aBinary variable of depression :Defined as participants who either self-reported a physician or clinician-diagnosed depression, or use of antidepressant medications, or CESD-10 = 10

^bBinary CESD-10 binary (= 10): Measured by CESD-10. Higher score represent more depressive symptoms. Total score ranges from 0 to 30. Depression was defined as CESD-10 = 10.

^cCESD-10 continuous: Measured by CESD-10. Higher score represent more depressive symptoms. Total score ranges from 0 to 30. Depression was defined as CESD-10 = 10.

Transitions in Religious Services Attendance in 1996 and 2000 with Subsequent Onset of Depression in 2004

Table 3

Marginal Structural Model									
Religious service attendance	Depression ^a		Presence of severe depressive symptoms ^b		Continuous level of depressive symptoms ^c		Std Err	P value	Reference
	Odds ratios	95% CI	Odds ratios	95% CI	Estimate	Reference			
< 1/wk. in 1996 and never in 2000	Reference		Reference		Reference	Reference	Reference	Reference	Reference
< 1/wk. in 1996 and <1/wk. in 2000	0.87	0.78, 0.96	0.86	0.76, 0.98	-0.21	0.08	0.01	0.01	0.01
< 1/wk. in 1996 and 1/wk. in 2000	0.90	0.77, 1.05	0.80	0.67, 0.97	-0.31	0.12	0.01	0.01	0.01
< 1/wk. in 1996 and >1/wk. in 2000	0.99	0.70, 1.39	1.18	0.81, 1.73	0.14	0.32	0.65	0.65	0.65
1/wk. in 1996 and never in 2000	1.33	1.09, 1.62	1.24	0.98, 1.57	0.30	0.18	0.09	0.09	0.09
1/wk. in 1996 and <1/wk. in 2000	1.05	0.91, 1.22	1.05	0.88, 1.24	0.02	0.11	0.88	0.88	0.88
1/wk. in 1996 and 1/wk. in 2000	0.82	0.72, 0.92	0.79	0.69, 0.91	-0.38	0.09	<.0001	<.0001	<.0001
1/wk. in 1996 and >1/wk. in 2000	0.80	0.70, 0.92	0.71	0.61, 0.84	-0.44	0.10	<.0001	<.0001	<.0001

1 Measured by CESD-10. Higher score represent more depressive symptoms. Total score ranges from 0 to 30. Depression was defined as CESD-10 ≥ 10.

Multivariate model adjusted for age (years), baseline depressive symptom in 1992(continuous), baseline religious service attendance in 1992 (never, less than once/week, once/week, more than once/week), live alone (yes, no), not employed in the past 2 years (yes, no), marital status (currently married, yes, no), nurses' education level (high school, bachelor, master), husband education level (under high school, high school graduate, college or above), postmenopausal hormone use (never, current, past), race/ethnic groups(Caucasian, African American, Asian, Hispanic, other), geographic region (middle, north, south), census tract level median family income (dollars per year, continuous), height (continuous), family history of Myocardial Infarction(yes, no), family history of cancer(yes, no), family history of diabetes(yes, no), body mass index (BMI,<18, 18.1–21, 21.1–23, 23.1–25, 25.1–27, 27.1–30, >30+ kg/m2), weight change since age 18 (kg), physical activity (MET hrs./wk.), hypertension(yes, no), hypercholesterolemia (yes, no), diabetes(yes, no), diet quality (Alternative Healthy Eating Index 2010, continuous), alcohol consumption (0, <5, 5.1–15, 15.4–30, >30 g/day), number of close friends (continuous), have someone close to talk to(yes, no), smoking status (never, current, past), physical/functional limitations (yes, no).

Marginal structural model with weights accounted for censoring and missing data.

^a Binary variable of depression :Defined as participants who either self-reported a physician or clinician-diagnosed depression, or use of antidepressant medications, or CESD-10 ≥ 10

^b Binary CESD-10 binary (10). Measured by CESD-10. Higher score represent more depressive symptoms. Total score ranges from 0 to 30. Depression was defined as CESD-10 ≥ 10.

^c CESD-10 continuous. Measured by CESD-10. Higher score represent more depressive symptoms. Total score ranges from 0 to 30. Depression was defined as CESD-10 ≥ 10.

Multivariate Associations Between Depression in 1996 and 2000 and Subsequent Frequent Religious Services Attendance (once/week) in 2004 and 2008

Table 4

Follow up period	Religious service attendance (1/week) in 2004		Religious service attendance (1/week) in 2008		
	1996-2004	Odds ratio	95% CI	Odds ratio	95% CI
Depression in 1996	No	Reference		Reference	
	Yes	0.91	0.83, 1.00	0.91	0.83, 1.00
Depression in 2000	No	Reference		Reference	
	Yes	0.74	0.68, 0.80	0.75	0.69, 0.82

Multivariate model adjusted for age (years), baseline depressive symptom in 1992(continuous), baseline religious service attendance in 1992 (never, less than once/week, once/week, more than once/week), live alone (yes, no), not employed in the past 2 years (yes, no), marital status (currently married, yes, no), nurses' education level (high school, bachelor, master), husband education level (under high school, high school graduate, college or above), postmenopausal hormone use (never, current, past), race/ethnic groups(Caucasian, African American, Asian, Hispanic, other), geographic region (middle, north, south), census tract level median family income (dollars per year, continuous), height (continuous), family history of Myocardial Infarction(yes, no), family history of cancer(yes, no), family history of diabetes(yes, no), body mass index (BMI,<18, 18.1-21, 21.1-23, 23.1-25, 25.1-27, 27.1-30, >30+ kg/m²), weight change since age 18 (kg), physical activity (MET hrs./wk.), hypertension(yes, no), hypercholesterolemia (yes, no), diabetes(yes, no), diet quality (Alternative Healthy Eating Index 2010, continuous), alcohol consumption (0, <5, 5.1-15, 15.4-30, >30 g/day), number of close friends (continuous), have someone close to talk to(yes, no), smoking status (never, current, past), physical/functional limitations (yes, no).

Marginal structural model with weights accounted for censoring and missing data.