



Deconstruction and gauge theories in AdS5

The Harvard community has made this article openly available. [Please share](#) how this access benefits you. Your story matters

Citation	Randall, Lisa, Yael Shadmi, and Neal Weiner. 2003. "Deconstruction and Gauge Theories in AdS5." Journal of High Energy Physics 2003 (1): 055–055. https://doi.org/10.1088/1126-6708/2003/01/055 .
Citable link	http://nrs.harvard.edu/urn-3:HUL.InstRepos:41417349
Terms of Use	This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA

Deconstruction and gauge theories in AdS_5

To cite this article: Lisa Randall *et al* JHEP01(2003)055

View the [article online](#) for updates and enhancements.

Related content

- [Anomalies and Fayet-Iliopoulos terms on warped orbifolds and large hierarchies](#)
Takayuki Hirayama and Koichi Yoshioka
- [Running of Gauge Couplings in \$AdS_5\$ via Deconstruction](#)
Adam Falkowski and Hyung Do Kim
- [Bulk gauge fields in warped space and localized supersymmetry breaking](#)
Zackaria Chacko and Eduardo Pontón

Recent citations

- [Exponential Hierarchies from Anderson Localization in Theory Space](#)
Nathaniel Craig and Dave Sutherland
- [Disassembling the clockwork mechanism](#)
Nathaniel Craig *et al*
- [Confronting Higgs couplings from \$U\(1\)^4\$ D-term extensions and natural SUSY at the LHC and ILC](#)
Moritz McGarrie *et al*

Deconstruction and gauge theories in AdS_5

Lisa Randall

*Jefferson Physical Laboratory, Harvard University
Cambridge, MA 02138, USA
E-mail: randall@physics.harvard.edu*

Yael Shadmi*

*Physics Department, Technion
Haifa 32000, Israel
E-mail: yshadmi@physics.technion.ac.il*

Neal Weiner

*Department of Physics, University of Washington
Seattle, WA, 98195, USA
E-mail: nealw@phys.washington.edu*

ABSTRACT: On a slice of AdS_5 , despite having a dimensionful coupling, gauge theories can exhibit logarithmic dependence on scale. In this paper, we utilize deconstruction to analyze the scaling behavior of the theory, both above and below the AdS curvature scale, and shed light on position-dependent regularizations of the theory. We comment on applications to geometries other than AdS.

KEYWORDS: Renormalization Group, Field Theories in Higher Dimensions, GUT.

*Incumbent of a Technion Management Career Development Chair.

Contents

1. Introduction	1
2. Gauge theories in AdS	2
3. The classical deconstructed theory	3
3.1 Couplings at intermediate energies in the deconstructed theory	5
3.2 Couplings at high energies in the deconstructed theory	6
4. Gauge coupling running in deconstructed RS	6
4.1 Fine latticizations and power law running	9
5. Position dependent regularizations from deconstruction	10
6. Conclusions	11

1. Introduction

In recent years, it has become clear that extra dimensions can play an important role in our understanding of the hierarchy problem [1, 2]. In [2], it was noted that the cutoff scale of the standard model could be $O(\text{TeV})$, and yet have weakly coupled gravity if the standard model fields were localized on a brane on the boundary of a slice of AdS_5 .

One seeming consequence of having a cutoff at a TeV is that it is impossible to discuss issues of the UV, such as high scale unification. However, this apparent obstacle no longer applies when the gauge sector of the standard model is placed in the bulk of AdS. In particular, in [3] the couplings of fields on the UV brane were studied, and it was demonstrated explicitly that despite the dimensionful bulk gauge coupling, the effective coupling of the zero mode runs logarithmically below the AdS curvature scale. In [4, 5], a position-dependent regulator motivated by holography was used to show that the predictions of perturbative high-scale unification were possible, even in theories where standard model fields experience a strong-coupling cutoff at the TeV scale. More recently, [6]–[12] have studied the behavior of gauge fields in AdS. All of these works agree that, at some level, gauge running in AdS is logarithmic, rather than power law, below the compactification scale. However, there nonetheless remains some confusion as to what constitutes a sensible regulator, and in what sense one can talk about a zero mode definitively above the IR brane scale.

Over the past year, “deconstruction” [13, 14] has appeared as a new tool for model building and understanding the properties of higher-dimensional gauge theories. (For early related work on the subject, see [15].) By viewing the higher-dimensional theory as a product of four-dimensional theories, one can easily study the properties of the theory at

intermediate energies. In this paper, we will employ deconstruction to give us an understanding of the properties of AdS gauge theories at energies above the scale of the IR brane.¹ We, too, will recover log running above a TeV, but we will also see how position dependent regulators naturally arise in the deconstructed framework. We also will recover cutoff sensitive “power-law running” effects above the curvature scale, which will appear as threshold effects associated with the cutoff of the theory. We will also see in what sense we can discuss a renormalized zero mode above the IR brane scale. Lastly, the results we achieve can be straightforwardly applied to different geometries.

2. Gauge theories in AdS

Five dimensional gauge field theories are defined as cutoff theories. Because the gauge coupling is dimensionful, there is an associated ultra-violet strong coupling scale. However, AdS gauge theories present a number of subtleties not found in flat extra dimensions.

In flat extra dimensions, we see the first Kaluza-Klein mode at a scale $1/R$. At a scattering energy $E \simeq n/R$, one exchanges n KK modes, giving a strong coupling scale $\Lambda \sim 4\pi/Rg_4^2 \sim 4\pi^2/g_5^2$. Note that we could live on a brane at any point in the space and this formula would remain essentially unchanged.

In AdS, the situation is quite different. If we consider the RS1 space with UV and IR cutoff branes, the mass of the first KK mode is set by the IR scale as is the spacing between KK modes. Thus, a scattering at a high scale $\Lambda \gg M_{IR}$ would naively be strongly coupled. The essential difference in AdS is that this is a position dependent question. In the language of KK modes, this is because the light KK modes are strongly coupled to the IR brane, but very weakly coupled to the UV brane.

This result is not surprising because it arises from the basic symmetries of AdS. Let us consider a U(1) gauge theory in AdS. We can write the metric as

$$ds^2 = e^{-2\sigma(y)} dx^2 - dy^2. \tag{2.1}$$

For now and unless otherwise noted, we take $\sigma(y) = ky$. With this metric, the gauge boson action is

$$S = -\frac{1}{4} \int d^4x dy \frac{1}{g_5^2} [G_{\mu\nu}^2 - 2e^{-2\sigma} G_{\mu 5}^2], \tag{2.2}$$

where all 4d contractions are with $\eta_{\mu\nu}$. Note that this action is invariant under the transformation

$$\begin{aligned} y &\rightarrow y + \delta, \\ x &\rightarrow e^{k\delta} x, \\ \partial_\mu &\rightarrow e^{-k\delta} \partial_\mu, \\ A_\mu &\rightarrow e^{-k\delta} A_\mu. \end{aligned} \tag{2.3}$$

The transformation of the gauge field is necessary to insure that gauge covariant derivatives of charged fields are also covariant under the rescaling.

¹Deconstructing gauge theories in RS was also discussed in [16, 17].

This symmetry tells us that a process with four-momentum p_μ at a point $y = y_0$ relates to an equivalent process with four-momentum $e^{-k\delta}p_\mu$ at a point $y' = y_0 + \delta$. Or (not surprisingly) the local (in y) strong coupling scale scales with the warp factor.

In going to the quantum theory, we have to be careful in how we define our couplings. In four-dimensional QED, for instance, we define the coupling e at a scale μ_0 , with the same e and the same μ_0 everywhere. This is a requirement for preserving the Poincaré invariance of the theory. In AdS we do not have five-dimensional Poincaré invariance so we must be careful to define our theory to preserve the symmetries.

One way we *cannot* define the theory is to specify all couplings at the scale of the UV brane. This is because over most of the space the gauge theory becomes strongly coupled below this scale, and hence this is not even a region in which the perturbative field theory is defined.

Likewise, we could specify the couplings at the scale of the IR brane, but this is troubling as we take the IR brane to infinity, and, moreover, it would spoil the symmetries of the theory. Hence, the most natural definition, the one that maintains the symmetry of the space, is to assume that the couplings are specified at a scale which scales with the warp factor in y . This will be essential in writing down the deconstructed quantum theory.

3. The classical deconstructed theory

Let us begin by reviewing the standard process of deconstruction. Our goal will be to write a five-dimensional $SU(n)$ as a four-dimensional, $SU(n)^N$ gauge theory with bifundamental fields, Q_i (see table 1) with²

$$\langle Q_i \rangle = v_i \text{diag}(1, 1, \dots, 1). \tag{3.1}$$

The lattice will more accurately reproduce intermediate scale physics with increasing N , the number of lattice sites. Most importantly, the theory is cut off at a local scale $\Lambda_j \sim v_j$, so that we can regulate our theory in a gauge-invariant fashion.

From the outset, deconstruction of an AdS gauge theory is quite different from the deconstruction of one in flat space. In flat space, if a fermion is localized at any point in the additional dimension(s), it will couple with comparable strength to each of the KK modes. Consequently, accurately reproducing the intermediate energy features in the deconstructed theory requires matching states and energies.

The AdS case is quite different. Above the local AdS curvature scale, the gauge theory quickly becomes strongly coupled. However, those KK modes are very weakly coupled to the UV brane, so in a deconstructed theory, these modes will be automatically grouped, depending on the coarseness of the latticization, into single modes whose effects will replace multiple modes in the extra dimensional theory. While this seems peculiar, it is precisely because those modes are so weakly coupled that this remains a true description of the features of the higher dimensional theory.

²We will comment on the supersymmetric version of the theory at end of this section.

	SU(n) ₁	SU(n) ₂	SU(n) ₃	⋯	SU(n) _N
Q ₁	□	□̄	1	⋯	1
Q ₂	1	□	□̄	⋯	1
	⋮	⋮	⋮	⋱	⋮
Q _{N-1}	1	1	1	⋯	□̄

Table 1: Field content for a deconstructed SU(n) theory.

We begin by discretizing the 5th dimension in (2.2). We have

$$S = -\frac{1}{4} \int d^4x \left[\sum_j \frac{a_j}{g_{5j}^2} G_{\mu\nu}^{(j)2} - 2 \sum_j \frac{a_j e^{-2\sigma_j}}{g_{5j}^2} G_{\mu 5}^{(j)2} \right], \tag{3.2}$$

where j denotes the lattice site, a_j is the lattice spacing near site j , $a_j = y_{j+1} - y_j$ and $j = 1, \dots, N$, and

$$G_{\mu 5}^{(j)} = \partial_\mu A_5^{(j)} - \frac{1}{a_j} \left(A_\mu^{(j+1)} - A_\mu^{(j)} \right). \tag{3.3}$$

To write down a lattice action we can replace A_5 by the link fields

$$U = e^{i \int_{y_j}^{y_{j+1}} dy A_5(x,y)}, \tag{3.4}$$

so that for small lattice spacing,

$$D_\mu U \sim i \left(a_j \partial_\mu A_5^j - A_\mu^j + A_\mu^{j+1} \right). \tag{3.5}$$

Let us now consider the action of the deconstructed theory,

$$S = \int d^4x \left[-\frac{1}{4} \sum_j \frac{1}{g_j^2} F_{\mu\nu}^{(j)2} + \sum_j \frac{1}{g_j^2} D_\mu Q_j^\dagger D^\mu Q_j \right]. \tag{3.6}$$

The peculiar normalization for the matter fields is convenient since they will ultimately supply the 5th component of the gauge fields. In the supersymmetric version of the theory, which we will later consider, this normalization allows for a holomorphic definition of the length of the 5th dimension, since the gauge coupling does not appear in the relation between this length and the VEVs.

Assuming all Q 's get VEVs we have $N - 1$ σ -models, coupled by the gauge interaction. Writing $Q_j = v_j \tilde{U}_j$, we choose the parameters of the deconstructed theory, v_j and g_j , so that the action (3.6) agrees with the 5d lattice action (3.2):

$$\begin{aligned} \frac{1}{g_j^2} &= \frac{a_j}{g_{5j}^2}, \\ \frac{v_j^2}{g_j^2} &= \frac{e^{-2\sigma_j}}{a_j g_{5j}^2}, \end{aligned} \tag{3.7}$$

so that,

$$v_j^2 = \frac{e^{-2\sigma_j}}{a_j^2}. \tag{3.8}$$

	SU(n) ₁	SU(n) ₂	SU(n) ₃	⋯	SU(n) _{N}
$\bar{q}_{i=1\dots n}$	\square	1	1	⋯	1
$q_{a=1\dots n}$	1	1	1	⋯	\square

Table 2: Adding vectorlike fields at the end sites cancels anomalies in the supersymmetric theory.

Here we see the effect of the AdS geometry. While a theory with five-dimensional Lorentz invariance naturally has a deconstructed description with all VEVs, v_j , equal, here the v_j 's scale with the warp factor. We also note that, with conventional normalization for the bifundamentals, the gauge coupling would appear in (3.8).

We thus have a “dictionary” between the $5d$ parameters — the gauge coupling, g_{5j} , and the lattice spacing a_j — and the parameters of the deconstructed theory, v_j and g_j .

One can consider different variations of this basic theory. For example, in order to study bulk matter, we can add matter fields for every SU(n) factor. In the supersymmetric version of the theory (with $\mathcal{N} = 1$ supersymmetry), one needs to add matter in order to cancel anomalies on the two ends of the chain. A simple possibility is to add fundamentals and antifundamentals as shown in table 2.

These fields can get mass through non-renormalizable operators [18], or through extra vectorlike matter [19]. However, they do not affect our basic analysis and results.

3.1 Couplings at intermediate energies in the deconstructed theory

To get a sense of the behavior of the deconstructed theory, let us begin by considering the exchange of a bulk gauge field coupling to a field on the UV brane at energies $R^{-1} \gg q \gg \text{TeV}$. Being confined to the UV brane is equivalent to being charged under the gauge group G_1 in the deconstructed theory.

At very low energies, the entire theory is higgsed to the diagonal. Then the two-point function in the Feynman-t’Hooft gauge for exchange of a bulk gauge boson between two fields on the UV brane is given simply by³

$$G_q(0)\eta^{\mu\nu} \simeq -i \frac{g^2}{N} \frac{\eta^{\mu\nu}}{q^2}. \tag{3.9}$$

At a higher energy $q \gg \text{TeV}$, we have resolved the independent gauge groups higgsed down below a scale q , and instead, the dominant contribution comes from the exchange of the diagonal gauge field of the groups which have been higgsed down at energies above q . This behavior is shown in figure 1. The effective number of groups contained in the diagonal at a scale q can be written approximately as

$$aN_{\text{eff}} \simeq R \log \left(\frac{q}{\Lambda} \right). \tag{3.10}$$

Now we can approximate the exchange of the gauge field by the diagonal mode, yielding

$$G_q(0)\eta^{\mu\nu} \approx -i \frac{g^2}{N_{\text{eff}}} \frac{\eta^{\mu\nu}}{q^2} \approx -i \frac{g_5^2 k}{\log(q/\Lambda)q^2} \eta^{\mu\nu}, \tag{3.11}$$

where we have used the relation $g_5^2 = g^2 a$.

³We use normalization such that the gauge coupling is included in the propagator for clarity.




Figure 1: Connection between RG flow in the continuum and deconstructed theory. A high energy theory has the massive modes (localized near the UV brane) integrated out to leave a different theory with a different UV brane with large kinetic terms for the bulk gauge field. For momentum q processes occurring on the UV brane, the dominant contribution to exchange comes from the diagonal mode of the shaded region, yielding an effective IR cutoff in the continuum theory.

It is notable that the dominant classical logarithmic scaling for the AdS gauge field two-point function falls out automatically from the deconstructed theory, without invoking any further assumptions about the holographic nature of the theory. This is an excellent indication that the deconstructed theory is effectively reproducing the physics of the five-dimensional theory.

3.2 Couplings at high energies in the deconstructed theory

At high energies, our coarse latticization is insufficient. If we want to understand physics near the strong coupling scale of the theory, we should latticize such that each $\alpha_j = g_j^2/4\pi \approx 4\pi$. In this case, adjacent VEVs are comparable, and in order to analyze the problem, one would need to explicitly diagonalize the gauge boson mass matrix, and to determine the couplings to the different modes. Clearly however, groups that are higgsed at scales much lower than the UV cutoff would be irrelevant. Groups higgsed at scales close to the cutoff would lead to running as in the presence of a flat extra dimension, with radius $R_{\text{eff}} \sim 1/k$. In the next section, where we will derive from holomorphy considerations the general form of the low-energy gauge coupling, we will see that these expectations are indeed borne out.

4. Gauge coupling running in deconstructed RS

Before we can talk about running of the gauge coupling, we must define what we mean, precisely. We cannot speak perturbatively about fields on the TeV brane scattering at energies well above a TeV. In what sense can we discuss running?

First, let us note that while physics on the TeV-brane is not perturbative above a TeV, physics at different positions, sufficiently close to the Planck brane, is perturbative. In particular, scattering on the Planck brane is perturbative at all energies up to the cutoff. At low energies, scattering proceeds through exchange of the gauge-field zero-mode.

This is true for fields living on the TeV brane, as well as for fields living on the Planck brane. In fact, because of the unbroken four-dimensional gauge invariance, the zero-mode couplings on both branes are equal. Thus, we can relate Green's functions on the TeV brane to Green's functions on the Planck brane at low energies. The latter can of course be extrapolated to high energies. Thus, we can relate measured gauge couplings on the TeV brane to those that *would* exist for fields on the UV brane at high energies.

The deconstructed theory is characterized by two sets of parameters: the dimensionless couplings g_j , and the dimensionful VEVs v_j . As discussed above, in order to mimic the continuum theory, we choose

$$v_j = v e^{-jw}, \tag{4.1}$$

with $w = ka$. In addition, the individual couplings are defined at the scale of the corresponding VEVs, that is,

$$g_j = g_j(v_j) = g, \tag{4.2}$$

or, in terms of the strong coupling scales of the individual groups,

$$\Lambda_j^b = v_j^b e^{-8\pi^2/g^2}, \tag{4.3}$$

where b is the one-loop beta function coefficient, which is the same for all groups. Examining the relations (3.7), we see that this indeed corresponds to taking $g_{5j} = \text{const}$ and $a_j = a$.⁴

It is instructive to start by considering a coarse lattice, with the lattice spacing larger than the AdS curvature: $a \gg k^{-1}$. Then we have

$$v_1 \gg v_2 \gg \dots \gg v_{N-1}. \tag{4.4}$$

This simplifies the analysis considerably: we can study the running by turning on the VEVs one at a time. It is amusing that the RS case turns out to be simpler to analyze than the flat case, where all VEVs are equal and therefore have to be considered together.

Let us now derive the coupling of the low energy theory. At the scale v_1 , $SU(n)_1 \times SU(n)_2$ is broken to the diagonal subgroup, which we label $SU(n)_{2D}$ (with the diagonal group of j groups labeled $SU(n)_{jD}$), with gauge coupling,

$$\frac{1}{g_{2D}^2(v_1)} = \frac{1}{g_1^2(v_1)} + \frac{1}{g_2^2(v_1)}. \tag{4.5}$$

We can now run down to the scale v_2 , where,

$$\frac{1}{g_{2D}^2(v_2)} = \frac{1}{g_1^2(v_1)} + \frac{1}{g_2^2(v_1)} + \frac{b}{8\pi^2} \ln \frac{v_2}{v_1}, \tag{4.6}$$

where we used the fact that the diagonal subgroups have the same beta function coefficients as the original group factors.

⁴Note that the same holds with conventional normalization for the matter fields: Taking the VEVs to warp and defining the individual couplings to equal a common value at *exponentially varying* scales corresponds to a 5d coupling and lattice spacing that have uniform values at *exponentially varying* scales.

Repeating this j times we find,

$$\frac{1}{g_{jD}^2(v_j)} = \frac{1}{g_1^2(v_1)} + \dots + \frac{1}{g_j^2(v_j)} + \frac{b}{8\pi^2} \ln \frac{v_j}{v_1}. \quad (4.7)$$

Using the fact that $k = -w^{-1} \ln v_j/v$, and $g_5^2 = ag^2$, this can be written as

$$\frac{1}{g_D^2(v_j)} = -\frac{1}{kg_5^2} \ln \frac{v_j}{v} + \frac{b}{8\pi^2} \ln \frac{\mu}{v} - \frac{b}{8\pi^2} ka. \quad (4.8)$$

Indeed, the low energy coupling depends logarithmically on the scale. In fact, we see that the low energy coupling contains two pieces with logarithmic energy dependence. The first term on the RHS of eqn. (4.8) is a classical piece, which reproduces the fact that the 4d and 5d couplings are related by the “radius”. In the deconstructed theory, it is obtained by summing over KK modes at tree level. The second piece is a loop effect. It is obtained from running the coupling between adjacent KK modes.

Let us now consider the supersymmetric theory. In the presence of $\mathcal{N} = 1$ supersymmetry, holomorphy and symmetries dictate the dependence of the diagonal $SU(n)$ strong coupling scale, Λ_D , on Λ_j , v_j , at any stage of the Higgsing. In particular, at low energies,

$$\Lambda_D^b \sim \frac{\Lambda_1^b \dots \Lambda_N^b}{N^b v_1^b \dots v_{N-1}^b}. \quad (4.9)$$

The factor of N^b is not dictated by symmetries, but can be found from explicit computation [20].⁵

The relation (4.9) allows us to obtain the low energy coupling for any profile of the VEVs. In particular, for the RS geometry, with (4.1), (4.3), we have at low energy

$$\frac{1}{g_L^2(\mu)} = \frac{N}{g^2} + \frac{b}{8\pi^2} \ln \frac{\mu}{v/N}. \quad (4.10)$$

Running from a scale below k , deconstruction yields ordinary logarithmic running with a beta function set by the four-dimensional gauge theory. In terms of the parameters of the theory, this gives us

$$\frac{1}{g_4^2(\mu)} \simeq -\frac{1}{kg_5^2} \ln \frac{v_{N-1}}{v} + \frac{b}{8\pi^2} \ln \frac{\mu}{v/N} = \frac{R}{g_5^2} + \frac{b}{8\pi^2} \ln \frac{\mu}{v/N}. \quad (4.11)$$

This indeed reproduces the expression we found for the non-supersymmetric theory. Again, we see the logarithmic dependence on the energy scale in both the tree-level and one-loop pieces.

⁵The factor N^b can also be motivated by the following argument: Consider the flat space theory. To properly reproduce the continuum theory, the scale of the unbroken diagonal subgroup should remain fixed as we take $N \rightarrow \infty$. Since v scales as N (recall v is the inverse lattice spacing), and g scales as \sqrt{N} (so that the classical diagonal coupling is fixed), the expression (4.9) should contain the factor N^b .

4.1 Fine latticizations and power law running

So far we have seen that a coarse latticization of AdS can generate the log running behavior which is characteristic of these gauge theories. But what about energies above k ? At these energies the space is approximately flat and there is a possible power law contribution between k and Λ .⁶

This arises from a subtlety which we have to this point ignored. We have defined our couplings g_j at the scale v_j . This was reasonable because the scales varied with energy, preserving the symmetries of the theory. Also, v_j is approximately the local cutoff, so if we believe the unification is only manifest in a microscopic theory (e.g. string theory), this is the approximately the cutoff scale (which is ambiguous).

These ambiguities make little difference when the coupling scales logarithmically with the cutoff, but is very important in the power law piece. To extract the behavior most straightforwardly, we will consider the supersymmetric theory, but the presence of the power law piece will not depend on the supersymmetric matter content. Consider again (4.9). If we state that the couplings unify at a scale $M_j \sim v_j$ we have

$$\begin{aligned} \Lambda_D^b &\sim \frac{\Lambda_1^b \dots \Lambda_N^b}{N^b v_1^b \dots v_{N-1}^b}, \\ \mu e^{8\pi^2/g_D^2(\mu)} &\sim \frac{\Lambda_1^b \dots \Lambda_N^b}{N^b M_1^b \dots M_{N-1}^b} \times \frac{M_1^b \dots M_{N-1}^b}{v_1^b \dots v_{N-1}^b}, \\ &\sim v^b e^{8\pi^2 N/g^2} e^{\gamma N b}, \end{aligned} \tag{4.12}$$

where $\gamma = \log(M_j/v_j N^{1/N})$ is independent of j .

Restating this in terms of gauge couplings, we have

$$\frac{1}{g_D^2(\mu)} = \frac{N}{g^2} + \frac{b}{8\pi^2} \log\left(\frac{v}{\mu}\right) + \frac{Nb\gamma}{8\pi^2}. \tag{4.13}$$

Note that we have made no assumptions about the background geometry to derive this result — it can be applied to any space. In AdS we recognize the first term as the usual $R/g_5^2 = \log(v/v_{N-1})/kg_5^2$ piece, and the second we recognize as the log running piece. The final piece is the power law running piece. Using the relation

$$N = \frac{R}{a} = \frac{v \log(v/v_{N-1})}{k}, \tag{4.14}$$

we can rewrite (4.13) as

$$\frac{1}{g_D^2(\mu)} = \frac{N}{g^2} + \frac{b}{8\pi^2} \log\left(\frac{v}{\mu}\right) + \frac{vb\gamma}{8\pi^2 k} \log\left(\frac{v}{v_{N-1}}\right). \tag{4.15}$$

Taking $v \sim \Lambda$, we see the linear cutoff dependence (“power law running”) explicitly. This term is worthy of two observations. The first is that it seems that merely by restating each g_j at v_j we could remove the power dependence. It is true that a suitable redefinition of g_5^2

⁶We are indebted to Nima Arkani-Hamed for very useful discussions on these issues.

can absorb this piece entirely. However, since we are ultimately considering the possibility of unification, we cannot do this as the separate redefinitions of couplings for $3 - 2 - 1$ would not be $SU(5)$ symmetric. Hence, in considering *differences* of couplings, this is a real effect.

The second observation is that there is essentially a free parameter in the theory, γ . This is not surprising because the theory has a linear divergence, implying cutoff sensitivity. We have employed a particular cutoff which has an ambiguity: namely the unification scale relative to the cutoff scale. This uncertainty is reasonable from an effective field theory perspective as it merely parameterizes our ignorance of the physics which generates the low energy field theory. In terms of the continuum theory, the upshot is that while there is true power law dependence on the cutoff, the coefficient of this term is unknown. That being said, it should also be noted that the relative strengths of these cutoff terms are still proportional to the relevant β functions.

Lastly, we note that here the power law piece can be subdominant to the log running piece. We would expect additional threshold effects comparable to the power law piece in any complete model, which could easily spoil quantitative unification. In AdS, because the log piece can dominate for unification scales below the strong coupling cutoff of the theory, the GUT-violating threshold effects are under control and one can still speak reliably about unification.

5. Position dependent regularizations from deconstruction

In [5], a position dependent cutoff was used to study the evolution of gauge couplings in a warped background. The regularization employed was an energy dependent IR brane, which was motivated by holography and the presence of strong coupling away from the UV brane at intermediate energy. Here, this regularization arises naturally from the deconstructed theory, without appealing to holography.

Let us begin by running the theory down from high energies. In the deconstructed theory, we gradually Higgs down more and more groups such that the highest energy group becomes more weakly coupled. At each level we integrate out the classical and quantum fluctuations of only the left-most groups in the quiver diagram. If we treat deconstruction as an invertible map, the resulting position space regularization is illustrated in figure 1.

In the deconstructed picture the appropriate regularization is obvious. One merely integrates out the heavy modes, which exist near the UV brane, and replaces the theory with a different one, with a new UV brane positioned nearer to the IR, but now with large boundary kinetic terms corresponding to the modes integrated out.

Of course, this relates the fundamental lagrangian with a lower energy effective theory. We also want to ask questions about the relation between low-energy and high-energy scatterings on the UV brane. Here, our interpretation of the above diagram is somewhat different. At any given momentum q , we are cut off from the sites higgsed at scales less than q . As a consequence, we should calculate loops only with the modes from the sites at energies above q . In particular, as we have already argued, the dominant contribution to the two-point function comes from the diagonal mode of the sites at energies above q .

Mapping the deconstructed theory back to the continuum, we see that we have introduced an IR cutoff on the theory, and the diagonal mode of the sites above q maps to the zero mode of the theory with this IR brane in place. Thus, while one cannot safely talk about couplings of the zero mode of the complete theory at high energies (because of strong couplings in the IR) one can reasonably talk about the zero mode of the theory with a different IR cutoff. It is notable that the deconstructed picture forces this regularization on us, and we need not appeal to holographic intuition.

Lastly, we note that these regularizations should also appear automatically in other, non-trivial geometries. In situations without an obvious regularization, the deconstructed approach, with VEVs following the warp factor, should give the appropriate position dependent regularization.

6. Conclusions

Extra dimensional theories with TeV-scale cutoffs invite careful analysis of questions regarding the ultra-violet. In the AdS slice of RS1, the introduction of gauge fields into the bulk offers the possibility of discussing short distance physics, such as coupling constant unification.

The application of deconstruction to this question obviates many of the questions raised in continuum theory analyses. Position dependent regulators appear automatically in the deconstructed framework, motivating their use in studying continuum theories.

Note added: while this work was being prepared, [21] appeared which addresses similar issues. The authors would also like to acknowledge unpublished work on this subject by N. Arkani-Hamed and A. Cohen.

Acknowledgments

The authors thank Kaustubh Agashe, Antonio Delgado, Walter Goldberger, Wittek Skiba, Veronica Sanz, Erich Poppitz and especially Yuri Shirman and Nima Arkani-Hamed for useful discussions. Y.S. and N.W. would like to thank the visitor program at Harvard University where this work was initiated, and the Aspen Center for Physics where much of this work was completed. The research of Y.S. was supported in part by the Technion V.P.R. Fund, the Henri Gutwirth Promotion of Research Fund, and by a Technion Fund for the Promotion of Research.

References

- [1] N. Arkani-Hamed, S. Dimopoulos and G.R. Dvali, *The hierarchy problem and new dimensions at a millimeter*, *Phys. Lett. B* **429** (1998) 263 [[hep-ph/9803315](#)].
- [2] L. Randall and R. Sundrum, *A large mass hierarchy from a small extra dimension*, *Phys. Rev. Lett.* **83** (1999) 3370 [[hep-ph/9905221](#)].
- [3] A. Pomarol, *Grand unified theories without the desert*, *Phys. Rev. Lett.* **85** (2000) 4004 [[hep-ph/0005293](#)].

- [4] L. Randall and M.D. Schwartz, *Unification and the hierarchy from AdS₅*, *Phys. Rev. Lett.* **88** (2002) 081801 [[hep-th/0108115](#)].
- [5] L. Randall and M.D. Schwartz, *Quantum field theory and unification in AdS₅*, *J. High Energy Phys.* **11** (2001) 003 [[hep-th/0108114](#)].
- [6] K.-W. Choi, H.D. Kim and Y.W. Kim, *Gauge coupling renormalization in orbifold field theories*, *J. High Energy Phys.* **11** (2002) 033 [[hep-ph/0202257](#)].
- [7] W.D. Goldberger and I.Z. Rothstein, *High energy field theory in truncated AdS backgrounds*, *Phys. Rev. Lett.* **89** (2002) 131601 [[hep-th/0204160](#)].
- [8] K. Agashe, A. Delgado and R. Sundrum, *Gauge coupling renormalization in RS1*, *Nucl. Phys.* **B 643** (2002) 172 [[hep-ph/0206099](#)].
- [9] K.-W. Choi, H.D. Kim and I.-W. Kim, *Radius dependent gauge unification in AdS₅*, [hep-ph/0207013](#).
- [10] R. Contino, P. Creminelli and E. Trincherini, *Holographic evolution of gauge couplings*, *J. High Energy Phys.* **10** (2002) 029 [[hep-th/0208002](#)].
- [11] W.D. Goldberger and I.Z. Rothstein, *Effective field theory and unification in AdS backgrounds*, [hep-th/0208060](#).
- [12] K.-W. Choi and I.-W. Kim, *One loop gauge couplings in AdS₅*, [hep-th/0208071](#).
- [13] N. Arkani-Hamed, A.G. Cohen and H. Georgi, *(De)constructing dimensions*, *Phys. Rev. Lett.* **86** (2001) 4757 [[hep-th/0104005](#)].
- [14] C.T. Hill, S. Pokorski and J. Wang, *Gauge invariant effective lagrangian for Kaluza-Klein modes*, *Phys. Rev.* **D 64** (2001) 105005 [[hep-th/0104035](#)].
- [15] M.B. Halpern and W. Siegel, *Electromagnetism as a strong interaction*, *Phys. Rev.* **D 11** (1975) 2967.
- [16] H.-C. Cheng, C.T. Hill and J. Wang, *Dynamical electroweak breaking and latticized extra dimensions*, *Phys. Rev.* **D 64** (2001) 095003 [[hep-ph/0105323](#)].
- [17] H. Abe, T. Kobayashi, N. Maru and K. Yoshioka, *Field localization in warped gauge theories*, [hep-ph/0205344](#).
- [18] C. Csáki, J. Erlich, C. Grojean and G.D. Kribs, *4d constructions of supersymmetric extra dimensions and gaugino mediation*, *Phys. Rev.* **D 65** (2002) 015003 [[hep-ph/0106044](#)].
- [19] W. Skiba and D. Smith, *Localized fermions and anomaly inflow via deconstruction*, *Phys. Rev.* **D 65** (2002) 095002 [[hep-ph/0201056](#)].
- [20] C. Csáki, J. Erlich, V. V. Khoze, E. Poppitz, Y. Shadmi and Y. Shirman, *Exact results in 5d from instantons and deconstruction*, *Phys. Rev.* **D 65** (2002) 085033 [[hep-th/0110188](#)].
- [21] A. Falkowski and H.D. Kim, *Running of gauge couplings in AdS₅ via deconstruction*, *J. High Energy Phys.* **08** (2002) 052 [[hep-ph/0208058](#)].