



DIGITAL ACCESS TO SCHOLARSHIP AT HARVARD

Association between Neighborhood Safety and Overweight Status among Urban Adolescents

The Harvard community has made this article openly available.
[Please share](#) how this access benefits you. Your story matters.

Citation	Duncan, Dustin T., Renee M. Johnson, Beth E. Molnar, and Deborah Azrael. 2009. Association between neighborhood safety and overweight status among urban adolescents. BMC Public Health 9:289.
Published Version	doi:10.1186/1471-2458-9-289
Accessed	August 19, 2018 3:02:43 PM EDT
Citable Link	http://nrs.harvard.edu/urn-3:HUL.InstRepos:4569480
Terms of Use	This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA

(Article begins on next page)

Research article

Open Access

Association between neighborhood safety and overweight status among urban adolescents

Dustin T Duncan*^{1,2}, Renee M Johnson^{2,3,4}, Beth E Molnar^{1,2} and Deborah Azrael^{2,3}

Address: ¹Department of Society, Human Development and Health, Harvard School of Public Health, Boston, MA, USA, ²Harvard Youth Violence Prevention Center, Harvard School of Public Health, Boston, MA, USA, ³Department of Health Policy and Management, Harvard School of Public Health, Boston, MA, USA and ⁴Department of Community Health Sciences, Boston University School of Public Health, Boston, MA, USA

Email: Dustin T Duncan* - dduncan@hsph.harvard.edu; Renee M Johnson - rjohnson@bu.edu; Beth E Molnar - bmolnar@hsph.harvard.edu; Deborah Azrael - azrael@hsph.harvard.edu

* Corresponding author

Published: 11 August 2009

Received: 2 March 2009

BMC Public Health 2009, 9:289 doi:10.1186/1471-2458-9-289

Accepted: 11 August 2009

This article is available from: <http://www.biomedcentral.com/1471-2458/9/289>

© 2009 Duncan et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Neighborhood safety may be an important social environmental determinant of overweight. We examined the relationship between perceived neighborhood safety and overweight status, and assessed the validity of reported neighborhood safety among a representative community sample of urban adolescents (who were racially and ethnically diverse).

Methods: Data come from the 2006 Boston Youth Survey, a cross-sectional study in which public high school students in Boston, MA completed a pencil-and-paper survey. The study used a two-stage, stratified sampling design whereby schools and then 9th–12th grade classrooms within schools were selected (the analytic sample included 1,140 students). Students reported their perceptions of neighborhood safety and several associated dimensions. With self-reported height and weight data, we computed body mass index (BMI, kg/m²) for the adolescents based on CDC growth charts. Chi-square statistics and corresponding *p*-values were computed to compare perceived neighborhood safety by the several associated dimensions. Prevalence ratios (PRs) and 95% confidence intervals (CI) were calculated to examine the association between perceived neighborhood safety and the prevalence of overweight status controlling for relevant covariates and school site.

Results: More than one-third (35.6%) of students said they always felt safe in their neighborhood, 43.9% said they sometimes felt safe, 11.6% rarely felt safe, and 8.9% never felt safe. Those students who reported that they rarely or never feel safe in their neighborhoods were more likely than those who said they always or sometimes feel safe to believe that gang violence was a serious problem in their neighborhood or school (68.0% vs. 44.1%, *p* < 0.001), and to have seen someone in their neighborhood assaulted with a weapon (other than a firearm) in the past 12 months (17.8% vs. 11.3%, *p* = 0.025). In the fully adjusted model (including grade and school) stratified by race/ethnicity, we found a statistically significant association between feeling unsafe in one's own neighborhood and overweight status among those in the Other race/ethnicity group [(PR = 1.56, (95% CI: 1.02, 2.40)].

Conclusion: Data suggest that perception of neighborhood safety may be associated with overweight status among urban adolescents in certain racial/ethnic groups. Policies and programs to address neighborhood safety may also be preventive for adolescent overweight.

Background

Adolescent overweight is a major public health priority because of the immediate and long-term health risks it poses and because of its rapidly increasing prevalence. Specifically, adolescent overweight is associated with numerous deleterious chronic health consequences in adolescence, such as hypertension [1-3], type 2 diabetes [1,3], and asthma [1,3-5], and in adulthood, such as hypertension [1,3,6], type 2 diabetes [1,3,7], coronary heart disease [1,7,8], and certain cancers [1,7-9]. Trend data indicate that overweight among adolescents has been dramatically increasing [10-12]. The prevalence of being overweight among 12–19 year olds, for example, has more than doubled in recent decades; going from 6.1% in 1971–1974 to 15.5% in 1999–2000 [11].

In an effort to thwart the overweight epidemic, research attention has increasingly been placed on social and environmental factors that give rise to overweight [10,13,14]. Neighborhood safety may be an important contributor to overweight, as it has been theorized that fear of violence and crime in the immediate social environment is a barrier to physical activity and a facilitator of sedentary behavior (two well-established predictors of overweight). Several studies among adolescents have shown an inverse association between neighborhood safety and physical activity [15-28], and others have demonstrated a positive association between neighborhood safety and sedentary behavior [17,18]. The few studies that have examined neighborhood safety as a predictor of overweight among adolescents have yielded inconsistent findings, with some finding a statistically significant inverse association [29,30] and others not having found such evidence [18,19,31,32]. These inconsistencies may be due to differences across study populations, in the measurement of neighborhood safety, and/or in the categorization of overweight status.

There are limitations with and gaps in the extant literature worth noting. First, most cross-sectional studies that examine neighborhood safety and overweight status among adolescents have used odds ratios for the studied associations. However, prevalence ratios are more appropriate because they provide a more valid parameter estimate when the prevalence of the outcome is common while the odds ratio is likely to overestimate the effect when there is a relatively high disease prevalence [33-36]. Given the high prevalence of adolescent overweight [10,11,37,38], prevalence ratios probably should be estimated when examining correlates of adolescent overweight. Second, few studies of neighborhood safety and adolescent overweight have been conducted among racially and ethnically diverse samples [19,31,32]. Generalizability of the previous research findings is therefore limited. Third, none of the existing studies in this area

have sought to evaluate the accuracy of the measure of perceived neighborhood safety. Threats to validity (e.g. misclassification of neighborhood safety) would likely result in an underestimation of effects due to non-differential misclassification bias.

The primary aim of the present study was to examine the association between neighborhood safety and overweight status among a representative sample of public high school students in Boston, MA. This study improves on earlier studies by using prevalence ratios as the parameter estimate, with a racially and ethnically diverse sample and by assessing the validity of perceived neighborhood safety (our secondary objective). To achieve the latter, we examined associations between perceived neighborhood safety and associated dimensions, including beliefs about gang activity and witnessed violence. Our evaluation of the validity of neighborhood safety is a novel contribution to the literature.

Methods

Sample Selection

Data for this investigation come from the 2006 *Boston Youth Survey* (BYS), a biennial survey of high school students (9th–12th graders) in selected Boston Public Schools. We used a two-stage, stratified random sampling strategy. The first sampling frame consisted of all 34 high schools in the Boston Public Schools system. Thirty schools were randomly selected for the survey, with a probability of selection proportional to each school's enrollment size. Eighteen schools agreed to participate.

Among the participating schools, we generated a numbered list of unique homeroom classes within each school. First, classrooms comprised of students with severe cognitive disabilities were excluded. Next, classrooms were stratified by grade, and then randomly selected for survey administration within each grade. Those classrooms that listed fewer than five students were skipped and the next randomly selected classroom was chosen. Selection continued until the total number of students to be surveyed ranged from 100–125 per school. In the two selected schools with total enrollments close to 100, all students in the school were invited to participate.

Data Collection

The BYS data collection instrument was developed by study staff. The instrument covered a range of topics (e.g., health behaviors, use of school and community resources, and indicators of positive youth development), and had a particular emphasis on violence. Items addressing violence inquired about aggressive behavior, victimization and assault, witnessed violence, fear of violence, and weapon carrying. The paper-and-pencil survey was administered in classrooms by trained staff in the spring

of 2006. Survey administrators completed a brief training program prior to going into the schools. All personnel underwent training in the ethical treatment of human subjects at the Harvard School of Public Health.

Surveys were not marked with any information that could identify an individual. Passive consent was sought from students' parents prior to survey administration. Any student whose parent sent back a form denying permission for the student to participate in the survey was not given one; this was the case for less than 1% of students. Survey administrators read an introduction and the informed consent statement prior to distributing the survey. Seventy of the 1,323 invited students (5.3%) declined to participate. Survey administrators remained in the room and were available to answer questions throughout the 50 minutes allotted for the survey. The Office of Human Research Administration at the Harvard School of Public Health approved all procedures for this research project.

Study Variables

The primary outcome variable was whether adolescents were overweight or at-risk for becoming overweight. To create this variable, we first calculated body mass index (BMI) using respondents' answers to items on height and weight, i.e., weight in kilograms divided by the square of height in meters. We then used BMI to classify respondents as underweight, healthy weight, at-risk for being overweight, or overweight, using age- and sex-specific BMI cut-offs based on Centers for Disease Control and Prevention (CDC) growth charts from the year 2000 [39]. Adolescents who were at or above the 85th percentile were classified as being at-risk for overweight, while those at or above the 95th percentile were classified as overweight. The four-level weight classification variable was subsequently reduced to a dichotomous variable of "at-risk/overweight" and "underweight/normal".

The primary predictor variable was adolescents' perceptions of neighborhood safety. We assessed it with a question designed to capture a global perception of neighborhood safety, i.e., "Do you feel safe in your neighborhood?". The item had the following four response options: always, sometimes, rarely, and never, and was dichotomized (i.e., always/sometimes and rarely/never) for analyses. To assess the validity of perceptions of neighborhood safety, we evaluated the association between perceptions of safety with: (1) beliefs about the seriousness of gang activity in the neighborhood or school, (2) having witnessed someone in the neighborhood being attacked with a weapon (other than a gun) in the past 12 months, and (3) having witnessed someone being physically attacked (i.e. punched, kicked, choked, or beaten up) in the neighborhood in the past 12 months.

Covariates included: age (≤ 14 – ≥ 18 years), grade level (9th–12th), sex (male, female), nativity (U.S. born, foreign-born), and race/ethnicity. To assess race, students were asked to indicate if they were: White; American Indian or Alaska Native; Asian; Black or African American; Native Hawaiian or Other Pacific Islander; Some Other Race, or any combination of those options. We combined Hispanic/Latino ethnicity and race to create a race/ethnicity variable with the following four levels: (1) Hispanic/Latino; (2) non-Hispanic Black/African American; (3) non-Hispanic White; and (4) Other, which includes non-Hispanic bi- or multi-racial students, Asians, American Indians, and other racial groups. To preserve respondent confidentiality, statistics by sub-groups within the "Other" category are not reported.

Statistical Analysis

Although there were 1,253 surveys collected in the 18 schools, the surveys of 38 students (3%) were excluded from data analysis: 35 because they left at least 80% of the items unanswered, and 3 because of erratic answering patterns. Of the remaining 1,215 respondents, 75 were restricted from the analysis sample because they did not answer the items on height and/or weight (and therefore we could not create a BMI variable for them) or because they did not answer the perceived neighborhood safety item. This resulted in an analytic sample size of 1,140.

Data analysis was performed using SAS statistical software version 9.1.3 [40]. First, we generated descriptive information for socio-demographic characteristics of the sample, perceived neighborhood safety, and correlates of perceived neighborhood safety (i.e., beliefs about gang violence, witnessed a physical attack with a weapon other than a gun in the past 12 months, and witnessed someone being physically assaulted in the past 12 months). Next, chi-square statistics and corresponding *p*-values were computed to assess group differences in overweight status by socio-demographic factors, and in perceived neighborhood safety by the associated dimensions. We implemented a multiple comparisons test for proportions using the methods established by Zar [41] for the socio-demographic factors; this was accomplished by using the COMPPROP macro in SAS [42].

We examined whether the association between neighborhood safety and overweight/at-risk for overweight status varied by sex by fitting sex-stratified models because we hypothesized different mechanisms for adolescent boys and girls. Prevalence ratios (PRs) and 95% confidence intervals (CI) were calculated to examine the bivariate association between perceived neighborhood safety and the prevalence of being overweight or at-risk for becoming overweight. We then fit a multiple regression model, in

which we adjusted for socio-demographic factors as appropriate based on group differences. If associated with both the predictor and outcome variables, we included them in the final model. In the final model, we controlled for clustering of students within schools by fitting a generalized estimating equation (GEE) model, with the cluster variable specified as school. Because overweight was analyzed as a dichotomous variable, we specified a binomial response distribution. We used a log link function to relate the expected value of the outcome to the predictor because the data are cross-sectional [43,44]. GEE models were fit using PROC GENMOD in SAS, with school specified as the subject in the REPEATED statement. Listwise deletion was used in regression analyses, i.e., individuals with missing data on any of the covariates were excluded from regression models. Statistical significance was determined by 95% CIs.

Results

More than half of the respondents in the analysis sample (n = 1,140) were female (57.8%), and 46.5% were non-Hispanic Black (Table 1). The mean age was 16.3 years (SD = 1.3). The majority of respondents (54.1%) were in the normal BMI range, 1.1% were underweight, 17.5% were at-risk for becoming overweight, and 27.3% were overweight. Almost one-third of the students were born outside of the U.S. (29.8%).

Although there was no difference in overweight/at-risk for overweight status by sex, age or nativity, there were statis-

tically significant group differences in weight status by race/ethnicity and by grade (Table 1). One-half of the Hispanic students were at-risk or overweight (49.6%), compared to 45.6% of Blacks, 39.3% of Whites, and 33.9% of those in the Other race/ethnicity group. Ninth graders were the most likely to be at-risk or overweight (50.2%), following by tenth graders (44.7%), twelfth graders (43.9%), and eleventh graders (39.1%). Multiple comparisons tests for proportions showed that the only statistically significant pairwise differences were between Hispanics and students in the Other race/ethnicity group, and between the ninth and eleventh grade students.

More than one-third (35.6%) of students said they always felt safe in their neighborhood, 43.9% said they sometimes felt safe, 11.6% rarely felt safe, and 8.9% never felt safe. There were no statistically significant differences in the proportion of students who said they rarely or never feel safe in their neighborhood by sex or age. There were, however, statistically significant differences in perceptions by race/ethnicity (p < 0.001), nativity (p < 0.05), and grade level (p < 0.015). Blacks, immigrants and twelfth graders were significantly more likely than others to report that they rarely or never felt safe in their neighborhoods. One-quarter of Blacks reported they rarely or never felt safe in their neighborhood, compared to 18.2% of Hispanics, 17.8% of Whites, and 9% of students in the Other race/ethnicity group. Compared to 18.7% of US-born students, nearly one-quarter of foreign-born students rarely or never felt safe in their neighborhood.

Table 1: Socio-demographic characteristics of respondents by overweight status

	Total n = 1,140 n (%)	Normal or underweight n = 629 n (%)	At risk or overweight n = 511 n (%)	χ ² Statistic (p-value)
Race/Ethnicity				10.62 (0.014)
Hispanic	335 (30.0)	169 (27.4)	166 (33.2)	
Black, non-Hispanic	520 (46.5)	283 (45.8)	237 (47.4)	
White, non-Hispanic	145 (13.0)	88 (14.2)	57 (11.4)	
Other ^a	118 (10.6)	78 (12.6)	40 (8.0)	
Sex				1.02 (0.311)
Male	481 (42.2)	257 (40.9)	224 (43.8)	
Female	659 (57.8)	372 (59.1)	287 (56.2)	
Nativity				3.54 (0.060)
US Born	788 (70.2)	420 (67.9)	368 (73.0)	
Foreign Born	335 (29.8)	199 (32.2)	136 (27.0)	
Grade				8.08 (0.044)
Ninth	325 (29.0)	162 (26.1)	163 (32.7)	
Tenth	293 (26.1)	162 (26.1)	131 (26.3)	
Eleventh	332 (29.6)	202 (32.5)	130 (26.3)	
Twelfth	171 (15.3)	96 (15.4)	75 (15.0)	

Note. Totals may not sum to 1,140 due to missing data.

^a The Other category includes non-Hispanics students who chose two or more races, or who were Asian, American Indian/Alaska Native, Native Hawaiian or Other Pacific Islander. Information about these groups are not reported separately to preserve confidentiality.

Those students who reported that they rarely or never feel safe in their neighborhoods were more likely than those who said they always or sometimes feel safe to believe that gang violence was a serious problem in their neighborhood or school, and to have seen someone in their neighborhood attacked with a weapon (other than a firearm) in the past 12 months (Table 2). This finding suggests that those who felt unsafe had more exposure to neighborhood violence. Adolescents' who reported that they rarely or never feel safe in their neighborhoods were no more likely than those who said they always or sometimes feel safe to have seen someone in their neighborhood physically assaulted in the past 12 months, indicating that seeing fights may be a frequent occurrence for adolescents regardless of neighborhood.

Next we ran regression models to estimate the effect of perceived neighborhood safety on overweight/at-risk for overweight status. Although the models were initially run stratified by sex, we present the pooled results because prevalence ratios were similar across both sexes. The first model, of the crude association between perception of neighborhood safety and overweight status, indicates that those who rarely or never felt safe in their neighborhoods were 1.21 times more likely to be at-risk for overweight or overweight as compared to those who always or sometimes felt safe in their neighborhoods (95% CI: 1.05, 1.40). Because race/ethnicity and grade level were associated with overweight status and with neighborhood safety, we adjusted for these factors in a second model; the magnitude of the association is attenuated by about 4% (PR = 1.16, 95% CI: 0.99, 1.35). In the third final model, we adjusted for race/ethnicity, grade level, and controlled for clustering of observations by school. Although we found that those who said they rarely or never feel safe in their neighborhoods were 1.16 times more likely to be overweight or at-risk for overweight as compared to adolescents who said they always or sometimes feel safe, this result was not statistically significant (95% CI = 0.97, 1.38). We reran the series of regression models stratifying by race/ethnicity since 1) we were interested in parameter estimates by race/ethnicity, 2) race/ethnicity was associated with both the independent variable and outcome variable, and 3) the distribution of students by race/eth-

nicity varied substantially by school (Table 3). Interestingly, those in the Other race/ethnicity group who reported rarely or never feeling safe in their neighborhood were more likely to be at-risk or overweight in the fully adjusted model (PR = 1.56, 95% CI: 1.02, 2.40).

Discussion

This population-based study of public high school students in Boston, MA adds to the accumulating body of evidence on the association between neighborhood safety and adolescent overweight. Our data suggest that feeling unsafe in one's neighborhood may be associated with an increased risk for overweight. However, this finding was only statistically significant among those within the Other race/ethnicity group. The Other group is comprised mainly of Asians and South Asians (65%), but also includes non-Hispanic bi- or multi-racial students, American Indians and Alaska Natives, Native Hawaiians and Other Pacific Islanders, and other students whose race did not fit into any of the other categories (e.g. those who were Guyanese, Belizean, or Brazilian). Thirty-eight percent of the students in the Other category were immigrants, compared to 34% of Hispanics, 31% of Blacks, and 12% of Whites.

Our findings are consistent with some previous studies that have found an association between neighborhood safety and adolescent overweight status [29,30], but inconsistent with others that did not find an association [18,19,31,32]. Few studies have examined the association between neighborhood safety and adolescent overweight stratified by race/ethnicity [32]. Our findings highlight the importance of considering the moderating effects of race/ethnicity in the association between neighborhood safety and overweight status. The magnitude of the association varied by race/ethnicity and the association was statistically significant for only one racial/ethnic group. We were surprised that the association held for only those within the Other race/ethnicity group; students in the Other group were the least likely to feel unsafe in their neighborhoods and were the least likely to be at-risk or overweight. We expected to find associations among Blacks and Hispanics, given that Black and Hispanic adolescents report particularly high levels of neighborhood violence expo-

Table 2: Perceptions of neighborhood safety by dimensions of neighborhood violence

	Believe there is a lot of gang violence in the neighborhood or school	Saw someone attacked with a weapon (other than a gun) in the neighborhood^b	Saw someone physically assaulted in the neighborhood^b
Rarely or never feel safe	68.0%	17.8%	25.0%
Always or sometimes feel safe	44.1%	11.3%	22.3%
p-value^a	<0.001	0.025	0.463

^ap for chi-squared test for perceived neighborhood safety by the associated dimensions.

^bTime referent is in the past 12 months.

Table 3: Prevalence ratios (and 95% confidence intervals) estimating overweight status among students who rarely or never feel safe in their neighborhood

	Model 1 PR (95% CI)	Model 2^a PR (95% CI)	Model 3^b PR (95% CI)
Hispanic	1.24 (0.98, 1.59)	1.16 (0.91, 1.49)	1.16 (0.89, 1.50)
Black, Non-Hispanic	1.09 (0.88, 1.34)	1.10 (0.99, 1.36)	1.10 (0.90, 1.34)
White, Non-Hispanic	1.42 (0.82, 2.46)	1.25 (0.71, 2.18)	1.23 (0.85, 1.79)
Other	1.54 (0.90, 2.64)	1.58 (0.91, 2.73)	1.56 (1.02, 2.40)

Note. PR = Prevalence Ratio. CI = Confidence Interval. Reference Category = Students who always or sometimes feel safe in their neighborhood

^a Adjusted for grade level using a generalized linear regression model.

^b Adjusted for grade level, using a generalized estimating equation model that accounted for clustering by school.

sure [45-49] and given that these adolescents are more likely to be overweight compared to Whites [10-12,37,38]. Interestingly, the magnitude of the association found among those within the Other race/ethnicity group was weaker in our study compared to the two other studies finding an association [29,30], perhaps a result of using prevalence ratios as the measure of effect (since these studies relied on odds ratios which likely produce inflated effect estimates when there is a relative high disease prevalence [33-36], such as adolescent overweight) [10,11,37,38] and/or because both studies found associations with traffic-related neighborhood safety (though the studies examined other safety-related variables). Because we examined general feelings of neighborhood safety, we can only speculate on which of the multiple aspects of neighborhood safety (e.g. violence/crime, traffic and road hazards, neighborhood disorder) might influence overweight status, but our interviews with students during the pilot testing phase strongly suggest that neighborhood violence is their primary neighborhood safety concern.

There are several pathways by which neighborhood safety might be related to adolescent overweight. One possible interpretation of our findings is that adolescents' concerns about neighborhood safety might decrease their willingness to engage in outdoor physical activity (e.g. walking and playing sports in their neighborhood), promote their use of non-ambulatory transportation options (e.g. use of buses, subways, and automobiles), and/or encourage sedentary behaviors (e.g. television watching, playing computer games, and playing video games in the home), all of which could contribute to being overweight [50,51]. Residing in an unsafe neighborhood might also increase stress (causing a release of cortisol) and result in overweight [52-55]. Evidence indicates that exposure to neighborhood violence, which is a potentially chronic traumatic stressor, is associated with increased cortisol secretion in adolescents [56].

Additionally, since we do not know whether students who perceived their neighborhoods to be unsafe *actually* live in unsafe neighborhoods and we are not aware of any study in this area that has conducted a validity check of percep-

tions of neighborhood safety, we explored whether students who reported feeling unsafe were more likely to have experienced neighborhood violence. We found that those students who felt unsafe in their neighborhoods were more likely than those who felt safe to believe that gang violence was a serious problem in their neighborhood or school and to have seen someone in their neighborhood attacked with a weapon (other than a firearm) in the past 12 months. However, they were not significantly more likely to have seen someone beaten up in their neighborhood in the past 12 months. These findings potentially indicate that witnessing a physical assault may not be a salient dimension of perceived neighborhood safety for adolescents, while neighborhood gang violence and seeing someone in their neighborhood assaulted with a weapon are important aspects of perceived neighborhood safety for them. This is an important contribution to the literature.

There is a need for additional research to clarify the role that neighborhood safety (including neighborhood violence) plays in the adolescent overweight epidemic and to understand salient aspects of perceptions of neighborhood safety. Perceived neighborhood safety is a complex, multidimensional psychosocial construct. Because there is no consensus on the definition of neighborhood safety in health research, qualitative research to explore the dimensions of neighborhood safety and to determine which dimensions of safety are most salient to adolescents at risk of overweight is warranted; this may vary by race/ethnicity. As all of the studies examining neighborhood safety and overweight in adolescents were cross-sectional, researchers should examine this association with prospective cohort designs. In addition, experimental research, e.g., cluster randomized trials (e.g. where neighborhoods might be randomly assigned to an intervention that improves safety) and natural experiments (e.g. new policies promoting police presence in certain neighborhoods to enhance safety), could be conducted (as they are the strongest evidence for temporality) to understand the effects of neighborhood safety on adolescent overweight. Neighborhood environmental interventions [57] and residential mobility experiments [58-60] hold promise to

reduce the prevalence of adolescent overweight. Beyond examining perceived neighborhood safety, research can also examine objective neighborhood safety (e.g. crime statistics to ascertain one's proximity to neighborhood violence) in relation to overweight. Including both subjective and objective reports of neighborhood safety in the same study, although these concepts are likely interconnected, can be beneficial. Each measure might capture distinct neighborhood features; thus, this strategy might ensure optimal measurements of neighborhood safety features.

This study has implications for primary and secondary prevention of adolescent overweight through the development of contextually-relevant interventions and policies. Adolescents in our sample report fear in their neighborhood and high levels of exposure to neighborhood violence, as others [45-49] have shown. This is concerning on its own and also because neighborhood safety may be a factor in adolescent overweight. Our study underscores the importance of policy-level overweight prevention strategies via reducing neighborhood safety concerns. A relevant policy intervention is crime prevention through environmental design [61,62], which would involve changes to the physical environment (such as elimination of hiding spots, landscaping trees and shrubs, and increased surveillance via increased lighting, closed-circuit television/surveillance cameras in public spaces, and/or security guards). Problem-oriented policing, i.e. increased local police attention in "hot spots" or high-crime locations, is another method used to reduce and prevent crime and violence [61,62]. Other potential strategies to reduce neighborhood safety concerns are revitalizing neighborhood watch programs to monitor criminal activity and liaising with police to enhance the protection of places where one can engage in physical activity (e.g. parks and recreation facilities). Furthermore, interventions that offer adolescents' safe havens (such as after-school programs) [63], those that are focused on community development (e.g. ensuring neighborhood resources [such as organizations, services, and employment opportunities]) [64,65], and building collective efficacy among community members [66] could prove beneficial to reduce neighborhood violence. Lastly, it is imperative that behavior change programs (e.g. behavioral weight loss programs) recognize the neighborhood social context of the participants. Physicians, for example, should recommend indoor physical activity for overweight prevention and weight management for adolescents who reside in unsafe neighborhoods.

Study Limitations

These findings should be interpreted in light of the limitations of our study. First, we relied on cross-sectional data;

thus, the study does not inform us about the direction of causation (e.g. whether the exposure preceded the outcome). However, despite the well-known limitations of cross-sectional data, our study hypotheses and directionality have intuitive appeal and were based on conclusions from past research. Additionally, we did not evaluate specific dimensions of neighborhood safety (as previously mentioned) and we did not evaluate objective measures of neighborhood safety (e.g. crime statistics or statistics on the number of sex offenders); we were particularly interested in understanding perceived neighborhood safety rather than the actual occurrence of neighborhood crime or violence. Third, we relied on self-reported height and weight data for BMI, which has the potential for misclassification because of inaccurate reporting. Past research, however, has found that adolescents can provide valid reports of height and weight [67]. Though the gold standard is to collect objectively measured height and weight data, this was not practical nor a central focus of the parent study. Residual confounding might also be a concern, as the survey might have excluded important confounding variables associated with both the independent variable and the dependent variable (e.g. household income, parental education and residential stability might be confounders), but we were unable to account for these variables in the adjusted regression analyses because they were not asked in the BYS data collection instrument. Due to expected high rates of non-response, we did not ask these questions. Finally, this study was conducted in one specific geographically-defined population; thus, these findings might only be generalizable to adolescents in comparable urban locations.

Conclusion

This study adds to the evidence base that neighborhood safety may be associated with overweight status among urban adolescents in certain racial/ethnic groups. Policies and programs should continue to be implemented to reduce neighborhood safety concerns (such as gang activity and witnessing violence) to prevent adolescent overweight.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

DTD conceived the study, assisted with the statistical analysis, interpreted the results, and drafted the manuscript. RMJ assisted with the study design, performed the statistical analysis, interpreted the results, and assisted with writing the manuscript. BEM and DA coordinated the overall survey implementation and data collection, and critically revised the manuscript for substantial intellectual content. All authors have read and approved the final manuscript.

Acknowledgements

DTD was supported in part by a National Institute of General Medical Sciences training grant (grant # 5R25GM055353-10; Ichiro Kawachi, MD, PhD, Principal Investigator). RMJ was supported in part by a National Center on Minority Health and Health Disparities grant (grant # IP20MD002290) awarded to the University of Massachusetts Boston (Celia Moore, PhD, Principal Investigator) and to the Harvard School of Public Health (Felton Earls, MD, Principal Investigator). The Boston Youth Survey 2006 (BYS) was funded by a grant from the Centers for Disease Control and Prevention (grant # U49CE00740) to the Harvard Youth Violence Prevention Center (David Hemenway, PhD, Principal Investigator). BYS was conducted in collaboration with the City of Boston and Mayor Thomas M. Menino. The survey would not have been possible without the participation of the faculty, staff, administrators, and students of Boston Public Schools. We also acknowledge the work of Daria Fanelli, Alicia Savannah, Angela Browne, and Steve Lippmann. We appreciate the assistance of Mary Vriniotis, MS with data collection and management. We thank David Hemenway, PhD and Matthew Miller, MD, MPH, ScD for commenting on an early version of this manuscript, and we thank Gheorghe Doros, PhD for assisting with statistical analysis.

References

- Fields AE: **Predictors and Consequences of Childhood Obesity.** In *Obesity Epidemiology: Methods and Applications* Edited by: Hu FB. Oxford, UK: Oxford University Press; 2008:416-436.
- McCarthy WJ, Yancey AK, Siegel JM, Wong WK, Ward A, Leslie J, Gonzalez E: **Correlation of obesity with elevated blood pressure among racial/ethnic minority children in two Los Angeles middle schools.** *Prev Chronic Dis* 2008, **5(2)**:A46.
- Daniels SR: **The consequences of childhood overweight and obesity.** *Future Child* 2006, **16(1)**:47-67.
- Guerra S, Wright AL, Morgan WJ, Sherrill DL, Holberg CJ, Martinez FD: **Persistence of asthma symptoms during adolescence: role of obesity and age at the onset of puberty.** *Am J Respir Crit Care Med* 2004, **170(1)**:78-85.
- Gilliland FD, Berhane K, Islam T, McConnell R, Gauderman WJ, Gilliland SS, Avol E, Peters JM: **Obesity and the risk of newly diagnosed asthma in school-age children.** *Am J Epidemiol* 2003, **158(5)**:406-15.
- Li L, Law C, Power C: **Body mass index throughout the life-course and blood pressure in mid-adult life: a birth cohort study.** *J Hypertens* 2007, **25(6)**:1215-23.
- Dietz WH: **Childhood weight affects adult morbidity and mortality.** *J Nutr* 1998, **128(2 Suppl)**:411S-414S.
- Must A, Jacques PF, Dallal GE, Bajema CJ, Dietz WH: **Long-term morbidity and mortality of overweight adolescents. A follow-up of the Harvard Growth Study of 1922 to 1935.** *N Engl J Med* 1992, **327(19)**:1350-5.
- Fairfield KM, Willett WC, Rosner BA, Manson JE, Speizer FE, Hankinson SE: **Obesity, weight gain, and ovarian cancer.** *Obstet Gynecol* 2002, **100(2)**:288-96.
- Anderson PM, Butcher KE: **Childhood obesity: trends and potential causes.** *Future Child* 2006, **16(1)**:19-45.
- Ogden CL, Flegal KM, Carroll MD, Johnson CL: **Prevalence and trends in overweight among US children and adolescents, 1999-2000.** *JAMA* 2002, **288(14)**:1728-32.
- Troiano RP, Flegal KM, Kuczmarski RJ, Campbell SM, Johnson CL: **Overweight prevalence and trends for children and adolescents. The National Health and Nutrition Examination Surveys, 1963 to 1991.** *Arch Pediatr Adolesc Med* 1995, **149(10)**:1085-91.
- Bennett GG, Wolin KY, Duncan DT: **Social Determinants of Obesity.** In *Obesity Epidemiology: Methods and Applications* Edited by: Hu FB. Oxford, UK: Oxford University Press; 2008:342-376.
- Sallis JF, Glanz K: **The role of built environments in physical activity, eating, and obesity in childhood.** *Future Child* 2006, **16(1)**:89-108.
- Carver A, Timperio A, Crawford D: **Perceptions of neighborhood safety and physical activity among youth: the CLAN study.** *J Phys Act Health* 2008, **5(3)**:430-44.
- Grow HM, Saelens BE, Kerr J, Durant NH, Norman GJ, Sallis JF: **Where are youth active? Roles of proximity, active transport, and built environment.** *Med Sci Sports Exerc* 2008, **40(12)**:2071-9.
- Singh GK, Yu SM, Siahpush M, Kogan MD: **High levels of physical inactivity and sedentary behaviors among US immigrant children and adolescents.** *Arch Pediatr Adolesc Med* 2008, **162(8)**:756-63.
- Veugelers P, Sithole F, Zhang S, Muhajarine N: **Neighborhood characteristics in relation to diet, physical activity and overweight of Canadian children.** *Int J Pediatr Obes* 2008, **3(3)**:152-9.
- Evenson KR, Scott MM, Cohen DA, Voorhees CC: **Girls' perception of neighborhood factors on physical activity, sedentary behavior, and BMI.** *Obesity* 2007, **15(2)**:430-45.
- Mota J, Gomes H, Almeida M, Ribeiro JC, Santos MP: **Leisure time physical activity, screen time, social background, and environmental variables in adolescents.** *Pediatr Exerc Sci* 2007, **19(3)**:279-90.
- Evenson KR, Birnbaum AS, Bedimo-Rung AL, Sallis JF, Voorhees CC, Ring K, Elder JP: **Girls' perception of physical environmental factors and transportation: reliability and association with physical activity and active transport to school.** *Int J Behav Nutr Phys Act* 2006, **3**:28.
- Heitzler CD, Martin SL, Duke J, Huhman M: **Correlates of physical activity in a national sample of children aged 9-13 years.** *Prev Med* 2006, **42(4)**:254-60.
- Li M, Dibley MJ, Sibbritt D, Yan H: **Factors associated with adolescents' physical inactivity in Xi'an City, China.** *Med Sci Sports Exerc* 2006, **38(12)**:2075-85.
- Carver A, Salmon J, Campbell K, Baur L, Garnett S, Crawford D: **How do perceptions of local neighborhood relate to adolescents' walking and cycling?** *Am J Health Promot* 2005, **20(2)**:139-47.
- Romero AJ: **Low-income neighborhood barriers and resources for adolescents' physical activity.** *J Adolesc Health* 2005, **36(3)**:253-9.
- Gomez JE, Johnson BA, Selva M, Sallis JF: **Violent crime and outdoor physical activity among inner-city youth.** *Prev Med* 2004, **39(5)**:876-81.
- Molnar BE, Gortmaker SL, Bull FC, Buka SL: **Unsafe to play? Neighborhood disorder and lack of safety predict reduced physical activity among urban children and adolescents.** *Am J Health Promot* 2004, **18(5)**:378-86.
- Gordon-Larsen P, McMurray RG, Popkin BM: **Determinants of adolescent physical activity and inactivity patterns.** *Pediatrics* 2000, **105(6)**:E83.
- Mota J, Delgado N, Almeida M, Ribeiro JC, Santos MP: **Physical Activity, Overweight, and Perceptions of Neighborhood Environments Among Portuguese Girls.** *J Phys Act Health* 2006, **3**:314-322.
- Timperio A, Salmon J, Telford A, Crawford D: **Perceptions of local neighbourhood environments and their relationship to childhood overweight and obesity.** *Int J Obes* 2005, **29(2)**:170-5.
- Grafova IB: **Overweight children: assessing the contribution of the built environment.** *Prev Med* 2008, **47(3)**:304-8.
- Ward DS, Dowda M, Trost SG, Felton GM, Dishman RK, Pate RR: **Physical activity correlates in adolescent girls who differ by weight status.** *Obesity* 2006, **14(1)**:97-105.
- Krieger N, Chen JT, Ware JH, Kaddour A: **Race/ethnicity and breast cancer estrogen receptor status: impact of class, missing data, and modeling assumptions.** *Cancer Causes Control* 2008, **19(10)**:1305-18.
- Behrens T, Taeger D, Wellmann J, Keil U: **Different methods to calculate effect estimates in cross-sectional studies. A comparison between prevalence odds ratio and prevalence ratio.** *Methods Inf Med* 2004, **43(5)**:505-9.
- Thompson ML, Myers JE, Kriebel D: **Prevalence odds ratio or prevalence ratio in the analysis of cross sectional data: what is to be done?** *Occup Environ Med* 1998, **55(4)**:272-7.
- Axelsson O, Fredriksson M, Ekberg K: **Use of the prevalence ratio v the prevalence odds ratio as a measure of risk in cross sectional studies.** *Occup Environ Med* 1994, **51(8)**:574.
- Ogden CL, Carroll MD, Curtin LR, McDowell MA, Tabak CJ, Flegal KM: **Prevalence of overweight and obesity in the United States, 1999-2004.** *JAMA* 2006, **295(13)**:1549-55.

38. Hedley AA, Ogden CL, Johnson CL, Carroll MD, Curtin LR, Flegal KM: **Prevalence of overweight and obesity among US children, adolescents, and adults, 1999–2002.** *JAMA* 2004, **291(23)**:2847-50.
39. Kuczmarski RJ, Ogden CL, Guo SS, Grummer-Strawn LM, Flegal KM, Mei Z, Wei R, Curtin LR, Roche AF, Johnson CL: **2000 CDC Growth Charts for the United States: methods and development.** *Vital Health Stat* 2002, **11(246)**:1-190.
40. SAS Institute Inc: **Statistical Applications Software, Version 9.1.3.** Cary, NC 2008.
41. Zar JH: **Biostatistical Analysis, Fourth Edition.** Prentice Hall; New Jersey; 1999:564.
42. Elliott AC, Reisch JS: **Implementing a multiple comparison test for proportions in a 2x2 crosstabulation in SAS®.** *Proceedings of the SAS User's Group International 31, San Francisco, CA, March 26–29, 2006. Paper 204-31*.
43. Spiegelman D, Hertzmark E: **Easy SAS calculations for risk or prevalence ratios and differences.** *Am J Epidemiol* 2005, **162(3)**:199-200.
44. Skov T, Deddens J, Petersen MR, Endahl L: **Prevalence proportion ratios: estimation and hypothesis testing.** *Int J Epidemiol* 1998, **27(1)**:91-5.
45. Buka SL, Stichick TL, Birdthistle I, Earls FJ: **Youth exposure to violence: prevalence, risks, and consequences.** *Am J Orthopsychiatry* 2001, **71(3)**:298-310.
46. Stein BD, Jaycox LH, Kataoka S, Rhodes HJ, Vestal KD: **Prevalence of child and adolescent exposure to community violence.** *Clin Child Fam Psychol Rev* 2003, **6(4)**:247-64.
47. Bain JE, Brown RT: **Adolescents as witnesses to violence.** *J Adolesc Health* 1996, **19(2)**:83-5.
48. Fitzpatrick KM, Boldizar JP: **The prevalence and consequences of exposure to violence among African-American youth.** *J Am Acad Child Adolesc Psychiatry* 1993, **32(2)**:424-30.
49. Schubiner H, Scott R, Tzelepis A: **Exposure to violence among inner-city youth.** *J Adolesc Health* 1993, **14(3)**:214-9.
50. Hu FB: **Physical Activity, Sedentary Behaviors, and Obesity.** In *Obesity Epidemiology: Methods and Applications* Edited by: Hu FB. Oxford, UK: Oxford University Press; 2008:301-319.
51. Must A, Tybor DJ: **Physical activity and sedentary behavior: a review of longitudinal studies of weight and adiposity in youth.** *Int J Obes* 2005, **29(Suppl 2)**:S84-96.
52. Hu FB: **Metabolic and Hormonal Predictors of Obesity.** In *Obesity Epidemiology: Methods and Applications* Edited by: Hu FB. Oxford, UK: Oxford University Press; 2008:377-398.
53. Roemmich JN, Smith JR, Epstein LH, Lambiase M: **Stress reactivity and adiposity of youth.** *Obesity* 2007, **15(9)**:2303-10.
54. Björntorp P: **Do stress reactions cause abdominal obesity and comorbidities?** *Obes Rev* 2001, **2(2)**:73-86.
55. Rosmond R, Dallman MF, Björntorp P: **Stress-related cortisol secretion in men: relationships with abdominal obesity and endocrine, metabolic and hemodynamic abnormalities.** *J Clin Endocrinol Metab* 1998, **83(6)**:1853-9.
56. Kliewer W: **Violence exposure and cortisol responses in urban youth.** *Int J Behav Med* 2006, **13(2)**:109-20.
57. Farley TA, Meriwether RA, Baker ET, Watkins LT, Johnson CC, Weber LS: **Safe play spaces to promote physical activity in inner-city children: results from a pilot study of an environmental intervention.** *Am J Public Health* 2007, **97(9)**:1625-31.
58. U.S. Department of Housing and Urban Development, Office of Policy Development and Research: **Moving to Opportunity Interim Impacts Evaluation.** 2003 [<http://www.huduser.org/Publications/pdf/MTOFullReport.pdf>]. Accessed September 15, 2008
59. Kingsley GT, Pettit KLS: **Have MTO Families Lost Access to Opportunity Neighborhoods Over Time? Three-City Study of Moving to Opportunity.** Brief No. 2 2008 [http://www.urban.org/UploadedPDF/411637_opportunity_neighborhoods.pdf]. Washington, DC: The Urban Institute Accessed September 15, 2008
60. Kling JR, Liebman JB, Katz LF: **Experimental analysis of neighborhood effects.** *Econometrica* 2007, **75(1)**:83-119.
61. Loukaitou-Sideris A, Eck JE: **Crime prevention and active living.** *Am J Health Promot* 2007, **21(4 Suppl)**:380-9.
62. Mair JS, Mair M: **Violence prevention and control through environmental modifications.** *Annu Rev Public Health* 2003, **24**:209-25.
63. Molnar BE, Roberts AL, Browne A, Gardener H, Buka SL: **What girls need: recommendations for preventing violence among urban girls in the US.** *Soc Sci Med* 2005, **60(10)**:2191-204.
64. Molnar BE, Cerda M, Roberts AL, Buka SL: **Effects of neighborhood resources on aggressive and delinquent behaviors among urban youths.** *Am J Public Health* 2008, **98(6)**:1086-93.
65. Yonas MA, O'Campo P, Burke JG, Gielen AC: **Neighborhood-level factors and youth violence: giving voice to the perceptions of prominent neighborhood individuals.** *Health Educ Behav* 2007, **34(4)**:669-85.
66. Sampson RJ, Raudenbush SW, Earls F: **Neighborhoods and violent crime: a multilevel study of collective efficacy.** *Science* 1997, **277(5328)**:918-24.
67. Goodman E, Hinden BR, Khandelwal S: **Accuracy of teen and parental reports of obesity and body mass index.** *Pediatrics* 2000, **106(1 Pt 1)**:52-8.

Pre-publication history

The pre-publication history for this paper can be accessed here:

<http://www.biomedcentral.com/1471-2458/9/289/prepub>

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

