



# A Method for the Preparation of Differentiated $\backslash(\text{trans}\backslash)$ -1,2-Diol Derivatives with Enantio- and Diastereocontrol

## Citation

Lim, Sang Min, Nicholas Hill, and Andrew G. Myers. 2009. A method for the preparation of differentiated  $\backslash(\text{trans}\backslash)$ -1,2-diol derivatives with enantio- and diastereocontrol. *Journal of the American Chemical Society* 131(16): 5763-5765.

## Published Version

doi:10.1021/ja901283q

## Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:5125262>

## Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Open Access Policy Articles, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#OAP>

## Share Your Story

The Harvard community has made this article openly available.  
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)


# A Method for the Preparation of Differentiated *trans*-1,2-Diol Derivatives with Enantio- and Diastereocontrol

Sang Min Lim, Nicholas Hill, and Andrew G. Myers\*

Department of Chemistry and Chemical Biology, Harvard University, Cambridge, Massachusetts 02138

RECEIVED DATE (automatically inserted by publisher); E-mail: myers@chemistry.harvard.edu

We describe a synthetic sequence that allows for the preparation of optically active *trans*-1,2-diol monosilyl ether derivatives from ketones, providing a new means for retrosynthetic simplification of differentiated diol and polyol targets. The sequence is exemplified in equation 1 and proceeds by silyl enol ether formation, Shi asymmetric epoxidation,<sup>1</sup> then regio- and stereospecific addition of hydride, methide, or higher alkylide. The tactical combination presented has not been integrated in synthetic problem solving, so far as we are aware, but has promise for broad application, we believe.



It has been shown that  $\alpha$ -hydroxy ketones can be prepared enantioselectively by application of the Shi asymmetric epoxidation protocol to *tert*-butyldimethylsilyl and trimethylsilyl enol ether substrates followed by hydrolysis.<sup>2</sup> Shi and coworkers noted that certain substrates were problematic in the transformation, especially silyl enol ethers derived from cyclic ketones (racemization and dimerization were specifically mentioned as complicating factors). This led them to direct their studies toward enol ester substrates, where the primary epoxidation products were sufficiently stable to be isolated and manipulated, providing an essential enablement of their general method for the enantiocontrolled synthesis of  $\alpha$ -acyloxy ketones.<sup>3</sup>

While prior research has shown that trimethylsilyloxy and *tert*-butyldimethylsilyloxy epoxides can be prepared as discrete intermediates, especially by epoxidation with non-acidic oxygen atom-transfer agents such as arylsulfonyloxaziridines<sup>4</sup> and dimethyldioxirane,<sup>5</sup> it is also evident from this prior work that the reactivity of the products complicate their isolation and handling, absent special stabilizing structural features. With these precedents in mind, and aware of the particular stability of *tert*-butyldiphenylsilyl ethers toward acid-catalyzed hydrolysis,<sup>6</sup> we undertook to reinvestigate the Shi asymmetric epoxidation of silyl enol ether derivatives with a focus on variation of the trialkylsilyl group. Here we show that *tert*-butyldiphenylsilyl enol ether derivatives of both cyclic and acyclic ketones undergo enantioselective epoxidation according to the Shi paradigm,<sup>1</sup> that the *tert*-butyldiphenylsilyloxy epoxide products can be isolated by simple extraction, and, most significantly, that these products undergo highly regio- and stereospecific addition reactions with certain hydride and alkylide donors (vide infra).

*tert*-Butyl(cyclohexenyloxy)diphenylsilyl ether (**1**), prepared from cyclohexanone in >95% yield under soft enolization conditions (*tert*-butyldiphenylsilyl triflate,<sup>7</sup> triethylamine, dichloromethane, 0 °C, 1 h)<sup>8</sup> or by deprotonation with potassium hexamethyldisilazide followed by trapping with the silyl chloride, is, as expected, substantially more stable than the corresponding triisopropylsilyl and *tert*-butyldimethylsilyl enol ethers<sup>6,9</sup> (in that order) under conditions such as exposure to silica gel (see Supporting Information). All three substrates were transformed into the corresponding silyloxy epoxides under typical Shi epoxidation conditions (0.30 equiv of Shi catalyst,<sup>10</sup> borate buffer, slow addition of Oxone and potassium carbonate solutions over 90 min at 0 °C) with similar, high enantioselectivities (91–93% ee, vide infra).<sup>2</sup> The products were efficiently extracted into pentane without decomposition.<sup>4</sup> Two-dimensional thin-layer chromatographic analysis of the product solutions revealed the following stability order: *tert*-butyldiphenylsilyl (compound **2**) >> triisopropylsilyl > *tert*-butyldimethylsilyl (see Supporting Information). Solutions of the epoxide **2** could be concentrated and the crude product stored neat for several weeks at –20 °C without evident decomposition. <sup>1</sup>H NMR spectral data were consistent with expectations for the proposed structure (**2**) and revealed that the product had been formed cleanly, with the only significant contaminant being a small amount of Shi catalyst that had partitioned into the pentane layer during the extraction. While future applications may benefit from or require that purified silyloxy epoxides be used, all of the transformations described herein were conducted using the pentane extracts of the Shi epoxidation reaction mixtures directly, without purification, or even concentration in many cases.

Among the new and useful reactions of silyloxy epoxides we have investigated is a simple regio- and stereoselective reduction with borane-THF, which gives rise to *trans*-diol monosilyl ethers.<sup>11</sup> In a specific illustration (eq 1 and entry 1, Table 1), addition of a 1.0 M solution of borane-THF (1.2 equiv) to an ice-cold solution of the pentane extracts containing the Shi epoxidation product **2**, stirring at 0 °C for 1 h, then careful quenching with an aqueous solution of tris(hydroxy-methyl)aminomethane, extractive isolation, and chromatographic purification afforded pure (1*R*,2*R*)-2-(*tert*-butyldiphenylsilyloxy)cyclohexanol **3** in 91% yield for the two-step (epoxidation-reduction) sequence. The product was shown to be of 93% ee by <sup>1</sup>H NMR analysis of the corresponding Mosher ester.<sup>12</sup> Similarly, the triisopropylsilyl and *tert*-butyldimethylsilyl enol ethers (entries 2 and 3, respectively, Table 1) were transformed directly into the corresponding *trans*-1,2-diol monosilyl ethers, in yields that reflected the stabilities of the silyloxy epoxide intermediates.

**Table 1.** Cyclic *trans*-1,2-Diol Monosilyl Ether Derivatives Formed by Shi Asymmetric Epoxidation<sup>a</sup> of Silyl Enol Ethers followed by Stereospecific Reduction with Borane-THF

Entry	Substrate	Product	Yield <sup>b</sup>	ee <sup>c</sup>
1			91	93
2			82	91
3			68	91
4			92	89
5			82	88
6			95	90
7			81	90
8 <sup>d</sup>			81	91
9 <sup>d</sup>			69	90


<sup>a</sup> Shi epoxidation conditions: 0.3 equiv Shi catalyst, 1.38 equiv Oxone, 5.8 equiv K<sub>2</sub>CO<sub>3</sub>, CH<sub>3</sub>CN-CH<sub>3</sub>OCH<sub>2</sub>OCH<sub>3</sub>-borate buffer, 0 °C, 2 h. <sup>b</sup> Isolated yields over two steps. <sup>c</sup> <sup>1</sup>H NMR analysis revealed that in all cases diastereoselectivities were > 20:1, favoring the *trans*-1,2-diol derivatives. <sup>d</sup> Enantiomeric excesses were determined by <sup>1</sup>H NMR analysis of the corresponding Mosher esters, except for entries 7 and 8, where ee's were determined by HPLC using a chiral column (see Supporting Information for details). <sup>d</sup> These entries involved slight procedural modifications; see Supporting Information for details.

As the examples of Table 1 reveal, the two-step Shi epoxidation-reduction sequence appears to hold promise as a general method for the enantio- and diastereocontrolled synthesis of differentiated cyclic *trans*-1,2-diols, and as entry 9 suggests, may also allow for simplification of certain polyol targets by simultaneous multiple application.<sup>13,14</sup> Because epimerization is unlikely to have occurred at any point during the transformations summarized in Table 1, we believe the ee

values presented there provide an accurate assessment of the enantioselectivities of the epoxidation step, which are typically above 90%, in keeping with Shi's prior observations.<sup>1</sup>

The stereochemistry of hydride addition supports a mechanism involving prior coordination of borane to the epoxide oxygen atom followed by epoxide opening and internal hydride transfer, as has been proposed for reductions of glycol epoxide-like substrates with borane-THF,<sup>15</sup> although the trajectories for hydride addition are presumably quite different.<sup>16</sup>


In further evaluating the scope of the epoxidation-reduction sequence we explored acyclic *tert*-butyldiphenylsilyl enol ethers as substrates and in this context gained mechanistic insight into the reduction process. Asymmetric epoxidation-reduction of *trans*-1-*tert*-butyldiphenylsilyloxypropene<sup>17</sup> afforded (*R*)-1-(*tert*-butyldimethylsilyloxy)propan-2-ol in 90% yield and 82% ee (eq 2). Employing BD<sub>3</sub>-THF in lieu of BH<sub>3</sub>-THF we found that the reduction proceeded with >95% stereospecificity (eq 3),<sup>18</sup> as observed in the cyclic substrate series (Table 1). Epoxidation-reduction (also with BD<sub>3</sub>-THF) of *cis*-1-*tert*-butyl-diphenylsilyloxypropene,<sup>19</sup> however, proceeded with substantially diminished stereospecificity (~33%), eq 4), which we rationalize in Figure 1.<sup>20</sup>




**Figure 1.** Reduction of the *cis*-*tert*-butyldiphenylsilyloxy epoxide **4** with BD<sub>3</sub>-THF provides evidence for a short-lived carbocationic intermediate. The data suggest that the rate of deuteride transfer is slightly more rapid than rotation about the internal C-C bond.

Two additional examples in the acyclic series, employing trisubstituted *Z*-*tert*-butyldiphenylsilyl enol ethers as substrates (each ≥ 14:1, *Z*:*E*, prepared by enolate formation with potassium hexamethyldisilazide followed by trapping

with *tert*-butyldiphenylsilyl chloride), suggest that the present method may have general value for the preparation of differentiated *anti*-1,2-diols (eqs 5 and 6).



Lastly, we have observed that *tert*-butyldiphenylsilyloxy epoxides react stereospecifically with trimethyl- and triethylaluminum to form differentiated *trans*-1,2-diol products (eqs 7–9). It is noteworthy that the tertiary hydroxyl groups of the product diols emerge bearing the *tert*-butyldiphenylsilyl protecting group. Thus far, we have not seen evidence of silyl group transfer. Here, too, substantial literature precedent exists for stereospecific additions of trialkylaluminum reagents to glycol epoxides<sup>21</sup> although, as discussed above in the context of hydride addition,<sup>16</sup> the stereoelectronic features of the present transformations are presumably very different and seemingly less favorable.




We imagine that the asymmetric epoxidation-reduction and epoxidation-alkylidene addition sequences presented herein will be useful for the preparation of a number of complex diol and polyol targets.

**Acknowledgment.** We thank the NSF (CHE-0749566), the NIH/NCI (CHE-0749566), the NIH/NIGMS (GM007598-30) (NH), the Kwanjeong Educational Foundation Fellowship (SML), the Eli Lilly Organic Chemistry Fellowship (SML), the Harvard College Research Program, Pfizer Inc., Amgen, and Merck & Co., Inc. for financial support of this research. We thank Dr. Richard Staples and Dr. Douglas Ho for X-ray crystallographic analyses.


**Supporting Information Available:** Detailed experimental procedures and characterization data for all new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

## References

- (a) Tu, Y.; Wang, Z.-X.; Shi, Y. *J. Am. Chem. Soc.* **1996**, *118*, 9806–9807. (b) Wang, Z.-X.; Tu, Y.; Frohn, M.; Zhang, J.-R.; Shi, Y. *J. Am. Chem. Soc.* **1997**, *119*, 11224–11235. (c) Frohn, M.; Shi, Y. *Synthesis* **2000**, 1979–2000. (d) Shi, Y. *Acc. Chem. Res.* **2004**, *37*, 488–496.
- (a) Adam, W.; Fell, R. T.; Saha-Möller, C. R.; Zhao, C.-G. *Tetrahedron: Asymmetry* **1998**, *9*, 397–401. (b) Zhu, Y.; Tu, Y.; Yu, H.; Shi, Y. *Tetrahedron Lett.* **1998**, *39*, 7819–7822. (c) Solladié-Cavallo, A.; Lupattelli, P.; Jierry, L.; Bovicelli, P.; Angeli, F.; Antonioletti, R.; Klein, A. *Tetrahedron Lett.* **2003**, *44*, 6523–6526.
- Zhu, Y.; Manske, K. J.; Shi, Y. *J. Am. Chem. Soc.* **1999**, *121*, 4080–4081.
- Davis, F. A.; Sheppard, A. C. *J. Org. Chem.* **1987**, *52*, 954–955.
- (a) Chenault, H. K.; Danishefsky, S. J. *J. Org. Chem.* **1989**, *54*, 4249–4250. (b) Adam, W.; Hadjiarapoglou, L.; Wang, X. *Tetrahedron Lett.* **1989**, *30*, 6497–6500. (c) Schaumann, E.; Tries, F. *Synthesis* **2002**, 191–194.
- Hanessian, S.; Lavallee, P. *Can. J. Chem.* **1975**, *53*, 2975–2977.
- Dimopoulos, P.; George, J.; Tocher, D. A.; Manaviar, S.; Hale, K. J. *Org. Lett.* **2005**, *7*, 5377–5380.
- For a different method of preparing compound **1**, see: Martel, A.; Leconte, S.; Dujardin, G.; Brown, E.; Maisonneuve, V.; Retoux, R. *Eur. J. Org. Chem.* **2002**, 514–525.
- Cunico, R. F.; Bedell, L. *J. Org. Chem.* **1980**, *45*, 4797–4798.
- The catalyst was prepared in two steps from D-Fructose as described in reference 1b.
- The transformation is functionally equivalent to an asymmetric hydroboration-oxidation of a silyl enol ether substrate. In the one example of such a process of which we are aware cyclohexanone trimethylsilyl enol ether was transformed into *trans*-1,2-cyclohexanediol monotrimethylsilyl ether with diisopinacampheylborane-alkaline hydrogen peroxide in 31% yield and 28% ee: Peterson, P. E.; Stepanian, M. *J. Org. Chem.* **1988**, *53*, 1903–1907.
- Dale, J. A.; Dull, D. L.; Mosher, H. S. *J. Org. Chem.* **1969**, *34*, 2543–2549. The absolute stereochemistry of product **3** was confirmed by comparison (optical rotation) with an authentic sample prepared by an independent route.
- The absolute stereochemistry of the product (entry 9, Table 1) was established by X-ray crystallographic analysis of the corresponding bis-*p*-bromophenyl ester derivative (see Supporting Information). Interestingly, we observed that when a pentane–ethyl acetate solution of the bis-epoxide intermediate of entry 9 was shaken briefly with a 1 N aqueous hydrochloric acid solution, the hemiketal monosilyl ether depicted was obtained as a stable substance. The structure was verified by X-ray crystallography (see Supporting Information).
- (a) Xiong, Z.; Corey, E. J. *J. Am. Chem. Soc.* **2000**, *122*, 9328–9329. (b) Lorenz, J. C.; Frohn, M.; Zhou, X.; Zhang, J.-R.; Tang, Y.; Burke, C.; Shi, Y. *J. Org. Chem.*, **2005**, *70*, 2904–2911.
- (a) Bazin, H. G.; Kerns, R. J.; Linhardt, R. J. *Tetrahedron Lett.* **1997**, *38*, 923–926. (b) Koyama, Y.; Yamaguchi, R.; Suzuki, K. *Angew. Chem. Int. Ed.* **2008**, *47*, 1084–1087.
- We speculate that hydride transfer in the two processes may involve different conformations of the respective six-membered rings:



- Prepared by iridium-catalyzed isomerization of allyloxy(*tert*-butyl)diphenylsilane: Ohmura, T.; Yamamoto, Y.; Miyaura, N.; *Organometallics* **1999**, *18*, 413–416.
- Using a commercial source of  $\text{BD}_3\text{-THF}$  with 11% hydrogen content (determined by  $^1\text{H}$  NMR analysis of the corresponding  $\text{BD}_3\text{-dimethylphenylphosphine}$  complex), we found the product of eq 5 was enriched in hydrogen (17% hydrogen content at the site of transfer), consistent with a kinetic isotope effect of approximately 1.6.
- Guha, S. K.; Shibayama, A.; Abe, D.; Sakaguchi, M.; Ukaji, Y.; Inomata, K. *Bull. Chem. Soc. Jpn.* **2004**, *77*, 2147–2157.
- The products of this two-step transformation were also formed with diminished enantioselectivity (~40% ee), in keeping with prior examples of Shi epoxidations of *cis*-1,2-disubstituted alkenes (see reference 1b).
- (a) Bailey, J. M.; Craig, D.; Gallagher, P. T. *Synlett* **1999**, 132–134. (b) Rainier, J. D.; Cox, J. M. *Org. Lett.* **2000**, *2*, 2707–2709.




---

We describe a synthetic sequence that allows for the preparation of optically active *trans*-1,2-diol monosilyl ether derivatives from ketones, providing a new means for retrosynthetic simplification of differentiated diol and polyol targets. The sequence involves silyl enol ether formation, Shi asymmetric epoxidation, then regio- and stereospecific addition of hydride, methide, or higher alkylide. The tactical combination presented has not been integrated in synthetic problem solving, so far as we are aware, but has promise for broad application, we believe.

---