



Experimental Observables Near a Nematic Quantum Critical Point in the Pnictide and Cuprate Superconductors

Citation

Xu, Cenke, Yang Qi, and Subir Sachdev. 2008. Experimental observables near a nematic quantum critical point in the pnictide and cuprate superconductors. *Physical Review B* 78(13): 134507.

Published Version

[doi://10.1103/PhysRevB.78.134507](https://doi.org/10.1103/PhysRevB.78.134507)

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:7462438>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Open Access Policy Articles, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#OAP>

Share Your Story

The Harvard community has made this article openly available. Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

Experimental observables near a nematic quantum critical point in the pnictide and cuprate superconductors

Kenke Xu, Yang Qi, and Subir Sachdev

Department of Physics, Harvard University, Cambridge, MA 02138

(Dated: August 7, 2008)

The newly discovered high temperature superconductor $\text{SmFeAs}(\text{O}_{1-x}\text{F}_x)$ shows a clear nematic transition where the square lattice of Fe ions has a rectangular distortion. Similar nematic ordering has also been observed in the cuprate superconductors. We provide a detailed theory of experimental observables near such a nematic transition: we calculate the scaling of specific heat, local density of states (LDOS) and NMR relaxation rate $1/T_1T$.

Rapid and important progress has been made in studies of Iron-oxypnictides superconductors. Various samples with similar FeAs plane and rare earths have been synthesized and several different compounds have shown superconductivity over 50K [1, 2, 3, 4, 5, 17], when the parent compound is doped by either fluorine or oxygen deficiency. Although many experimental facts, including the pairing symmetry, are still under debate, all these different samples share common features: a tetragonal-monoclinic (orthorhombic) lattice distortion and $(\pi, 0)$ spin density wave (SDW) which commonly exist in the undoped samples, and “compete” with superconductivity at finite doping. The SDW and lattice distortion were first observed in $\text{LaFeAs}(\text{O}_{1-x}\text{F}_x)$ by elastic neutron scattering and X-ray spectroscopy [6]. Later on this phenomenon was confirmed in many other samples with La replaced by Ce [7], Sm [17, 18] and Nd [19, 20], and also in oxygen free materials BaFe_2As_2 [8, 9], SrFe_2As_2 [12, 13], and CaFe_2As_2 [15, 16]. In all the samples, at the lattice distortion temperature T_{c1} , the resistivity shows a λ shaped anomaly; therefore the λ anomaly of resistivity can be taken as a measure of the lattice distortion in experiments.

Both lattice distortion and SDW are suppressed under doping, but, in general, the lattice distortion occurs at higher temperature than the SDW. In $\text{SmFeAsO}_{1-x}\text{F}_x$, the lattice distortion and superconductivity coexist in a finite range of doping; the lattice distortion temperature seems to vanish within the superconducting phase [17, 18], while the coexistence between SDW and superconductor was never observed. In Ref. [21, 22], the lattice distortion is attributed to anisotropic antiferromagnetic correlation between electrons along x and y directions, without developing long range SDW. Since this order deforms the electron Fermi surface, equivalently, it can also be interpreted as electronic nematic order. This nematic order has Ising symmetry, therefore the transition temperature is controlled by the intralayer spin interaction, while the long range SDW is controlled by the interlayer spin interaction which is much weaker. Therefore the nematic transition (lattice distortion) occurs at a higher temperature than the SDW in general, and unless very close to the critical point, the nematic transition at

finite temperature should belong to the 2d Ising universality class. The distance between the lattice distortion temperature and the SDW temperature depends on the anisotropy between ab plane and c axis, which can be checked by comparing the anisotropy of different samples.

The intimate relation between the structure distortion and SDW phase proposed by Ref. [21, 22] has gained support from recent experiments. It is suggested by detailed X-ray, neutron and Mössbauer spectroscopy studies that both the lattice distortion transition and the SDW transition of $\text{LaFeAs}(\text{O}_{1-x}\text{F}_x)$ are second order [23], where the two transitions occur separately. However, in AFe_2As_2 with $\text{A} = \text{Sr}, \text{Eu}, \text{Ba}, \text{Ca}$, the structure distortion and SDW occur at the same temperature, and the structure distortion becomes a first order transition [10, 12, 13, 14, 16] (or a very steep second order transition [24]). These results suggest that the SDW and structure distortion are indeed strongly interacting with each other, and the structure distortion is probably induced by magnetism.

We also note that the importance of nematic ordering has also been discussed recently in the context of the cuprate superconductors [25, 26]. Our results below are presented in the context of the pnictides, but all of the scaling properties of the experimental observables apply equally to the cuprates.

We focus on the zero temperature nematic phase transition at finite doping, motivated by the experimental suggestion of the existence of structure distortion critical point within the superconducting phase of sample $\text{SmFeAsO}_{1-x}\text{F}_x$ [17, 18]. By contrast, the SDW phase shows no overlap with the superconducting phase in all the samples studied so far, therefore we will generally ignore it except for noting that the transition from the SDW to the nematic order is likely an $z = 1$ $\text{O}(3)$ transition, based on the fact that the SDW order wave vector is independent of doping [7], so the low energy particle-hole excitations at the SDW wave vector vanishes rapidly with small doping and hence make no contribution to the damping of the SDW order parameter [21]. The universality class of the nematic transition strongly depends on the pairing symmetry of the superconducting phase. If it




FIG. 1: The global phase diagram. The blue, red and green curves are phase boundaries of SDW, nematic (structure distortion) and superconductivity respectively.

is an s -wave superconductor without nodes, the transition of nematic order will be an ordinary 3D Ising transition; while if the superconductor is d -wave, the gapless nodal particles may change the universality class of the nematic transition. The recent STM [27] and Andreev reflection measurement [28] suggest that $\text{SmFeAsO}_{1-x}\text{F}_x$ has nodes in the cooper pair, and the spin susceptibility measured by Knight shift will tell us whether it is a p -wave or d -wave pairing. In our current work we assume a d -wave pairing. The universal behavior of the nematic transition in a d -wave superconductor was first studied in Ref. [25]. Recently the same theory was studied carefully, and it was shown that in the infrared limit there is a special fixed point with logarithmically diverging velocity anisotropy of the nodal particles [26]. In the current work we will calculate experimentally relevant quantities close to this nematic quantum critical point. The global phase diagram is shown in Fig. 1.

The low energy Lagrangian describing the nematic order and nodal particle reads [25]

$$\begin{aligned}
 L &= L_\Psi + L_\phi + L_{\Psi\phi}, \\
 L_\Psi &= \sum_{a=1}^{N_f} \Psi_{1a}^\dagger (\partial_\tau - iv_f \partial_x \tau^z - iv_\Delta \partial_y \tau^x) \Psi_{1a} \\
 &\quad + \Psi_{2a}^\dagger (\partial_\tau - iv_f \partial_y \tau^z - iv_\Delta \partial_x \tau^x) \Psi_{2a}, \\
 L_\phi &= \frac{1}{2} (\partial_\tau \phi)^2 + \frac{c^2}{2} (\nabla \phi)^2 + \frac{r}{2} \phi^2 + \frac{u_0}{4!} \phi^4, \\
 L_{\Psi\phi} &= \lambda_0 \phi \sum_{a=1}^{N_f} (\Psi_{1a}^\dagger \tau^x \Psi_{1a} + \Psi_{2a}^\dagger \tau^x \Psi_{2a}). \quad (1)
 \end{aligned}$$

The Nambu fermion Ψ is defined in the standard convention: $\Psi_{1a} = (f_{1a}, \epsilon_{ab} f_{3b}^\dagger)$ and $\Psi_{2a} = (f_{2a}, \epsilon_{ab} f_{4b}^\dagger)$. a and b are spin indices, f_1, f_2, f_3 and f_4 are slow fermion modes at nodal points $(Q, Q), (-Q, Q), (-Q, -Q)$ and




FIG. 2: The Feynman diagram used in this work, the dashed lines are the propagators of nematic order parameter ϕ . a , the self-energy correction to fermion Ψ ; b , the self-energy correction to ϕ ; c , the vertex correction to fermion bilinear $\Psi_i^\dagger T_A \Psi_j$; d , the vertex correction to fermion bilinear $\Psi_i^\dagger T_A \Psi_j$.




FIG. 3: Plot of local density of states with $v_{\Delta 0}/v_{f0} = 1/5$, $v_{\Delta 0}/v_{f0} = 1/10$, $v_{\Delta 0}/v_{f0} = 1/20$ from the top to bottom. The horizontal axis is in scale of ω/T_c . The curves are $y = x^{0.91}$ (red), $y = x^{0.935}$ (blue), $y = x^{0.955}$ (green), $y = x$ (black).

$(Q, -Q)$ respectively. If the system develops long range order of ϕ , the four nodal points of the d -wave superconductor will be shifted and break the C_{4v} symmetry down to C_{2v} due to the coupling $L_{\Psi\phi}$ [25]. The lagrangian (1) is not Lorentz invariant because in the real system v_Δ/v_f is in general not unity. Also, the coupling $L_{\Psi\phi}$ breaks the Lorentz invariance, since $\Psi^\dagger \tau^x \Psi$ is only one component of the space-time current of the Dirac fermion. Therefore a realistic scaling procedure is to allow v_Δ and v_f flow independently under renormalization group (RG).

The nematic transition fixed point in Ref. [26] was obtained by expansion of $v_\Delta/(v_f N_f)$, assuming a small initial value $v_{\Delta 0}/v_{f0}$, and in the current situation $N_f = 2$. The RG flow of velocities is obtained by calculating the one-loop self-energy in Fig. 2a, with dressed ϕ propagator at order $1/N_f$ in Fig. 2b. The renormalization

condition is chosen to be keeping the coupling constant λ_0 in $L_{\Psi\phi}$ invariant under RG flow. After the one-loop correction, the flow of the self-energy and velocities reads

$$\begin{aligned}\frac{d\Sigma_1}{d\ln\Lambda} &= C_1(-i\omega) + C_2v_fk_x\tau^z + C_3v_\Delta k_y\tau^x, \\ \frac{dv_f}{d\ln\Lambda} &= (C_1 - C_2)v_f, \\ \frac{dv_\Delta}{d\ln\Lambda} &= (C_1 - C_3)v_\Delta, \\ \frac{d(v_\Delta/v_f)}{d\ln\Lambda} &= (C_2 - C_3)(v_\Delta/v_f).\end{aligned}\quad (2)$$

Λ is the momentum cut-off. C_1 , C_2 and C_3 are functions of v_Δ/v_f and N_f , their detailed forms are given in the appendix. Using these RG equations, we are ready to calculate the scaling of the local density of states (LDOS) accessible by scanning tunneling microscope (STM):

$$\begin{aligned}\rho(\omega) &\sim \int \frac{dk_x dk_y}{(2\pi)^2} \text{Tr}\{\text{Im}G_{1,ret}(v_f k_x, v_\Delta k_y, \omega) \\ &+ \text{Im}G_{2,ret}(v_f k_y, v_\Delta k_x, \omega)\} \\ &= \frac{1}{v_\Delta v_f} \text{Tr}\left\{ \int \frac{dk'_x dk'_y}{(2\pi)^2} \text{Im}G_{1,ret}(k'_x, k'_y, \omega) \right. \\ &+ \left. \int \frac{dk'_x dk'_y}{(2\pi)^2} \text{Im}G_{2,ret}(k'_y, k'_x, \omega) \right\}.\end{aligned}\quad (3)$$

G_1 and G_2 are retarded single particle propagator for Ψ_1 and Ψ_2 . The RG equations in Eq. 2 are calculated by rescaling momentum cut-off. Since v_Δ/v_f is much smaller than 1 and flows to zero under RG, for frequency ω , the corresponding momentum scale is $\tilde{p} = \omega/v_f$. Therefore $d\ln\omega/d\ln\tilde{p} = 1 + d\ln v_f/d\ln\tilde{p}$. Now the scaling equation for $\rho(\omega)$ reads:

$$\begin{aligned}\frac{d\ln\rho}{d\ln\omega} &= \frac{d\ln\rho}{d\ln\tilde{p}} \times \frac{d\ln\omega}{d\ln\tilde{p}} = -\frac{d\ln\rho}{d\ln\Lambda} \times \left(1 - \frac{d\ln v_f}{d\ln\Lambda}\right) \\ &= \frac{(1 - C_1) - \frac{d\ln\left(\frac{1}{v_f v_\Delta}\right)}{d\ln\Lambda}}{1 - \frac{d\ln v_f}{d\ln\Lambda}} \\ &= \frac{1 - C_3 - C_2 + C_1}{1 + C_2 - C_1}.\end{aligned}\quad (4)$$

The ultraviolet cut-off of the theory is taken to be the transition temperature T_c at the critical doping of nematic transition. Although v_Δ will be renormalized to be zero in the infrared limit, the expansion of C_2 and C_3 with small v_Δ/v_f given in Ref. [26] shows that v_Δ approaches zero slowly with energy scale, therefore for the experimentally relevant energy scale, one cannot naively take the fixed point value of v_Δ . Instead, we have to integrate Eq. 4 numerically from the ultraviolet cut-off, and the result at certain frequency ω depends on the initial value of v_Δ/v_{F0} . The results of $\rho(\omega)$ with initial




FIG. 4: Plot of nodal fermion contribution to specific heat with $v_\Delta/v_{f0} = 1/5$, $v_\Delta/v_{f0} = 1/10$, $v_\Delta/v_{f0} = 1/20$ from the top to bottom. The horizontal axis is in scale of T/T_c . The curves are $y = x^{1.86}$ (red), $y = x^{1.91}$ (blue), $y = x^{1.945}$ (green), $y = x^2$ (black).

value $v_\Delta/v_{F0} = 1/5, 1/10, 1/20$ are plotted in Fig. 3, for frequency between $e^{-4} < \omega/T_c < e^{-1}$. Unlike ordinary d -wave superconductor with LDOS $\rho(\omega) \sim \omega$, in all three plots the LDOS scales with frequency as

$$\rho(\omega) \sim \omega^\alpha, \quad \alpha < 1. \quad (5)$$

This simple power law relation fits well for the frequency range $e^{-4} < \omega/T_c < e^{-1}$, and it can be checked by STM technique on samples in the quantum critical regime.

The fluctuation of nematic order parameter certainly affects the thermal dynamical quantities. The free energy $\mathcal{F} = T \ln \mathcal{Z}/V$ has scaling dimension $d + z$, the singular part of the free energy can be written as

$$\mathcal{F} \sim (\xi_\tau \xi_x \xi_y)^{-1}. \quad (6)$$

Now temperature is taken to be the infrared cut-off: $\xi_\tau \sim 1/T$, and the spatial correlation length can be estimated to be $\xi_x \sim v_x \xi_\tau$, $\xi_y \sim v_y \xi_\tau$. The anisotropic velocity of Ψ_i leads to the following contribution to the free energy:

$$\begin{aligned}\mathcal{F}_\Psi &\sim \frac{1}{v_f v_\Delta} T^3, \\ \frac{d\ln C_\Psi}{d\ln T} &= \frac{2 + \frac{d\ln\left(\frac{1}{v_f v_\Delta}\right)}{d\ln T}}{1 + C_2 - C_1} \\ &= \frac{2 + 2C_1 - C_2 - C_3}{1 + C_2 - C_1}.\end{aligned}\quad (7)$$

The ϕ contribution to the free energy and specific heat can be estimated in the same manner, although there is no velocity anisotropy for the ϕ field. The velocity of ϕ in the large N_f limit can be evaluated by calculating the one loop correction to the self-energy of ϕ . In the case of small v_Δ/v_f , the velocity of ϕ can be taken to be v_f

isotropically:

$$\begin{aligned} \mathcal{F}_\phi &\sim \frac{1}{v_f^2} T^3 \\ \frac{d \ln C_\phi}{d \ln T} &= \frac{2 + 2C_1 - 2C_2}{1 + C_2 - C_1}. \end{aligned} \quad (8)$$

The solutions of Eq. 7 with different $v_{\Delta 0}/v_{F0}$ are plotted in Fig. 4. The equations are solved for the experimentally relevant temperature range $e^{-4} < T/T_c < e^{-1}$. In general, the nodal fermions contribute more to the specific heat compared with ϕ field, because v_Δ scales stronger with temperature compared with v_f . For $e^{-4} < T/T_c < e^{-1}$, the scaling of specific heat is

$$C \sim T^\beta, \quad \beta < 2, \quad (9)$$

which distinguishes the current situation from the ordinary d -wave superconductor with nodes.

Another way to probe the density of states is the NMR relaxation rate $1/T_1$, which is related to the following Green function:

$$F(\omega) \sim \int dq_x dq_y \frac{1}{\omega} \chi''(q_x, q_y, \omega), \quad (10)$$

in the limit of $\omega \rightarrow 0$. The scaling of $1/(T_1 T)$ with temperature is the same as the scaling of $F(\omega)$ with ω , as T and ω can both serve as infrared cut-off of the theory. The momentum integrated susceptibility should involve spin density at various ‘‘slow’’ momenta. At the low energy theory of the nodal particles, following fermion bilinears are low energy spin density modes that have universal scalings:

$$\begin{aligned} \vec{q} &= (0, 0), \quad \Psi_1^\dagger \sigma^a \Psi_1 + \Psi_2^\dagger \sigma^a \Psi_2; \\ \vec{q}_{12A} &= (2Q, 0), \quad \Psi_2^\dagger \sigma^a \Psi_1; \\ -\vec{q}_{12A} &= (-2Q, 0), \quad \Psi_1^\dagger \sigma^a \Psi_2; \\ \vec{q}_{12B} &= (0, 2Q), \quad \Psi_1^t \tau^y \sigma^y \sigma^a \Psi_2; \\ -\vec{q}_{12B} &= (0, -2Q), \quad \Psi_2^t \tau^y \sigma^a \sigma^y \Psi_1^*; \\ \vec{q}_{11} &= (2Q, 2Q), \quad \Psi_1^t \tau^y \sigma^y \sigma^a \Psi_1; \\ -\vec{q}_{11} &= (-2Q, -2Q), \quad \Psi_1^t \tau^y \sigma^a \sigma^y \Psi_1^*; \\ \vec{q}_{22} &= (-2Q, 2Q), \quad \Psi_2^t \tau^y \sigma^y \sigma^a \Psi_2; \\ -\vec{q}_{22} &= (2Q, -2Q), \quad \Psi_2^t \tau^y \sigma^a \sigma^y \Psi_2^* \end{aligned} \quad (11)$$

σ^a are three spin Pauli matrices. To evaluate $F(\omega)$ we need to calculate the correlation of all the fermion bilinears above. The susceptibility χ gains fermion self-energy correction as in Fig. 2a as well as vertex corrections Fig 2c and Fig. 2d for vertices $\Psi_i^\dagger T_A \Psi_j$ and $\Psi_i^t T_A \Psi_j$ respectively. For a general fermion bilinear with flavor matrix




FIG. 5: Plot of $1/(T_1 T)$ with T from contribution of spin density at $\vec{q} = (0, 0)$, with $v_{\Delta 0}/v_{f0} = 1/5$, $v_{\Delta 0}/v_{f0} = 1/10$, $v_{\Delta 0}/v_{f0} = 1/20$ from the top to bottom. The horizontal axis is in scale of T/T_c . The curves are $y = x^{1.74}$ (red), $y = x^{1.83}$ (blue), $y = x^{1.89}$ (green), $y = x^2$ (black).

T_A , the vertex correction RG equation is conventionally written as

$$\frac{dT_A}{d \ln \Lambda} = C_A T_A. \quad (12)$$

C_A is a function of v_Δ/v_f . The details of calculations are given in the appendix, the results are

$$\begin{aligned} C_0 &= -C_1, \quad C_{\tau^x} = C_4 = -C_3, \\ C_{\tau^y} &= C_3 - C_1 - C_2, \quad C_{\tau^z} = -C_2 \\ C_{12A} &= C_{12B} = -0.3486 \frac{v_\Delta/v_f}{N_f} + \dots \\ C_{11} &= C_{22} = -C_{\tau^y}. \end{aligned} \quad (13)$$

C_0 is the vertex correction for fermion bilinear $\Psi_i^\dagger \Psi_i$ and $\Psi_i^\dagger \sigma^a \Psi_i$, because spin is a good quantum number spin Pauli matrices do not change the vertex correction. C_{12A} and C_{12B} are vertex corrections to $\Psi_1^\dagger \sigma^a \Psi_2$ and $\Psi_1^t \tau^y \sigma^y \sigma^a \Psi_2$, C_{11} and C_{22} are vertex corrections to $\Psi_1^t \tau^y \sigma^y \sigma^a \Psi_1$ and $\Psi_2^t \tau^y \sigma^y \sigma^a \Psi_2$ respectively.

After taking into account of both self-energy and vertex corrections, the scaling equation of the relaxation rate reads:

$$\begin{aligned} \frac{d \ln F(\omega)}{d \ln \omega} &= \frac{2 - 2C_1 - 2C_A + 2 \frac{d \ln \left(\frac{1}{v_f v_\Delta} \right)}{d \ln \omega}}{1 + C_2 - C_1} \\ &= \frac{2 - 2C_A - 2C_2 - 2C_3 + 2C_1}{1 + C_2 - C_1}. \end{aligned} \quad (14)$$

Equation (14) leads to the following scaling equation for $1/(T_1 T)$ with temperature:

$$\frac{d \ln 1/(T_1 T)}{d \ln T} = \frac{2 - 2C_A - 2C_2 - 2C_3 + 2C_1}{1 + C_2 - C_1}. \quad (15)$$




FIG. 6: Plot of $1/(T_1T)$ with T from contribution of spin density at $\vec{q} = (2Q, 0)$ and $\vec{q} = (0, 2Q)$, with $v_{\Delta 0}/v_{f0} = 1/5$, $v_{\Delta 0}/v_{f0} = 1/10$, $v_{\Delta 0}/v_{f0} = 1/20$ from the top to bottom. The horizontal axis is in scale of T/T_c . The curves are $y = x^{1.88}$ (red), $y = x^{1.9}$ (blue), $y = x^{1.93}$ (green), $y = x^2$ (black).




FIG. 7: Plot of $1/(T_1T)$ with T from contribution of spin density at $\vec{q} = (2Q, 2Q)$ and $\vec{q} = (2Q, -2Q)$, with $v_{\Delta 0}/v_{f0} = 1/5$, $v_{\Delta 0}/v_{f0} = 1/10$, $v_{\Delta 0}/v_{f0} = 1/20$ from the bottom to the top. The horizontal axis is in scale of T/T_c . The curves are $y = x^{2.08}$ (red), $y = x^{2.05}$ (blue), $y = x^{2.025}$ (green), $y = x^2$ (black).

The contribution from different fermion bilinear components listed in Eq. 11 should be solved individually, and the solutions of each component are plotted in Fig. 5, 6 and 7. One can see that at small frequency the spin density at $(0, 0)$ makes the most substantial contribution, which makes the scaling of $1/(T_1T)$ differ from the ordinary d -wave superconductor:

$$\frac{1}{T_1T} \sim T^\gamma, \quad \gamma < 2. \quad (16)$$

The superconductivity can be fully suppressed by strong enough magnetic field. Since the nematic quantum critical point is most likely in the underdoped regime with low T_c , it is possible to apply strong enough inplane magnetic field $H > H_{c2}$ in experiments. With supercon-

ductivity fully suppressed by inplane magnetic field, the universality class of the nematic transition becomes the $z = 3$ theory described in Ref. [21]. The $z = 3$ quantum critical point leads to a large number of low energy excitations, which will contribute to thermal dynamics and transport. The standard mean field theory leads to following results at low temperature [21]:

$$C \sim T^{2/3}, \quad \rho \sim T^{4/3}. \quad (17)$$

These results will finally crossover to $d = 3$ scalings close enough to the critical point:

$$C \sim T, \quad \rho \sim T^{5/3}. \quad (18)$$

Similar $z = 3$ nematic quantum critical point was discussed in Ref. [29].

In a summary, in this work we computed the scaling of physical quantities close to a nematic quantum critical point in a d -wave superconducting phase of the newly discovered material $\text{SmFeAsO}_{1-x}\text{F}_x$, motivated by recent experiments on the polycrystal sample. For the experimentally relevant energy range, the scaling of LDOS, specific heat, NMR relaxation rate all deviates from an ordinary d -wave superconductor. These results also apply, essentially unchanged, to the cuprate superconductors [25, 26]. This research is supported by the NSF under grant DMR-0537077.

APPENDIX

In this section we shall calculate the vertex corrections to the fermion bilinears listed in Eq. 11. For a general vertex $\Psi_i^\dagger T_A \Psi_j$, the vertex correction is calculated according to Feynman diagram in Fig. 2c:

$$\begin{aligned} \mathcal{T}_A = T_A + \frac{1}{N_f} \int \frac{d^2p}{(2\pi)^2} \frac{d\Omega}{2\pi} [\tau^x G_i(\Omega, p) \\ \times T_A G_j(\Omega, p) \tau^x] \frac{1}{\Gamma_2(\Omega, p)}. \end{aligned} \quad (19)$$

For a general vertex $\Psi_i^t T_A \Psi_j$, the vertex correction is calculated according to Feynman diagram in Fig. 2d:

$$\begin{aligned} \mathcal{T}_A = T_A + \frac{1}{N_f} \int \frac{d^2p}{(2\pi)^2} \frac{d\Omega}{2\pi} (-1) [\tau^x G_i(\Omega, p) \\ \times T_A G_j(\Omega, p) \tau^x] \frac{1}{\Gamma_2(\Omega, p)}. \end{aligned} \quad (20)$$

$\Gamma_2(\Omega, p)$ is the self-energy of ϕ from integrating out fermions:

$$\begin{aligned} \Gamma_2(\omega, p) = \Pi_2(k_x, k_y, \omega) + \Pi_2(k_y, k_x, \omega), \\ \Pi_2(k_x, k_y, \omega) = \frac{1}{16v_f v_\Delta} \frac{\omega^2 + v_f^2 k_x^2}{(\omega^2 + v_f^2 k_x^2 + v_\Delta^2 k_y^2)^{1/2}}. \end{aligned} \quad (21)$$

Π_2 is reminiscent of the vacuum polarization of the 2+1d QED, with a gauge invariant form $\Pi_{\mu\nu}(p) \sim p(\delta_{\mu\nu} - p_\mu p_\nu/p^2)$, and $\Pi_2 = \Pi_{xx}(p)$.

The RG equation is defined as the change of parameters after rescaling the cut-off. The cut-off is introduced in a smooth function $\mathcal{K}(p^2/\Lambda^2)$, with $\mathcal{K}(0) = 1$ and falls rapidly when $y \rightarrow 1$. Now following the notation in Ref. [26], we change the momentum space integral to cylindrical coordinates $p_\mu = y\Lambda(v_f x, \cos\theta, \sin\theta)$, and the RG equations in Eq. 2 and Eq. 12 are obtained when the cut-off is reduced from Λ to $\Lambda - d\Lambda$, with

$$\begin{aligned}
C_1 &= -C_0 = \frac{2(v_\Delta/v_f)}{\pi^3 N_f} \int_{-\infty}^{+\infty} dx \int_0^{2\pi} d\theta \\
&\times \frac{(x^2 - \cos^2\theta - (v_\Delta/v_f)^2 \sin^2\theta)}{(x^2 + \cos^2\theta + (v_\Delta/v_f)^2 \sin^2\theta)^2} \mathcal{G}(x, \theta); \\
C_2 &= -C_{\tau z} = \frac{2(v_\Delta/v_f)}{\pi^3 N_f} \int_{-\infty}^{+\infty} dx \int_0^{2\pi} d\theta \\
&\times \frac{(-x^2 + \cos^2\theta - (v_\Delta/v_f)^2 \sin^2\theta)}{(x^2 + \cos^2\theta + (v_\Delta/v_f)^2 \sin^2\theta)^2} \mathcal{G}(x, \theta); \\
C_3 &= -C_{\tau x} = \frac{2(v_\Delta/v_f)}{\pi^3 N_f} \int_{-\infty}^{+\infty} dx \int_0^{2\pi} d\theta \\
&\times \frac{(x^2 + \cos^2\theta - (v_\Delta/v_f)^2 \sin^2\theta)}{(x^2 + \cos^2\theta + (v_\Delta/v_f)^2 \sin^2\theta)^2} \mathcal{G}(x, \theta); \\
C_{\tau y} &= -C_{11} = -C_{22} = \frac{2(v_\Delta/v_f)}{\pi^3 N_f} \int_{-\infty}^{+\infty} dx \int_0^{2\pi} d\theta \\
&\times \frac{(x^2 + \cos^2\theta + (v_\Delta/v_f)^2 \sin^2\theta)}{(x^2 + \cos^2\theta + (v_\Delta/v_f)^2 \sin^2\theta)^2} \mathcal{G}(x, \theta) \\
&= C_3 - C_1 - C_2; \\
C_{12A} &= C_{12B} = \frac{2(v_\Delta/v_f)}{\pi^3 N_f} \int_{-\infty}^{+\infty} dx \int_0^{2\pi} d\theta \\
&\times \frac{1}{x^2 + \cos^2\theta + (v_\Delta/v_f)^2 \sin^2\theta} \\
&\times \frac{1}{x^2 + (v_\Delta/v_f)^2 \cos^2\theta + \sin^2\theta} \mathcal{G}(x, \theta); \\
\mathcal{G}^{-1}(x, \theta) &= \frac{x^2 + \sin^2\theta}{\sqrt{x^2 + (v_\Delta/v_f)^2 \cos^2\theta + \sin^2\theta}} \\
&+ \frac{x^2 + \cos^2\theta}{\sqrt{x^2 + (v_\Delta/v_f)^2 \sin^2\theta + \cos^2\theta}}. \quad (22)
\end{aligned}$$

It was noted in Ref. [26] that function C_3 has a rather singular form in the small v_Δ/v_f limit:

$$C_3 = \frac{8}{\pi^2} [\ln(v_f/v_\Delta) - 0.9601] \frac{v_\Delta/v_f}{N_f} + \dots \quad (23)$$

Plugging this function into Eq. 2, one can see that v_Δ/v_f approaches zero a little faster than the ordinary marginally irrelevant operators, but for the experimentally relevant energy scale, v_Δ/v_f still flows slowly. Therefore all the plots in our paper, though integrated from a complicated equation, can be fit with a simple power law.

-
- [1] Linjun Li, Yuke Li, Zhi Ren, Xiao Lin, Yongkang Luo, Zengwei Zhu, Mi He, Xiangfan Xu, Guanghan Cao, Zhu'an Xu, arXiv:0806.1675 (2008).
 - [2] Zhi-An Ren, Jie Yang, Wei Lu, Wei Yi, Guang-Can Che, Xiao-Li Dong, Li-Ling Sun, Zhong-Xian Zhao, arXiv:0803.4283 (2008).
 - [3] Zhi-An Ren, Guang-Can Che, Xiao-Li Dong, Jie Yang, Wei Lu, Wei Yi, Xiao-Li Shen, Zheng-Cai Li, Li-Ling Sun, Fang Zhou, Zhong-Xian Zhao, EuroPhys. Lett. **83**, 17002 (2008).
 - [4] X. H. Chen, T. Wu, G. Wu, R. H. Liu, H. Chen, D. F. Fang, Nature **453**, 761 (2008).
 - [5] Linjun Li, Yuke Li, Zhi Ren, Xiao Lin, Yongkang Luo, Zengwei Zhu, Mi He, Xiangfan Xu, Guanghan Cao, Zhu'an Xu, arXiv:0806.1675 (2008).
 - [6] Clarina de la Cruz, Q. Huang, J. W. Lynn, Jiying Li, W. Ratcliff II, J. L. Zarestky, H. A. Mook, G. F. Chen, J. L. Luo, N. L. Wang, Pengcheng Dai, Nature **453**, 899 (2008).
 - [7] Jun Zhao, Q. Huang, Clarina de la Cruz, Shiliang Li, J. W. Lynn, Y. Chen, M. A. Green, G. F. Chen, G. Li, Z. Li, J. L. Luo, N. L. Wang, Pengcheng Dai, arXiv:0806.2585 (2008).
 - [8] Marianne Rotter, Marcus Tegel, Dirk Johrendt, arXiv:0805.4630 (2008).
 - [9] N. Ni, S. L. Bud'ko, A. Kreyssig, S. Nandi, G. E. Rustan, A. I. Goldman, S. Gupta, J. D. Corbett, A. Kracher, P. C. Canfield, arXiv:0806.1874 (2008).
 - [10] Q. Huang, Y. Qiu, Wei Bao, J.W. Lynn, M.A. Green, Y. Chen, T. Wu, G. Wu, X.H. Chen, arXiv:0806.2776 (2008).
 - [11] Marianne Rotter, Marcus Tegel, Inga Schellenberg, Wilfried Hermes, Rainer Pötz, Dirk Johrendt, arXiv:0805.4021 (2008).
 - [12] C. Krellner, N. Caroca-Canales, A. Jesche, H. Rosner, A. Ormeci, C. Geibel, arXiv:0806.1043 (2008).
 - [13] J.-Q. Yan, A. Kreyssig, S. Nandi, N. Ni, S. L. Bud'ko, A. Kracher, R. J. McQueeney, R. W. McCallum, T. A. Lograsso, A. I. Goldman, P. C. Canfield, arXiv:0806.2711 (2008).
 - [14] Jun Zhao, W. Ratcliff II, J. W. Lynn, G. F. Chen, J. L. Luo, N. L. Wang, Jiangping Hu, Pengcheng Dai, arXiv:0807.1077 (2008).
 - [15] Milton S. Torikachvili, Sergey L. Bud'ko, Ni Ni, and Paul C. Canfield, arXiv:0807.0616 (2008).
 - [16] A.I. Goldman, D.N. Argyriou, B. Ouladdiaf, T. Chatterji, A. Kreyssig, S. Nandi, N. Ni, S. L. Bud'ko, P.C. Canfield, R. J. McQueeney, arXiv:0807.1525 (2008).
 - [17] R. H. Liu, G. Wu, T. Wu, D. F. Fang, H. Chen, S. Y. Li, K. Liu, Y. L. Xie, X. F. Wang, R. L. Yang, L. Ding, C. He, D. L. Feng, X. H. Chen, arXiv:0804.2105 (2008).

- [18] Serena Margadonna, Yasuhiro Takabayashi, Martin T. McDonald, Michela Brunelli, G. Wu, R. H. Liu, X. H. Chen, Kosmas Prassides, arXiv:0806.3962 (2008).
- [19] Y. Qiu, Wei Bao, Q. Huang, J.W. Lynn, T. Yildirim, J. Simmons, Y.C. Gasparovic, J. Li, M. Green, T. Wu, G. Wu, X.H. Chen, arXiv:0806.2195 (2008).
- [20] Ying Chen, J. W. Lynn, J. Li, G. Li, G. F. Chen, J. L. Luo, N. L. Wang, Pengcheng Dai, C. dela Cruz, H. A. Mook, arXiv:0807.0662 (2008).
- [21] Cenke Xu, Markus Mueller, Subir Sachdev, Phys. Rev. B **78**, 020501(R) (2008).
- [22] Chen Fang, Hong Yao, Wei-Feng Tsai, JiangPing Hu, Steven A. Kivelson, Phys. Rev. B **77** 224509 (2008).
- [23] M. A. McGuire, A. D. Christianson, A. S. Sefat, B. C. Sales, M. D. Lumsden, R. Jin, E. A. Payzant, D. Mandrus, Y. Luan, V. Keppens, V. Varadarajan, J. W. Brill, R. P. Hermann, M. T. Sougrati, F. Grandjean, G. J. Long, arXiv:0806.3878 (2008).
- [24] Marcus Tegel, Marianne Rotter, Veronika Weiss, Falko M. Schappacher, Rainer Poettgen, Dirk Johrendt, arXiv:0806.4782 (2008).
- [25] Eun-Ah Kim, Michael J. Lawler, Paul Oreto, Subir Sachdev, Eduardo Fradkin, Steven A. Kivelson, Phys. Rev. B **77**, 184514 (2008).
- [26] Yejin Huh, Subir Sachdev, arXiv:0806.0002 (2008).
- [27] Oded Millo, Itay Asulin, Ofer Yuli, Israel Felner, Zhi-An Ren, Xiao-Li Shen, Guang-Can Che, Zhong-Xian Zhao, arXiv:0807.0359 (2008).
- [28] Yonglei Wang, Lei Shan, Lei Fang, Peng Cheng, Cong Ren, Hai-Hu Wen, arXiv:0806.1986 (2008).
- [29] Vadim Oganesyan, Steven Kivelson, Eduardo Fradkin, Phys. Rev. B **64**, 195109 (2001).