all immunotherapy drugs
Patients with matching 1:N
Patients who switched to
Vedolizumab (switch cohort) or
who did not switch (matching N)
to Vedolizumab (control cohort)
P-value (joint matching)
P-value (post matching)
all autoimmune diseases
Acute rheumatic heart disease
Alopecia
Ankylosing spondylitis
Arthropathy
Autoimmune disease NEC
Autoimmune hemolytic anemias
Behcet's syndrome
Biliary cirrhosis
Bullous dermatoses
Celiac disease
Chronic hepatitis
Dermatomyositis
diabetes mellitus
Diffuse diseases of connective tissue
Diseases of hair and hair follicles
Encephalitis
Endocarditis
Giant cell arteritis
Glomerulonephritis
Granulomatosis with polyangiitis
Graves' disease
Hypothyroidism NOS
IBD
Inflammatory and toxic neuropathy
Multiple sclerosis
Myalgia and myositis NOS
Myeloid disorders
Nephritis
Other diseases of central nervous system
Other diseases of lung
Other disorders of the immune mechanism
Other endocrine disorders
Other immunological findings
Other peripheral diffuse diseases of connective tissue
Pemphigus and pemphigoid
Peripheral autonomic neuropathy
Polyarteritis nodosa and allied conditions
Polymyalgia Rheumatica
Polymyositis
Primary thrombocytopenia
Psoriasis vulgaris
Psoriatic arthropathy
RA
Raynaud's syndrome
Reiter's disease
Rheumatic fever / chorea
Sarcoidosis
Sicca syndrome
Systemic lupus erythematosus
Systemic sclerosis
Thrombotic microangiopathy
Thyroiditis
Thromboses with or without polar
Toxic erythema
Vitiligo
Wegener's granulomatosis
post matching
pre matching
Num. at risk