

1
2 Addis, D.R. & Schacter, D.L. (2012). *Frontiers in Human Neuroscience*, 5, 173. doi:
3 10.3389/fnhum.2011.00173.

4
5 **The hippocampus and imagining the future: Where do we stand?**

6
7 Donna Rose Addis^{1*} and Daniel L. Schacter²

8
9
10 ¹ARC Centre of Excellence in Cognition and its Disorders, Centre for Brain Research and
11 Department of Psychology, The University of Auckland, New Zealand

12 ²Department of Psychology, Harvard University, MA, USA

13
14 *Corresponding author:

15 Dr. Donna Rose Addis
16 Department of Psychology, The University of Auckland
17 Private Bag 92019,
18 Auckland 1142,
19 New Zealand
20 d.addis@auckland.ac.nz

21
22 Running title: Hippocampus and imagining the future

23 Keywords: Hippocampus, future, imagination, simulation, episodic, autobiographical

24 **Abstract**

25

26 Recent neuroimaging work has demonstrated that the hippocampus is engaged when
27 imagining the future, in some cases more than when remembering the past. It is possible that
28 this hippocampal activation reflects recombining details into coherent scenarios and/or the
29 encoding of these scenarios into memory for later use. However, inconsistent findings have
30 emerged from recent studies of future simulation in patients with memory loss and
31 hippocampal damage. Thus, it remains an open question as to whether the hippocampus is
32 necessary for future simulation. In this review, we consider the findings from patient studies
33 and the neuroimaging literature with respect to a new framework that highlights three
34 component processes of simulation: accessing episodic details, recombining details, and
35 encoding simulations. We attempt to reconcile these discrepancies between neuroimaging
36 and patient studies by suggesting that different component processes of future simulation may
37 be differentially affected by hippocampal damage.

38

39 **1. Introduction**

40

41 In daily life, particularly during the unoccupied moments, we often revert to our inner mental
42 world and engage with our aspects of our lives outside of the present. Mentally projecting
43 ourselves back into the past or forwards into the future can take make forms – a cursory
44 thought, a vague image, or a vivid and consuming scenario. There has been increasing
45 interest in understanding the ways in which remembering and future thinking are similar or
46 different, both in terms of cognitive and neural processes, and whether such characteristics
47 are evident for various forms of past and future thinking (for recent reviews, see Schacter,
48 Addis, & Buckner, 2008; Szpunar, 2010). These studies have been informed by a closely
49 related line of neuroimaging research showing that when people are consumed by various
50 forms of thoughts and images, these internally-directed cognitive activities are accompanied
51 by a characteristic pattern of neural activity - known as the default network (Buckner,
52 Andrews-Hanna, & Schacter, 2008; Spreng, Mar, & Kim, 2009).

53

54 This network, which includes many regions traditionally associated with memory, such as the
55 hippocampus, is also up-regulated by tasks that specifically require a focus on remembering
56 and imagining personal experiences (Buckner & Carroll, 2007; Schacter, Addis, & Buckner,
57 2007; Spreng et al., 2009). Motivated by findings that remembering and imagining engage
58 the same ‘common core network’, we advanced the *constructive episodic simulation*
59 *hypothesis*, which holds that the common neural activity for past and future reflects a reliance
60 on memory to provide the details comprising both remembered and imagined event
61 representations (Schacter & Addis, 2007). In that theory, as well as in this review, we focus
62 on a particularly vivid form of future thinking: the imaginative construction or simulation of
63 scenarios that might occur in one’s future. We hypothesized that the flexible use of episodic
64 details from memory during imaginative simulations of the future can help to understand
65 constructive aspects of memory, such as its susceptibility to distortion (see also Schacter,
66 Guerin, & St. Jacques, 2011). Like autobiographical memories of past experiences, these
67 simulations are considered “episodic” in nature because they represent the self engaging in a
68 specific event in a particular spatiotemporal context. And although the emphasis here is
69 primarily on simulations located in the imagined future, primarily because of the adaptive
70 value of such simulations for maximizing future success (Ingvar, 1985; Schacter & Addis,
71 2007; Suddendorf & Corballis, 1997, 2007; Szpunar, 2010), simulations can also focus on
72 present or past events; indeed, we have argued that many of the same processes discussed
73 here are likely also applicable under those conditions (Addis, Pan, Vu, Laiser, & Schacter,
74 2009).

75

76 One of the more compelling and even unexpected findings from research on the neural
77 underpinnings of episodic simulations is that the hippocampus, a region traditionally thought
78 of as a “memory region”, can be engaged to a greater degree when imagining than
79 remembering (e.g., Addis, Wong, & Schacter, 2007; for reviews, see Buckner, 2010; Schacter
80 & Addis, 2009). Such findings raise the question of what is unique about episodic simulation
81 or future thinking that recruits the hippocampus. In very general terms, it would appear that
82 more intensive processing is required when imagining future events relative to retrieving past
83 events, because the former requires construction of a novel event, whereas the latter involves
84 retrieval of an already established event. However, determining what specific component
85 processes underlie this ‘more intensive processing’, and which such processes rely on the
86 hippocampus, is necessary to better understand this future>past effect. A number of candidate

87 cognitive processes exist. Although both remembering and imagining typically involve the
 88 reactivation of memories and episodic details comprising these memories, only imagining
 89 requires the additional step of recombining such details into a new arrangement – the
 90 imagined scenario. It is plausible that this recombination process would engage the
 91 hippocampus, given its role in relational memory processes that link together disparate bits of
 92 information (Eichenbaum, 2001). Also, if these newly constructed scenarios are ever to be
 93 accessed in future, they need to be encoded and stored in memory (Ingvar, 1985). In this
 94 review, we will discuss the conditions under which a hippocampal future>past effect
 95 emerges, and also consider recent work investigating whether hippocampal activation during
 96 future thinking reflects access to episodic details, recombining these details to construct
 97 specific scenarios, and/or the encoding of these scenarios into memory.

98
 99 A related line of enquiry is to determine not only whether the hippocampus is active during
 100 future simulation but whether it makes a critical and necessary contribution. While it has
 101 been long established that a functioning hippocampus is necessary for the retrieval of detailed
 102 autobiographical memories (for a review, see Moscovitch et al., 2005), it is less clear whether
 103 this is the case for future simulation (see Table 1 for a summary of patient cases discussed
 104 herein). While some patients with hippocampal damage and impaired episodic memory also
 105 exhibit difficulties in imagining detailed and coherent future events (Andelman, Hoofien,
 106 Goldberg, Aizenstein, & Neufeld, 2010; Hassabis, Kumaran, Vann, & Maguire, 2007; Race,
 107 Keane, & Verfaellie, 2011), other studies do not report imagination deficits in such patients.
 108 Spared simulation abilities in the context of hippocampal damage and memory loss have been
 109 reported in an adult developmental amnesic patient (Maguire, Vargha-Khadem, & Hassabis,
 110 2010), a group of developmental amnesic school-aged children (Cooper, Vargha-Khadem,
 111 Gadian, & Maguire, 2011; see also, Hurley, Maguire, & Vargha-Khadem, in press), and a
 112 group of adult patients with bilateral hippocampal damage (Squire et al., 2010).

113
 114 --Insert Table 1 about here--
 115

116 Such findings imply that a fully intact hippocampus may not be required for future
 117 simulation. However, the inconsistent results yielded from these studies raise a number of
 118 important questions. Does the temporal extent of amnesia influence the degree to which
 119 imagined scenarios can be constructed? Does the age of onset of hippocampal damage affect
 120 the degree of impairment? Does the location of the damage within the hippocampus influence
 121 the pattern of spared and impaired abilities? Can residual hippocampal tissue support future
 122 simulation? Are particular simulation tasks better able to detect deficits? In considering the
 123 findings from patient studies in conjunction with those from neuroimaging literature, we will
 124 attempt to reconcile these discrepant results by suggesting that different component processes
 125 of future simulation may be differentially affected by hippocampal damage: although the
 126 processes of accessing and recombining details to construct and encode a future event are
 127 inherently related processes in healthy individuals, it is possible that in the damaged brain
 128 these processes are, to some extent, dissociable.

129

130 **2. Access to memory details: the episodic fodder for future simulations**

131

132 In recent years, neuroimaging has provided evidence to suggest that imagining the future
 133 relies on much of the same neural machinery as remembering the past. One hypothesis that
 134 such findings motivate is that memories must be reactivated in order to extract the
 135 information needed to ‘flesh out’ detailed simulations. Indeed, if simulations involve the

136 projection of the self in time beyond the present (Buckner & Carroll, 2007) and are to be
137 meaningful for that individual, then personally-relevant episodic details from memory are
138 needed. Such elements would include the major components of an episode, including the
139 people, places and objects previously encountered by the individual. In their scene
140 construction hypothesis, Hassabis and Maguire (2007) argue that spatial information is
141 particularly important. A spatial framework provides a platform upon which to build the
142 scenario, and without this, an imagined event would likely lack a sense of coherence.

143

144 Although common hippocampal activity for past and future events is suggestive of access to
145 mnemonic information during both tasks, it is not conclusive. Addis and Schacter (2008)
146 examined whether hippocampal responses during remembering and imagining were
147 modulated by subjective ratings of the detail comprising these events. Activity in the
148 posterior hippocampus correlated with detail ratings for both past and future events,
149 consistent with the idea that both tasks require access to episodic details. Moreover, Weiler,
150 Suchan and Daum (2010b) found activity in the posterior hippocampus was associated with
151 both past and future events, though the responses had differing timecourses. Nevertheless, the
152 location of this neural response dovetails with studies implicating the posterior hippocampus
153 in retrieval as opposed to encoding (Lepage, Habib, & Tulving, 1998; Prince, Daselaar, &
154 Cabeza, 2005; Schacter & Wagner, 1999), in the reinstatement of previous conditions
155 (Giovanello, Schnyer, & Verfaellie, 2009; Preston, Shrager, Dudukovic, & Gabrieli, 2004),
156 and in the amount of detail comprising autobiographical memory (Addis, Moscovitch,
157 Crawley, & McAndrews, 2004).

158

159 However, the most convincing evidence that access to episodic details may be *necessary* for
160 future simulations comes from studies of patients with memory loss (see Table 1). One of the
161 early observations of a link between past and future thinking came from Tulving (1985). In a
162 discussion of K.C., a patient with dense autobiographical amnesia resulting from a head
163 injury, it was also noted that K.C. exhibited difficulties in imagining specific episodes in his
164 personal future. Similarly, amnesic patient D.B., who sustained brain damage as a result of
165 cardiac arrest and anoxia, cannot remember or imagine personal events (Klein, Loftus, &
166 Kihlstrom, 2002). In both cases, the neuroanatomical damage is not restricted to the
167 hippocampus (patient KC has damage in and beyond the hippocampus, including extensive
168 prefrontal damage, Rosenbaum et al., 2005; no neuroanatomical findings have been reported
169 for patient D.B.). Nevertheless, these reports raised the possibility that there is a link between
170 remembering and imagining – that being able to access details from episodic memory may be
171 an important and perhaps necessary condition of the successful construction of episodic
172 simulations.

173

174 Similar results have been reported in patients with damage reported to be limited to the
175 hippocampus. Hassabis et al. (2007) found that four out of five patients with hippocampal
176 amnesia could not construct imaginary scenarios of everyday scenes: their constructions
177 contained significantly less content than those of controls, and the details that were generated
178 were not well integrated. Although the authors also found that providing patients with details
179 did not improve their performance, the provided information was semantic in nature and
180 therefore may not have been sufficient to support imaginings that have an episodic basis. One
181 critical issue is whether these patients have damage circumscribed to the hippocampus.
182 Although Maguire and Hassabis (2011) state these patients were “specifically selected” for
183 damaged restricted to the hippocampus, Squire and colleagues (Squire, McDuff, & Frascino,
184 2011) disagree with this assessment. They argue that aspects of the clinical profiles of these

185 patients (e.g., generalized atrophy, seizures, personality change) suggest the presence of
186 damage outside of the hippocampus. They also note that the one patient in the Hassabis et al.
187 (2007) study who did not exhibit imagination deficits had a different etiology
188 (meningoencephalitis and recurrent meningitis, versus limbic encephalitis in the four other
189 patients), as well as residual hippocampal tissue and function (Hassabis et al., 2007).

190

191 Race and colleagues (2011) examined the ability to remember and imagine in a group of
192 eight amnesic patients with medial temporal damage. This study is important for two reasons.
193 First, the paradigm included a condition in which participants were required to construct
194 narratives when the details did not have to be retrieved from memory but were presented as
195 pictures (also see Gaesser, Sacchetti, Addis, & Schacter, 2011). When completing the past
196 and future tasks, amnesic patients generated significantly fewer episodic details than did
197 controls, and the number of episodic details for past and future narratives was correlated.
198 Critically, hippocampal damage did not disrupt the ability to construct a narrative in the
199 picture condition, where access to episodic memory was not required. Moreover,
200 performance on the picture narrative task was not correlated with performance on the future
201 task. Second, although the etiology and extent of damage varied across the eight patients,
202 there was one patient in whom damage was confirmed as being limited to the hippocampus.
203 Importantly, the performance of this patient mirrored that of the other patients who had some
204 degree of extra-hippocampal temporal damage, suggesting that damage to the hippocampus
205 alone is sufficient to disrupt future simulation. Together, the observations from this study
206 further support the notion that in the context of hippocampal damage, it is an inability to
207 access details in episodic memory, and not more general deficits in narrative ability, that
208 underlies deficient episodic simulation performance.

209

210 While studies of amnesia give insight into the ability to simulate when there is little, or no,
211 access to episodic details, studies of aging -- where deficits in accessing past events are
212 present but comparatively milder -- have also provided relevant evidence. In a series of
213 studies, we have examined the ability to remember and imagine in healthy and also in
214 pathological aging (i.e., patients in the early stages of Alzheimer's disease), in which
215 autobiographical memory is typically affected (Levine, Svoboda, Hay, Winocur, &
216 Moscovitch, 2002), and hippocampal atrophy and dysfunction are also evident (Hedden &
217 Gabrieli, 2004). In these studies, we had participants generate memories of past events and
218 simulations of future events in response to word cues and found that the number of episodic
219 details comprising events in older or demented adults was reduced relative to appropriate
220 control groups (for a review, see Schacter, Gaesser, & Addis, 2010). Moreover, the number
221 of episodic details for past events is strongly correlated with the number of details comprising
222 future events. These correlations are consistently evident across old and young (Addis,
223 Musicaro, Pan, & Schacter, 2010; Addis, Wong, & Schacter, 2008), and across demented and
224 healthy older adults (Addis, Sacchetti, Ally, Budson, & Schacter, 2009), and exist even when
225 controlling other factors that may more generally influence the detail of narratives, such as
226 cognitive decline and verbal fluency (Addis, Sacchetti, et al., 2009). The deficits in episodic
227 remembering and imagining that we have documented in older adults do also extend to a
228 picture description task that does not require episodic memory (Gaesser et al., 2011).
229 Nonetheless, we also found that the age deficits in remembering and imagining were still
230 observed after controlling for general narrative abilities, as measured by this picture
231 description task. Neuroimaging evidence suggests that the reduction in episodic detail when
232 older adults describe past and future events may be related to dysfunction in the regions
233 supporting episodic detail, including the hippocampus (Addis, Roberts, & Schacter, 2011).

234

235 What is to be made, then, of patients with memory loss who can still imagine the future?
236 Such findings appear to speak against the idea that access to memories is a critical precursor
237 to future simulation. As noted earlier, Squire et al. (2010) reported that a group of patients
238 with damage to the hippocampus showed an intact ability to create detailed imaginary future
239 events. However, although these patients have hippocampal damage, it is notable that their
240 degree of retrograde amnesia is minimal: these patients can retrieve events from the remote
241 past, and only exhibit a mild (and non-significant) deficit for retrieving memories from the
242 recent past. Thus, the results of this study could also be interpreted as supporting the notion
243 that access to the past – even in the context of hippocampal damage – can provide a basis for
244 imagining the future.

245

246 However, there are reported cases of hippocampal damage that has differentially affected
247 remembering but not imagining. For instance, Maguire and colleagues reported that
248 developmentally amnesic patients who sustained hippocampal damage early in life can
249 construct imaginary scenarios (Maguire et al., 2010; Hurley et al., in press; but see, Kwan,
250 Carson, Addis, & Rosenbaum, 2010). Moreover, as noted earlier, one of the patients from the
251 Hassabis et al. (2007) study could also complete their scene construction task. Interestingly,
252 some of these patients have been noted to have residual hippocampal tissue that appears to be
253 functional, in that it is activated during memory tasks (Maguire et al., 2010), although such
254 activation has not yet been shown during future simulation. These researchers also report
255 normal imagination abilities in a group of children with hippocampal damage and amnesia
256 (Cooper et al., 2011), further suggesting that the time of onset of the amnesia may be an
257 important consideration. It is possible that with early damage, these patients develop other
258 strategies or rely either on residual episodic memories or detailed semantic information to
259 construct scenarios (Cooper et al., 2011).

260

261 It is also notable that these findings have emerged using the scene construction task. Hassabis
262 et al. (2007) mention that this task was designed to “increase the dependence of constructions
263 on generalized semantic memory representations”. On each trial, a sentence cue (e.g.,
264 “Imagine you are lying on a white sandy beach”) is provided to take participants into a
265 generic scene; it is very likely that this scene can then be fleshed out with semantic detail.
266 Thus, it is possible that these patients are able to complete this particular imagination task
267 using detailed yet semantic representations of how certain scenes or episodes unfold, rather
268 than extracting information from their own experiences. However, when the task requires
269 creation of a specific and novel episode, similar patients (e.g., with developmental amnesia)
270 show simulation deficits – particularly in the amount of episodic detail generated (Kwan et
271 al., 2010). Although amnesics may generate fewer episodic details relative to controls, they
272 sometimes show little or no reduction in the number of semantic details comprising their
273 event narratives (Race et al., 2011). It has also been shown in other studies that patients with
274 episodic, but not semantic, memory deficits can successfully complete future thinking tasks
275 that are based primarily on general knowledge (e.g., non-personal future tasks; Klein et al.,
276 2002).

277

278 When faced with reduced or no access to episodic memory, it may be a natural compensation
279 strategy to rely on semantic information to aid in describing autobiographical events. Using a
280 scoring technique that specifically parses episodic from non-episodic information (Levine et
281 al., 2002), we have also found that although older adults show a decline in the amount of
282 episodic detail comprising their past and future events, they show a corresponding increase in

283 the amount of non-episodic, conceptual information (Addis et al., 2010; Addis et al., 2008;
 284 note also that this pattern extends to picture description; Gaesser et al., 2011). In line with
 285 this finding, older adults also show an increase, relative to young, in their recruitment of
 286 lateral temporal regions during autobiographical tasks (Addis et al., in revision); these regions
 287 are thought to mediate semantic and conceptual autobiographical information (Addis,
 288 McIntosh, Moscovitch, Crawley, & McAndrews, 2004; Graham, Lee, Brett, & Patterson,
 289 2003).

290

291 Another key question is whether access to episodic details is *sufficient* for future simulation
 292 to occur. It is likely that this ability is only a starting point; once episodic details are
 293 extracted, they still have to be used in a meaningful way, which we have argued requires
 294 additional processes such as detail recombination (e.g., Addis & Schacter, 2008; Schacter &
 295 Addis, 2009). Nonetheless, the findings discussed earlier of intact future simulation
 296 performance in hippocampal amnesics with relatively preserved autobiographical memories
 297 (Squire et al., 2010) suggests that access to episodic details may be sufficient for future
 298 simulation. By contrast, Andelman et al. (2010) reported a case study of a patient, M.C., with
 299 a bilateral hippocampal lesion and loss of autobiographical memory restricted only to the past
 300 3 years. Thus, at 27 years of age, she still had approximately 20 years of episodic memories
 301 to draw upon when completing a future simulation task. M.C. was, however, unable to do so:
 302 when asked to describe her personal future, her responses were vague and general, or she
 303 reported that she simply didn't know. Because there was no quantitative assessment of future
 304 simulation performance in this case, the results must be interpreted cautiously. Still, they
 305 raise the possibility that while access to episodic details may be necessary in order to
 306 construct episodic simulations, it may not be sufficient.

307

308 **3. Detail recombination: constructing a coherent scenario**

309

310 As we have reviewed above, being able to access details from episodic memory can be
 311 conceptualized as an initial stage in the process of episodic simulation. Of course, having a
 312 jumble of details is useless if they cannot be recombined and integrated appropriately. We
 313 have argued that 'detail recombination' is critical to imagining coherent scenarios – the kinds
 314 of simulations one creates when thinking about experiences relevant in their daily lives.
 315 Given the role of the hippocampus, particularly the anterior hippocampus, in relational
 316 processing, we have argued that this region is likely critical in the ability to form coherent
 317 scenarios (e.g., Addis & Schacter, 2008; Schacter & Addis, 2009).

318

319 This proposal is based on an integration of findings from various neuroimaging studies. An
 320 early meta-analysis of medial temporal activity during memory tasks reported that the
 321 anterior portion of the hippocampus appears to be particularly responsive to tasks with
 322 relational demands (Schacter & Wagner, 1999); subsequent work has further supported this
 323 anterior localization of relational memory processes (e.g., Chua, Schacter, Rand-Giovannetti,
 324 & Sperling, 2007; Giovanello, Schnyer, & Verfaellie, 2004; Jackson & Schacter, 2004;
 325 Kirwan & Stark, 2004; Staresina & Davachi, 2008, 2009). The role of this region within the
 326 realm of relational memory may be further refined, based on findings from Preston et al.
 327 (2004; see also Heckers, Zalesak, Weiss, Ditman, & Titone, 2004). This work suggests that
 328 the anterior hippocampus may be particularly involved in the recombination of details
 329 extracted from various memories. Using a transitive inference paradigm, participants first
 330 learned to associate one set of items (faces, A) with another set of items (houses, B). They
 331 then learned to associate those same houses (B) with a new set of items (novel faces, C).

332 During the scanning session, seeing items (A, B, or C) taken from any of the memories (A-B,
333 B-C) resulted in posterior hippocampal activity, further implicating the posterior
334 hippocampus in retrieval or reinstatement. However, seeing novel rearrangements of such
335 details (A-C) resulted in selective anterior hippocampal activity. This recombination process
336 can be considered analogous to future simulation, where we argue details extracted from
337 different memories that may have not been encountered together in reality, are rearranged in
338 imagination – and similarly, this recombination process should also engage the anterior
339 hippocampus.

340

341 More recently, Staresina and Davachi (2009) investigated hippocampal responses to the
342 process of integrating details across time and space. They identified a region in the anterior
343 hippocampus that was more responsive when details were presented in a spatiotemporally
344 discontinuous manner (i.e., separated across time and space) and required integration, relative
345 to when details were presented in a contiguous, integrated form. Conceptually, we suggest
346 that this process again maps onto the kind of recombination thought to occur during
347 simulation: an integration of details from memories formed in different spatiotemporal
348 contexts.

349

350 The findings of Preston et al. (2004) and Staresina and Davachi (2009) dovetail with those
351 from a neuroimaging study of past and future detail. In that study, we (Addis & Schacter,
352 2008) found common responses to detail of past and future events in posterior hippocampus,
353 but the anterior hippocampus was responsive only to the amount of detail comprising future
354 events – which are presumably recombined across spatiotemporally distinct experiences.
355 Interestingly, we have replicated the finding of differential future activity within the anterior
356 hippocampus across a number of studies using autobiographical cuing (e.g., Addis, Wong, et
357 al., 2007; adapted from Crovitz & Schiffman, 1974) and experimental recombination
358 paradigms (Addis, Pan, et al., 2009). While the cuing task requires an individual to generate
359 future events from generic cues (nouns), the experimental recombination paradigm uses
360 random rearrangements of episodic details (persons, places, objects) taken from the
361 individual's own memories, thus ensuring that detail recombination occurs. Moreover, these
362 paradigms enable examination of activity during the initial construction of the future event
363 when the cue is presented, and the subsequent elaboration of the event once it is in mind.
364 With this approach, we have found that over the course of a simulation trial, this activity
365 typically emerges during the initial construction phase rather than being evident throughout
366 the duration of a simulation trial (Addis, Cheng, Roberts, & Schacter, 2011; Addis, Pan, et
367 al., 2009; Addis, Wong, et al., 2007; Martin, Schacter, Corballis, & Addis, 2011). This
368 temporal pattern suggests that the differential future-related activity is associated with
369 processes occurring early in the construction of future events, when detail recombination
370 would be expected to occur. Other labs have also reported similar future>past effects in the
371 anterior hippocampus. For instance, Weiler and colleagues (Weiler, Suchan, & Daum, 2010a)
372 found that imagining future events that had a low probability of occurring during the
373 upcoming holidays was associated with more anterior hippocampal activity than events with
374 a higher probability of occurring. The authors suggested that perhaps low probability events
375 place a higher demand on the binding of disparate event features relative to high probability
376 events that may be already planned.

377

378 Determining the boundary conditions of the future>past effect will provide a better
379 understanding of whether detail recombination is important for engaging the anterior aspect
380 of the hippocampus. Importantly, we have recently shown that this effect is limited to certain

381 types of future events. We examined hippocampal activity when imagining specific (unique)
 382 and general (routine) future events, hypothesizing that constructing a specific future event
 383 should place greater demand on recombining details and hippocampal resources relative to
 384 constructing a generic future event that more closely relies on conceptual knowledge about
 385 routines (Addis et al., 2011). Indeed, our analysis supported this hypothesis, demonstrating
 386 that hippocampal activity was strongest when imagining specific future events relative to
 387 more generic and routinized ones. Participant ratings confirmed that specific future events
 388 were more detailed and novel than general future events, further suggesting that the process
 389 of constructing an event that is both detailed *and* novel engages the anterior hippocampal
 390 region. Additionally, because these findings suggest that the hippocampus is not strongly
 391 engaged by constructing generic future events, it may not be surprising that patients with
 392 hippocampal damage can imagine the future in a gist-like, conceptual manner.

393

394 These observations from neuroimaging studies suggest that dysfunction in the hippocampus
 395 may result in deficits in recombining details. Several findings suggest the presence of such
 396 difficulties. Hassabis et al. (2007) found that not only did the events constructed by
 397 hippocampal amnesics lack content overall, but the details they did generate were not well
 398 integrated and lacked a spatial coherence. In healthy older adults who show some degree of
 399 structural and functional dysfunction in the hippocampus (Hedden & Gabrieli, 2004), we
 400 found that the integration of memory details into simulations was reduced relative to young
 401 adults (Addis et al., 2010). Using the experimental recombination paradigm, we
 402 experimentally ‘extracted’ person, place and object details from different past events; random
 403 recombinations of a participant’s memory details were later presented during a future
 404 simulation task. Importantly, each future simulation was required to include the person, place
 405 and object details presented. While both groups were able to include all three details in the
 406 simulations, the young group was better able to integrate these three details into the same
 407 imagined spatiotemporal context. In contrast, older adults integrated on average two of the
 408 three details into the same spatiotemporal context, and then often touched on the third detail
 409 in a separate context, essentially resulting in a series of ‘mini-events’. These findings
 410 suggests that even with experimental support to access details from various episodic
 411 memories, the ability to integrate these details into a coherent scenario with a specific
 412 temporal and spatial context may be reduced in populations with compromised hippocampal
 413 function.

414

415 Again, one might raise the question that if the hippocampus is necessary for detail
 416 recombination, how is it that some patients with hippocampal damage can imagine seemingly
 417 coherent future events? One issue is that not every study of future simulation in patients
 418 includes a measure of detail integration or spatial coherence and thus in instances where
 419 hippocampal patients can successfully imagine, it can be difficult to determine whether the
 420 scenarios constructed were in fact coherent. Maguire and Hassabis (2011) argue that the
 421 number of spatial references produced by the patients studied by Squire et al. (2010) appear
 422 reduced relative to the typical level of controls, suggesting that these patients may have been
 423 creating primarily semantic representations. Moreover, it is possible to imagine a future event
 424 with minimal, if any, detail recombination: one can “recast” past events into the future. It is
 425 possible that paradigms using single cues may elicit recasting. For instance, if shown the cue
 426 word “car”, one might recall a relevant experience (“my car breaking down and my husband
 427 picking me up”) and then imagine that experience unfolding in the same way in future. In
 428 many protocols, it is ensured that participants are generated novel scenarios (e.g., Addis,
 429 Wong, et al., 2007, 2008; Hassabis et al., 2007), but this is not always done or reported. In

430 order to circumvent this possibility, we designed an experimental recombination paradigm in
 431 which participants are required to recombine details extracted from their own past events
 432 (Addis et al., 2009). Although this paradigm has been employed with older adults (Addis et
 433 al., 2010), replicating our findings using the cue word paradigm, it has not yet been used to
 434 assess recombination abilities in patients with circumscribed hippocampal damage. The
 435 results of such a study would be of considerable interest.

436

437 **4. Memory for the future: encoding future simulations**

438

439 Differential engagement of the anterior hippocampus may also reflect the process of encoding
 440 newly-imagined scenarios. Indeed, the anterior portion of the hippocampus has been
 441 implicated in encoding (Schacter & Wagner, 1999; Spaniol et al., 2009), particularly for
 442 relational (e.g., Chua et al., 2007; Jackson & Schacter, 2004; Kirwan & Stark, 2004;
 443 Staresina & Davachi, 2008, 2009) and novel (Kohler, Danckert, Gati, & Menon, 2005)
 444 information. If the adaptive significance of simulating several alternative “behavioral modes”
 445 is to maximize success in anticipated situations (Ingvar, 1985) and flexible planning (Boyer,
 446 2008), then retaining this “fitness-relevant” information in memory for future reference is a
 447 necessary step. Nairne, Thompson and Pandeirada (2007) investigated whether information
 448 relevant to survival is remembered better than survival-irrelevant information. In that study,
 449 participants judged whether items were relevant to survival (having provisions and
 450 protection) or moving (moving to a foreign country) situations, or judged the items for
 451 pleasantness. In line with the idea that we are tuned to remember fitness-relevant information,
 452 subsequent memory performance was boosted for items rated as survival-relevant.
 453 Interestingly, more recent work using a variant of the paradigm developed by Nairne and
 454 colleagues suggests that the much of the benefit of “survival processing” may be attributable
 455 to the engagement of encoding processes that support planning for the future (Klein,
 456 Robertson, & Delton, 2010).

457

458 Three kinds of evidence demonstrate the adaptive value of simulations. First, it is well
 459 established that simulations play an important role in psychological well-being. Being able to
 460 generate specific and detailed simulations of future events can enhance one’s ability to cope
 461 with upcoming situations (Brown, MacLeod, Tata, & Goddard, 2002; Taylor, Pham, Rivkin,
 462 & Armor, 1998; Taylor & Schneider, 1989). For instance, creating simulations about positive
 463 future outcomes can improve emotion regulation, resulting in decreased amounts of worry
 464 related to upcoming future events (Brown et al., 2002). In addition to helping one cope with
 465 the prospect of an upcoming event, mentally simulating appropriate actions for future
 466 stressful situations can enhance one’s ability to cope if and when those situations arise
 467 (Taylor & Schneider, 1989).

468

469 Second, simulations are used when attempting to solve open-ended or ill-defined problems,
 470 where different possible solution paths need to be mentally evaluated. Using the Means-Ends
 471 Problem Solving Test, Sheldon and colleagues (Sheldon, McAndrews, & Moscovitch, 2011)
 472 examined the ability of older adults and patients with temporal lobe epilepsy to solve open-
 473 ended social problems. Both of these groups are known to have some degree of impairment
 474 on tasks of autobiographical memory (Addis, Moscovitch, & McAndrews, 2007; Levine et
 475 al., 2002; St-Laurent, Moscovitch, Levine, & McAndrews, 2009); older adults are also known
 476 to show reduced performance on episodic simulation tasks (Addis et al., 2008). It was found
 477 that when simulating solutions to ill-defined problems, both groups generated fewer relevant
 478 steps than controls. This finding suggests that without full access to episodic memory and the

479 ability to generate detailed simulations, the effectiveness of problem solving is reduced (for
480 relevant neuroimaging evidence, see Gerlach, Spreng, Gilmore, & Schacter, 2011; Spreng,
481 Stevens, Chamberlain, Gilmore, & Schacter, 2010).

482

483 Third, recent studies have demonstrated that episodic simulation has a significant impact on
484 temporal discounting of future rewards: when people imagine experiencing a reward in the
485 future, they show an increased tendency to favor rewards that produce greater long-term
486 payoffs, thereby countering the normal tendency to devalue delayed rewards (Benoit, Gilbert,
487 & Burgess, 2011; Peters & Büchel, 2010). Interestingly, fMRI data reveal that these effects
488 of episodic simulation on temporal discounting are associated with increased coupling
489 between activity in the hippocampus and prefrontal regions involved in reward representation
490 (Benoit et al., 2011; Peters & Büchel, 2010). Related studies have shown that varying the
491 manner in which memory is queried can also influence temporal discounting toward long-
492 term payoffs when memory queries emphasize the production of patient (vs. impatient)
493 thoughts (Weber et al., 2007). It would be interesting to approach effects of episodic
494 simulation on temporal discounting from the theoretical perspective of query theory
495 (Johnson, Haubl, & Keinan, 2007) and to determine whether the memory-based effects on
496 temporal discounting have a similar neural basis to those shown for episodic simulation.

497

498 In order to influence future behaviors and realize these adaptive benefits of simulation, it is
499 important that simulations are encoded and maintained in memory (Ingvar, 1985; Szpunar,
500 Addis, & Schacter, in press). There is indirect evidence to support this idea. For instance,
501 individuals tend to act in a way that is consistent with or constrained by how they have
502 imagined themselves in those situations (Johnson & Sherman, 1990), implying that some
503 record of that simulation influences later behavior. There is typically a high correspondence
504 of stated intentions and subsequent behavior (Fishbein & Ajzen, 1980). Consider also
505 prospective memory, where an intention is encoded into memory and later accessed and
506 implemented when triggered by a target event or time cue. It is likely that the intentions
507 involved in prospective memory range in the degree to which they draw upon simulations.
508 Particularly relevant to the idea of episodic simulation is the process of forming
509 “implementation intentions” (Gollwitzer, 1999) which involve imagining and rehearsing a
510 plan with reference to the specific future context in which it will be executed. Research has
511 shown that creating implementation intentions significantly increases the likelihood of
512 carrying out that intention (Chasteen, Park, & Schwarz, 2001; Orbell, Hodgkins, & Sheeran,
513 1997), again suggesting that these simulations are not only stored in memory but do influence
514 future behavior. Poppenk and colleagues (Poppenk, Moscovitch, McIntosh, Ozcelik, & Craik,
515 2010) directly investigated the process of encoding intentions, using fMRI to see whether
516 later memory for intentions was associated with hippocampal activity during encoding. They
517 found that successful encoding of intentions engaged the hippocampus, as did the encoding of
518 other forms of information, such as present actions. But unique to the prospective task was
519 the recruitment of frontopolar cortex, consistent with finding that damage to this region
520 results in deficits of prospective memory (e.g., Burgess, Veitch, de Lacy Costello, & Shallice,
521 2000).

522

523 If the involvement of the hippocampus in future simulation is only to encode imagined
524 scenarios, then hippocampal damage would not necessarily result in an inability to construct
525 simulations – just an inability to encode and retain them. There are some data to suggest that
526 this might be the case (see Table 1). For instance, although children with hippocampal
527 damage can imagine scenarios, when asked to recall them the following day, they do so with

528 less accuracy and consistency than healthy controls (Cooper et al., 2011). Additionally, adults
 529 with hippocampal damage appear to repeat themselves more than controls when describing
 530 future events, possibly indicative of a failure to sufficiently encode the scenario as it is
 531 constructed (Squire et al., 2010).

532

533 We conducted an fMRI study (Martin et al., 2011) to investigate whether hippocampal
 534 activity during future simulation is indeed related to successful encoding by incorporating the
 535 experimental recombination (Addis, Pan, et al., 2009) and subsequent memory (e.g., Wagner
 536 et al., 1998) paradigms. During scanning, participants were presented with random
 537 recombinations of person, location, and object details taken from their own memories and for
 538 each set of details, they imagined a novel future event involving all three details. After
 539 scanning, participants completed an unexpected cued recall test, in which they were showed
 540 two details and had to recall the third. By this design, we had an objective measure of
 541 whether the critical details comprising each simulation were successfully encoded. As
 542 predicted, successfully encoded simulations were associated with greater activity in the
 543 anterior right hippocampus than simulations that were later forgotten. Moreover, the posterior
 544 right hippocampus was also modulated by encoding success. A functional connectivity
 545 analysis revealed that both the anterior and posterior hippocampus exhibited connectivity
 546 with each other and a wider brain network (including medial prefrontal and medial parietal
 547 regions) during successful encoding. When encoding was not successful, the posterior
 548 hippocampus did not show this pattern of connectivity. However, it is interesting to note that
 549 during unsuccessful encoding, the anterior region still exhibited connectivity with the wider
 550 core network. It is possible that this neural pattern reflects the attempt to construct a
 551 simulation, even if it is ultimately not encoded sufficiently to be recalled later. We also
 552 found that the imagined events that were later-remembered were on average more detailed
 553 than later-forgotten ones, and activity in regions exhibiting an encoding effect was also
 554 modulated by the level of detail. Together, these observations suggest that constructing a
 555 memorable scenario may be related, at least in part, to how well the composite details were
 556 retrieved from memory and recombined.

557

558 **5. Future directions: mapping component processes to hippocampal regions**

559

560 Considering together the patient and neuroimaging data reviewed here, there appears to be
 561 evidence supporting the idea that there are three important component processes involved in
 562 the simulation of episodic future events. First, details stored in episodic memory with which
 563 to furnish the simulation must be accessed. Second, the details extracted from various
 564 memories need to be recombined and integrated into a spatiotemporal context in order imbue
 565 a simulation with a sense of coherence. Third, if a simulation is to influence and guide future
 566 behaviours, it needs to be successfully encoded into memory. The evidence reviewed herein
 567 suggests that these different processes all rely, to some extent, on the hippocampus. It
 568 remains an open and important question as to whether different subregions of the
 569 hippocampus are specifically associated with specific component processes. While the
 570 posterior hippocampus likely supports the retrieval of previously experienced details,
 571 particularly those spatial in nature, the anterior hippocampus supports the recombination of
 572 extracted details into a coherent scenario, and both regions support successful encoding.

573

574 This framework may be able to inform the debate on whether hippocampal damage disrupts
 575 the ability to imagine the future (Maguire & Hassabis, 2011; Squire et al., 2010). It is critical
 576 that future research on patients with hippocampal damage employ more refined experimental

577 designs to probe whether detail access, detail recombination and/or encoding of simulations
578 is disrupted. The case study approach may particularly important here. There is considerable
579 variance of performance across patients with hippocampal damage, and it will be important to
580 understand the specific patterns of spared and impaired sub-processes within each case.
581 Moreover, it is likely that the nature and location of damage to the hippocampus is critical.
582 Differential impairments of the construction and/or encoding of future simulations may
583 emerge depending on the nature of the hippocampal damage: whether it is confined to the
584 anterior and/or posterior aspects, affects primarily the right hippocampus, affects the entirety
585 of the structure, or extends beyond its boundaries. Moreover, it will be critical in future
586 studies to ascertain whether damage in amnesic patients is restricted to the hippocampus or
587 extends more broadly.

588
589 Another challenge will be to find ways in which to differentiate the process of recombining
590 details to construct a simulation and the encoding of those simulations. These processes are
591 closely related in two ways: cognitively, with more detailed simulations being more
592 successfully encoded; and neurally, with both processes engaging the anterior right
593 hippocampus. As such, they may be difficult to disentangle. One fruitful avenue may be to
594 investigate whether detail recombination and successful encoding are mediated by specific
595 hippocampal subfields. The hippocampal formation is a circuit comprised of several
596 anatomically-distinct subregions, including the dentate gyrus, three cornu ammonis
597 (CA₁/CA₂/CA₃) areas, and the subiculum. Recent work suggests a functional distinction
598 between the input structures into the hippocampus (dentate gyrus/CA₂/CA₃) and the output
599 (subiculum/CA₁). Specifically, while the input structures appear to be involved in encoding,
600 the output structures may be more involved in binding (Carr, Rissman, & Wagner, 2010).
601 Moreover, the finding that the dentate gyrus is involved in encoding is consistent with the
602 hypothesis that the ability to form temporal associations among new experiences that happen
603 close together in time is ultimately dependent upon the continuous production of new-born
604 granule cells in the dentate gyrus (Aimone, Wiles, & Gage, 2006; Deng, Aimone, & Gage,
605 2010). Extrapolating these findings to the realm of future simulation, it is possible that detail
606 recombination during future simulation may be differentially associated with CA₁/subiculum,
607 and successful encoding with dentate gyrus/CA₂/CA₃. Recent developments in ultra-high-
608 field 7T MRI to obtain exceptionally high resolution images of hippocampal subfield
609 anatomy – including distinct layers within subfields (e.g. Kerchner et al., 2010) – will no
610 doubt facilitate more detailed investigations of the roles of different hippocampal subfields.

611
612 Neuroimaging studies to date suggest there may also be lateralization effects in the
613 hippocampal activity that is differentially associated with future thinking. Specifically, we
614 initially reported that hippocampal activity common to past and future events was evident in
615 the left hippocampus, but that the future>past effect was specific to the right hippocampus
616 (Addis, Wong, et al., 2007). A number of other studies finding future-related activity also
617 report a right lateralization (Addis et al., 2011; Martin et al., 2011; Weiler et al., 2010a,
618 2010b), although some studies report such activity is bilateral (Addis, Pan, et al., 2009).
619 Interestingly, a patient with damage that affected only the right hippocampus exhibited
620 difficulties in generating detailed future simulations (Race et al., 2011), suggesting the right
621 hippocampus may indeed be critical to this ability. However, it remains to be determined
622 what specific contribution the right hippocampus might be making to future simulation.

623
624 The research considered here is in an early stage of development. It is only during the past
625 few years that studies examining the contribution of the hippocampus to imagining the future

626 have begun in earnest, and it is clear that much remains to be learned. Further integration of
 627 this new line of work with more firmly established research on hippocampal contributions to
 628 memory encoding and retrieval will be critical to advancing our understanding, as will
 629 integration with animal studies of such related phenomena as prospective coding in the
 630 hippocampus (e.g., Ferbinteanu & Shapiro, 2003; Foster & Wilson, 2006; Johnson & Redish,
 631 2007; for discussion, see Buckner, 2010). We are hopeful that these kinds of studies will
 632 help to increase our understanding of the neural and cognitive processes that link memory
 633 and imagination, and in so doing, provide new insights into how the future depends on the
 634 past.

635

636

637 **Acknowledgements**

638

639 Preparation of this manuscript was supported in part by the Australian Research Council
 640 Centre of Excellence in Cognition and its Disorders (CE110001021), a Royal Society
 641 Marsden Grant (UOA0810) and a Grant-in-Aid for Research and Study Leave from The
 642 University of Auckland awarded to DRA, and a National Institute of Mental Health grant
 643 (MH060941) awarded to D.L.S.

644

645 **References**

646

647 Addis, D. R., Cheng, T., Roberts, R. P., & Schacter, D. L. (2011). Hippocampal contributions
 648 to the episodic simulation of specific and general future events. *Hippocampus* 21,
 649 1045-1052.

650 Addis, D. R., McIntosh, A. R., Moscovitch, M., Crawley, A. P., & McAndrews, M. P.
 651 (2004). Characterizing spatial and temporal features of autobiographical memory
 652 retrieval networks: a partial least squares approach. *NeuroImage* 23, 1460-1471.

653 Addis, D. R., Moscovitch, M., Crawley, A. P., & McAndrews, M. P. (2004). Recollective
 654 qualities modulate hippocampal activation during autobiographical memory retrieval.
 655 *Hippocampus* 14, 752-762.

656 Addis, D. R., Moscovitch, M., & McAndrews, M. P. (2007). Consequences of hippocampal
 657 damage across the autobiographical memory retrieval network in patients with left
 658 temporal lobe epilepsy. *Brain* 130, 2327-2342.

659 Addis, D. R., Musicaro, R., Pan, L., & Schacter, D. L. (2010). Episodic simulation of past
 660 and future events in older adults: Evidence from an experimental recombination task.
 661 *Psychol Aging* 25, 369-376.

662 Addis, D. R., Pan, L., Vu, M.-A., Laiser, N., & Schacter, D. L. (2009). Constructive episodic
 663 simulation of the future and the past: Distinct subsystems of a core brain network
 664 mediate imagining and remembering. *Neuropsychologia* 47, 2222-2238.

665 Addis, D. R., Roberts, R. P., & Schacter, D. L. (2011). Age-related neural changes in
 666 autobiographical remembering and imagining. *Neuropsychologia* 49, 3656-3669.

667 Addis, D. R., Sacchetti, D. C., Ally, B. A., Budson, A. E., & Schacter, D. L. (2009). Episodic
 668 simulation of future events is impaired in mild Alzheimer's disease. *Neuropsychologia*
 669 47, 2660-2671.

670 Addis, D. R., & Schacter, D. L. (2008). Effects of detail and temporal distance of past and
 671 future events on the engagement of a common neural network. *Hippocampus* 18, 227-
 672 237.

- 673 Addis, D. R., Wong, A. T., & Schacter, D. L. (2007). Remembering the past and imagining
674 the future: Common and distinct neural substrates during event construction and
675 elaboration. *Neuropsychologia* 45, 1363-1377.
- 676 Addis, D. R., Wong, A. T., & Schacter, D. L. (2008). Age-related changes in the episodic
677 simulation of future events. *Psychol Sci* 19, 33-41.
- 678 Aimone, J. B., Wiles, J., & Gage, F. H. (2006). Potential role for adult neurogenesis in the
679 encoding of time in new memories. *Nat Neurosci* 9, 723-727.
- 680 Andelman, F., Hoofien, D., Goldberg, I., Aizenstein, O., & Neufeld, M. Y. (2010). Bilateral
681 hippocampal lesion and a selective impairment of the ability for mental time travel.
682 *Neurocase* 16, 426-435.
- 683 Bayley, P. J., Gold, J. J., Hopkins, R. O., & Squire, L. R. (2005). The Neuroanatomy of
684 Remote Memory. *Neuron* 46, 799-810.
- 685 Benoit, R. G., Gilbert, S. J., & Burgess, P. W. (2011). A neural mechanism mediating the
686 impact of episodic prospection on farsighted decisions. *J Neurosci* 31, 6771-6779.
- 687 Boyer, P. (2008). Evolutionary economics of mental time travel? *Trends Cogn Sci* 12, 219-
688 224.
- 689 Brown, G. P., MacLeod, A. K., Tata, P., & Goddard, L. (2002). Worry and the simulation of
690 future outcomes. *Anxiety, Stress and Coping* 15, 1-17.
- 691 Buckner, R. L. (2010). The role of the hippocampus in prediction and imagination. *Annu Rev*
692 *Psychol* 61, 27-48.
- 693 Buckner, R. L., Andrews-Hanna, J. R., & Schacter, D. L. (2008). The brain's default
694 network: anatomy, function, and relevance to disease. *Ann N Y Acad Sci* 1124, 1-38.
- 695 Buckner, R. L., & Carroll, D. C. (2007). Self-projection and the brain. *Trends Cogn Sci* 11,
696 49-57.
- 697 Burgess, P. W., Veitch, E., de Lacy Costello, A., & Shallice, T. (2000). The cognitive and
698 neuroanatomical correlates of multitasking. *Neuropsychologia* 38, 848-863.
- 699 Carr, V. A., Rissman, J., & Wagner, A. D. (2010). Imaging the human medial temporal lobe
700 with high-resolution fMRI. *Neuron* 65, 298-308.
- 701 Chan, D., Henley, S. M. D, Rossor, M. N., Warrington, E. K. (2007). Extensive and
702 Temporally Ungraded Retrograde Amnesia in Encephalitis Associated With
703 Antibodies to Voltage-Gated Potassium Channels. *Arch Neurol* 64, 404-410
- 704 Chasteen, A. L., Park, D. C., & Schwarz, N. (2001). Implementation intentions and
705 facilitation of prospective memory. *Psychol Sci* 12, 457-461.
- 706 Chua, E., Schacter, D. L., Rand-Giovannetti, E., & Sperling, R. A. (2007). Evidence for a
707 specific role of the anterior hippocampal region in successful associative encoding.
708 *Hippocampus* 17, 1071-1080.
- 709 Cooper, J. M., Vargha-Khadem, F., Gadian, D. G., & Maguire, E. A. (2011). The effect of
710 hippocampal damage in children on recalling the past and imagining new experiences.
711 *Neuropsychologia* 49, 1843-1850.
- 712 Crovitz, H. F., & Schiffman, H. (1974). Frequency of episodic memories as a function of
713 their age. *Psychon Bull Rev* 4, 517-518.
- 714 Deng, W., Aimone, J. B., & Gage, F. H. (2010). New neurons and new memories: how does
715 adult hippocampal neurogenesis affect learning and memory? *Nat Rev Neurosci* 11,
716 339-350.
- 717 Eichenbaum, H. (2001). The hippocampus and declarative memory: cognitive mechanisms
718 and neural codes. *Behav Brain Res* 127, 199-207.
- 719 Ferbinteanu, J., & Shapiro, M. L. (2003). Prospective and retrospective memory coding in the
720 hippocampus. *Neuron* 40, 1227-1239.

- 721 Fishbein, M., & Ajzen, I. (1980). *Belief, Attitude, Intention, and Behavior*. Reading, MA:
722 Addison-Wesley.
- 723 Foster, D. J., & Wilson, M. A. (2006). Reverse replay of behavioural sequences in
724 hippocampal place cells during the awake state. *Nature* 440, 680–683.
- 725 Gaesser, D., Sacchetti, D. C., Addis, D. R., & Schacter, D. L. (2011). Characterizing age-
726 related changes in remembering the past and imagining the future. *Psychol Aging* 26,
727 80-84.
- 728 Gerlach, K. D., Spreng, R. N., Gilmore, A. W., & Schacter, D. L. (2011). Solving future
729 problems: Default network and executive activity associated with goal-directed
730 mental simulations. *NeuroImage* 55, 1816-1824.
- 731 Giovanello, K. S., Schnyer, D., & Verfaellie, M. (2004). A critical role for the anterior
732 hippocampus in relational memory: evidence from an fMRI study comparing
733 associative and item recognition. *Hippocampus* 14, 5-8.
- 734 Giovanello, K. S., Schnyer, D., & Verfaellie, M. (2009). Distinct hippocampal regions make
735 unique contributions to relational memory. *Hippocampus* 19, 111-117.
- 736 Gold, J. J., & Squire, L. R. (2005). Quantifying medial temporal lobe damage in memory-
737 impaired patients. *Hippocampus* 15, 79-85.
- 738 Gollwitzer, P. M. (1999). Implementation intentions: Strong effects of simple plans. *Am*
739 *Psychol* 54, 493-503.
- 740 Graham, K. S., Lee, A. C., Brett, M., & Patterson, K. (2003). The neural basis of
741 autobiographical and semantic memory: new evidence from three PET studies. *Cogn*
742 *Affect Behav Neurosci* 3, 234-254.
- 743 Hassabis, D., Kumaran, D., Vann, S. D., & Maguire, E. A. (2007). Patients with hippocampal
744 amnesia cannot imagine new experiences. *Proc Natl Acad Sci USA* 104, 1726-1731.
- 745 Hassabis, D., & Maguire, E. A. (2007). Deconstructing episodic memory with construction.
746 *Trends Cogn Sci* 11, 299-306.
- 747 Heckers, S., Zalesak, M., Weiss, A. P., Ditman, T., & Titone, D. (2004). Hippocampal
748 activation during transitive inference in humans. *Hippocampus* 14, 153-162.
- 749 Hedden, T., & Gabrieli, J. D. (2004). Insights into the ageing mind: A view from cognitive
750 neuroscience. *Nat Rev Neurosci* 5, 87-96.
- 751 Hurley, N.C., Maguire, E.A., & Vargha-Khadem, F. (in press). Patient HC with
752 developmental amnesia can construct future scenarios. *Neuropsychologia*.
- 753 Ingvar, D. H. (1985). ‘Memory of the future’: an essay on the temporal organization of
754 conscious awareness. *Hum Neurobiol* 4, 127–136.
- 755 Jackson, O., 3rd, & Schacter, D. L. (2004). Encoding activity in anterior medial temporal
756 lobe supports subsequent associative recognition. *NeuroImage* 21, 456-462.
- 757 Johnson, A., & Redish, A. D. (2007). Neural ensembles in CA3 transiently encode paths
758 forward of the animal at a decision point. *J Neurosci* 27, 12176–12189.
- 759 Johnson, M. K., & Sherman, S. J. (1990). “Constructing and reconstructing the past and the
760 future in the present,” in *Handbook of Motivation and Cognition: Foundations of*
761 *Social Behavior Vol. 2*, eds. E. T. Higgins and R. M. Sorrentino (New York: The
762 Guilford Press), 482-526.
- 763 Kirwan, C. B., & Stark, C. E. (2004). Medial temporal lobe activation during encoding and
764 retrieval of novel face-name pairs. *Hippocampus* 14, 919-930.
- 765 Klein, S. B., Loftus, J., & Kihlstrom, J. F. (2002). Memory and temporal experience: The
766 effects of episodic memory loss on an amnesic patient's ability to remember the past
767 and imagine the future. *Social Cogn* 20, 353-379.
- 768 Klein, S. B., Robertson, T. E., & Delton, A. W. (2010). Facing the future: memory as an
769 evolved system for planning future acts. *Mem Cognit* 38, 13-22.

- 770 Kohler, S., Danckert, S., Gati, J. S., & Menon, R. S. (2005). Novelty responses to relational
771 and non-relational information in the hippocampus and the parahippocampal region: a
772 comparison based on event-related fMRI. *Hippocampus* 15, 763-774.
- 773 Kwan, D., Carson, N., Addis, D. R., & Rosenbaum, R. S. (2010). Deficits in past
774 remembering extend to future imagining in a case of developmental amnesia.
775 *Neuropsychologia* 48, 3179-3186.
- 776 Lepage, M., Habib, R., & Tulving, E. (1998). Hippocampal PET activations of memory
777 encoding and retrieval: the HIPER model. *Hippocampus* 8, 313-322.
- 778 Levine, B., Svoboda, E., Hay, J. F., Winocur, G., & Moscovitch, M. (2002). Aging and
779 autobiographical memory: dissociating episodic from semantic retrieval. *Psychol*
780 *Aging* 17, 677-689.
- 781 Levy, D. A., Bayley, P. J., & Squire, L. R. (2004). The anatomy of semantic knowledge:
782 medial vs. lateral temporal lobe. *Proc Natl Acad Sci USA* 101, 6710-6715.
- 783 Maguire, E. A., & Hassabis, D. (2011). Role of the hippocampus in imagination and future
784 thinking. *Proc Natl Acad Sci USA* 108, E39.
- 785 Maguire, E. A., Nannery, R., Spiers, H. J. (2006). Navigation around London by a taxi driver
786 with bilateral hippocampal lesions. *Brain* 129, 2894-2907.
- 787 Maguire, E. A., Vargha-Khadem, F., & Hassabis, D. (2010). Imagining fictitious and future
788 experiences: evidence from developmental amnesia. *Neuropsychologia* 48, 3187-
789 3192.
- 790 Maguire, E. A., Vargha-Khadem, F., & Mishkin, M. (2001). The effects of bilateral
791 hippocampal damage on fMRI regional activations and interactions during memory
792 retrieval. *Brain* 124, 1156-1170.
- 793 Martin, V. C., Schacter, D. L., Corballis, M. C., & Addis, D. R. (2011). A role for the
794 hippocampus in encoding simulations of future events. *Proc Natl Acad Sci USA* 108,
795 13858-13863.
- 796 McKenna, P., & Gerhand, S. (2002). Preserved semantic learning in an amnesic patient.
797 *Cortex* 38, 37-58.
- 798 Moscovitch, M., Rosenbaum, R. S., Gilboa, A., Addis, D. R., Westmacott, R., Grady, C., . . .
799 Nadel, L. (2005). Functional neuroanatomy of remote episodic, semantic and spatial
800 memory: A unified account based on multiple trace theory. *J Anat* 207, 35-66.
- 801 Nairne, J. S., Thompson, S. R., & Pandeirada, J. N. S. (2007). Adaptive memory: Survival
802 processing enhances retention. *J Exp Psychol Gen* 33, 263-273.
- 803 Orbell, S., Hodgkins, S., & Sheeran, P. (1997). Implementation intentions and the theory of
804 planned behavior. *Pers Soc Psychol Bull* 23, 945-954.
- 805 Peters, J., & Büchel, C. (2010). Episodic future thinking reduces reward delay discounting
806 through an enhancement of prefrontal-mediotemporal interactions. *Neuron* 66, 138-
807 148.
- 808 Poppenk, J., Moscovitch, M., McIntosh, A. R., Ozcelik, E., & Craik, F. I. M. (2010).
809 Encoding the future: Successful processing of intentions engages predictive brain
810 networks. *NeuroImage* 49, 905-913.
- 811 Preston, A. R., Shrager, Y., Dudukovic, N. M., & Gabrieli, J. D. (2004). Hippocampal
812 contribution to the novel use of relational information in declarative memory.
813 *Hippocampus* 14, 148-152.
- 814 Prince, S. E., Daselaar, S. M., & Cabeza, R. (2005). Neural correlates of relational memory:
815 successful encoding and retrieval of semantic and perceptual associations. *J Neurosci*
816 25, 1203-1210.

- 817 Race, E., Keane, M. M., & Verfaellie, M. (2011). Medial temporal lobe damage causes
 818 deficits in episodic memory and episodic future thinking not attributable to deficits in
 819 narrative construction. *J Neurosci* 31, 10262-10269.
- 820 Rosenbaum, R. S., Köhler, S., Schacter, D. L., Moscovitch, M., Westmacott, R., Black, S. E.,
 821 . . . Tulving, E. (2005). The case of K.C.: contributions of a memory-impaired person
 822 to memory theory. *Neuropsychologia* 43, 989-1021.
- 823 Samarasekera, S. R., Vincent, A., Welch, J. L., Jackson, M., Nichols, P., & Griffiths, T. D.
 824 (2006). Course and outcome of acute limbic encephalitis with negative voltage-gated
 825 potassium channel antibodies. *J Neurol Neurosurg Psychiatry* 78, 391-394.
- 826 Schacter, D. L., & Addis, D. R. (2007). The cognitive neuroscience of constructive memory:
 827 Remembering the past and imagining the future. *Philos Trans R Soc Lond B Biol Sci*
 828 362, 773-786.
- 829 Schacter, D. L., & Addis, D. R. (2009). On the nature of medial temporal lobe contributions
 830 to the constructive simulation of future events. *Philos Trans R Soc Lond B Biol Sci*
 831 364, 1245-1253.
- 832 Schacter, D. L., Addis, D. R., & Buckner, R. L. (2007). The prospective brain: Remembering
 833 the past to imagine the future. *Nat Rev Neurosci* 8, 657-661.
- 834 Schacter, D. L., Addis, D. R., & Buckner, R. L. (2008). Episodic simulation of future events:
 835 concepts, data, and applications. *Ann N Y Acad Sci* 1124, 39-60.
- 836 Schacter, D. L., Gaesser, B., & Addis, D. R. (2011). "Age-related changes in the episodic
 837 simulation of past and future events," in *Successful Remembering and Successful*
 838 *Forgetting: A Festschrift in Honor of Robert A. Bjork*, ed. A. S. Benjamin (New
 839 York: Psychology Press), 505-525.
- 840 Schacter, D.L., Guerin, S.A., & St. Jacques, P.L. (2011). Memory distortion: An adaptive
 841 perspective. *Trends Cogn Sci* 15, 467-474.
- 842 Schacter, D. L., & Wagner, A. D. (1999). Medial temporal lobe activations in fMRI and PET
 843 studies of episodic encoding and retrieval. *Hippocampus* 9, 7-24.
- 844 Sheldon, S., McAndrews, M. P., & Moscovitch, M. (2011). Episodic memory processes
 845 mediated by the medial temporal lobes contribute to open-ended problem solving.
 846 *Neuropsychologia* 49, 2439-2447.
- 847 Spaniol, J., Davidson, P. S., Kim, A. S., Han, H., Moscovitch, M., & Grady, C. L. (2009).
 848 Event-related fMRI studies of episodic encoding and retrieval: meta-analyses using
 849 activation likelihood estimation. *Neuropsychologia* 47, 1765-1779.
- 850 Spreng, R. N., Mar, R. A., & Kim, A. S. N. (2009). The common neural basis of
 851 autobiographical memory, prospection, navigation, theory of mind and the default
 852 mode: a quantitative meta-analysis. *J Cogn Neurosci* 21, 489-510.
- 853 Spreng, R. N., Stevens, W. D., Chamberlain, J. P., Gilmore, A. W., & Schacter, D. L. (2010).
 854 Default network activity, coupled with the frontoparietal control network, supports
 855 goal-directed cognition. *NeuroImage* 53, 303-317.
- 856 Squire, L. R., McDuff, S. G., & Frascino, J. C. (2011). Reply to Maguire and Hassabis:
 857 Autobiographical memory and future imagining. *Proc Natl Acad Sci USA* 108, E40
- 858 Squire, L. R., van der Horst, A. S., McDuff, S. G. R., Frascino, J. C., Hopkinse, R. O., &
 859 Mauldin, K. N. (2010). Role of the hippocampus in remembering the past and
 860 imagining the future. *Proc Natl Acad Sci USA* 107, 19044-19048.
- 861 St-Laurent, M., Moscovitch, M., Levine, B., & McAndrews, M. P. (2009). Determinants of
 862 autobiographical memory in patients with unilateral temporal lobe epilepsy or
 863 excisions. *Neuropsychologia* 47, 2211-2221.

- 864 Staresina, B. P., & Davachi, L. (2008). Selective and shared contributions of the
 865 hippocampus and perirhinal cortex to episodic item and associative encoding. *J Cogn*
 866 *Neurosci* 20, 1478-1489.
- 867 Staresina, B. P., & Davachi, L. (2009). Mind the gap: binding experiences across space and
 868 time in the human hippocampus. *Neuron* 63, 267-276.
- 869 Suddendorf, T., & Corballis, M. C. (1997). Mental time travel and the evolution of the human
 870 mind. *Genet Soc Gen Psychol Monogr* 123, 133-167.
- 871 Suddendorf, T., & Corballis, M. C. (2007). The evolution of foresight: What is mental time
 872 travel, and is it unique to humans? *Behav Brain Sci* 30, 299-351.
- 873 Szpunar, K. K. (2010). Episodic future thought: An emerging concept. *Perspectives on*
 874 *Psychol Sci* 5, 142-162.
- 875 Szpubar, K.K. Addis, D.R., & Schacter, D.L. (in press). Memory for emotional simulations:
 876 Remembering a rosy future. *Psychol Sci*.
- 877 Taylor, S. E., Pham, L. B., Rivkin, I. D., & Armor, D. A. (1998). Harnessing the imagination:
 878 Mental simulation, self-regulation and coping. *American Psychologist*, 53, 429-439.
- 879 Taylor, S. E., & Schneider, S. K. (1989). Coping and the simulation of events. *Social Cogn* 7,
 880 174-194.
- 881 Tulving, E. (1985). Memory and consciousness. *Canadian Psychologist* 25, 1-12.
- 882 Vharga-Khadem, F., Gadian, D. G., Watkins, K. E., Connelly, A., Van Paesschen, W., &
 883 Mishkin, M. (1997). Differential effects of early hippocampal pathology on episodic
 884 and semantic memory. *Science* 277, 376-380.
- 885 Wagner, A. D., Schacter, D. L., Rotte, M., Koutstaal, W., Maril, A., Dale, A. M., . . .
 886 Buckner, R. L. (1998). Building memories: remembering and forgetting of verbal
 887 experiences as predicted by brain activity. *Science* 281, 1188-1191.
- 888 Weiler, J. A., Suchan, B., & Daum, I. (2010a). Foreseeing the Future: Occurrence Probability
 889 of Imagined Future Events Modulates Hippocampal Activation. *Hippocampus* 20,
 890 685-690.
- 891 Weiler, J. A., Suchan, B., & Daum, I. (2010b). When the future becomes the past:
 892 Differences in brain activation patterns for episodic memory and episodic future
 893 thinking. *Behav Brain Res* 212, 196-203.