



The Innate Immune Protein Nod2 Binds Directly to MDP, a Bacterial Cell Wall Fragment

Citation

Grimes, Catherine Leimkuhler, Lushanti De Zoysa Ariyananda, James E. Melnyk, and Erin K. O'Shea. 2012. The innate immune protein nod2 binds directly to mdp, a bacterial cell wall fragment. *Journal of the American Chemical Society* 134(33): 13535-13537.

Published version

<https://doi.org/10.1021/ja303883c>

Link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:11210616>

Terms of use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material (LAA), as set forth at

<https://harvardwiki.atlassian.net/wiki/external/NGY5NDE4ZjgzNTc5NDQzMGIzZWZhMGFIOWI2M2EwYTg>

Accessibility

<https://accessibility.huit.harvard.edu/digital-accessibility-policy>

Share Your Story

The Harvard community has made this article openly available.

Please share how this access benefits you. [Submit a story](#)

The innate immune protein Nod2 binds directly to MDP, a bacterial cell wall fragment

Catherine Leimkuhler Grimes*[§], Lushanti De Zoysa Ariyananda[§], James E. Melnyk[§], and Erin K. O'Shea[‡]

[§]Department of Chemistry and Biochemistry, University of Delaware, Newark, DE 19716. [‡] Howard Hughes Medical Institute, Harvard Faculty of Arts and Sciences Center for Systems Biology, Departments of Molecular and Cellular Biology and of Chemistry and Chemical Biology, Harvard University, Cambridge, MA 02138.

innate immunity, nod-like-receptors, bacterial cell wall, Crohn's disease, muramyl dipeptide (MDP)

Supporting Information Placeholder

ABSTRACT: Mammalian Nod2 is an intracellular protein that is implicated in the innate immune response to the bacterial cell wall and is associated with the development of Crohn's disease, Blau syndrome and gastrointestinal cancers. Nod2 is required for an immune response to muramyl dipeptide (MDP), an immunostimulatory fragment of bacterial cell wall, but it is not known if MDP binds directly to Nod2. We report the expression and purification of human Nod2 from insect cells. Using novel MDP-self-assembled monolayers (SAMs) we provide the first biochemical evidence for a direct, high affinity interaction between Nod2 and MDP.

The innate immune system is the body's first line of defense against invading pathogens^{1,2}. This ancient system has evolved to exist in a symbiotic relationship with commensal bacteria and at the same time to recognize and destroy virulent bacteria¹⁻³. Chronic inflammatory diseases such as asthma, rheumatoid arthritis, and Crohn's disease are thought to arise from an inappropriate innate immune response to bacteria⁴⁻⁹. Chronic inflammation has also been shown to lead to a variety of types of cancers including those affecting gastric, colon, and lung¹⁰.

Mammalian Nod2 is an intracellular protein that is involved in the signaling response to bacterial cell wall fragments^{11,12}. Mutations in Nod2 correlate with the development of Crohn's disease, a chronic inflammatory disease of the gastrointestinal tract¹³⁻¹⁵. In order to generate the proper immunologic response, the Nod2 signaling pathway must recognize bacteria. The biochemical mechanism by which Nod2 detects bacteria is not known, but it has been proposed that Nod2 senses bacterial cell wall fragments directly. Nod2 could sense bacterial cell wall fragments through at least three mechanisms: (1) a direct interaction; (2) a mediated interaction; or (3) a signaling relay. The mechanism of activation remains unresolved because until now the proper tools to probe the mechanism have been unavailable.

Nod2 is essential for the cellular response to a small fragment of bacterial cell wall, muramyl dipeptide, consisting of one carbohydrate and two amino acids (**MDP-(D)**, **1**, Figure 1)^{12,16}. MDP is found in both Gram-negative and Gram-positive bacteria. Cellular and *in vivo* assays have shown that when mammalian cells expressing Nod2 are treated with MDP, an inflammatory response is activated via the NF- κ B and MAP

kinase pathways¹⁷⁻²⁰. Moreover, the response is not observed if a diastereomer of **1**, (**MDP-(L)**, **2**, Figure 1), is used in the cellular assays. For this reason, MDP is often referred to in the literature as the "ligand" for Nod2 signaling²¹. However, there is no biochemical or biophysical data that demonstrates an interaction between the two molecules²². The purpose of our investigation was to determine if Nod2 and MDP interact *in vitro*. We developed an expression system and a biochemical assay using synthetic probes to investigate this question.




Figure 1: Muramyl Dipeptides: MDP-(D) is the biologically relevant isomer; MDP-(L) is a synthetic diastereomer of the compound found in Nature.

Human Nod2 is a large protein (1040 residues, 110 kD) with multiple domains: two N-terminal caspase recruitment domains (CARDs), a central nucleotide oligomerization domain (NOD) and a C-terminal leucine rich repeat (LRR) domain¹⁶. To determine if Nod2 interacts directly with MDP we first expressed a Flag-tagged version of Nod2 using baculovirus-infected Sf21 cells (Figure S1a in the Supporting Information) with a yield of 1 mg/L. CD spectroscopy and limited proteolysis experiments are consistent with Nod2 being a folded protein (Figure S1b and S1c in the Supporting Information).

With purified Nod2 in hand, a Surface Plasmon Resonance (SPR) assay was developed to assess binding to MDP. Initial attempts to develop a SPR assay with biotinylated-MDP²³ failed, as we observed significant non-specific binding of Nod2 to the streptavidin/biotin chip lacking MDP (Figure S2). In order to develop a SPR assay, we coupled 6-amino MDP (**3** & **4**, Figure 1) directly to the chip without the use of biotin. **3** & **4** are synthetic intermediates of the biotinylated -MDPs and have been shown to activate Nod2 in the appropriate manner²³. Using methodology developed by Whitesides and co-workers, we prepared carboxy-terminated alkane thiol self assembled monolayers (SAMs) and then used on-chip NHS/EDC activation of the carboxylic acid (Figure 2)²⁴ to couple 6-amino-MDPs to the chip surface.

Carboxylic Acid


MDP - Chip

3-Lane: $R = \text{H}_2\text{N}(\text{CH}_2)_3\text{COOH}$


4-Lane: $R = \text{H}_2\text{N}(\text{CH}_2)_4\text{COOH}$

Control-Lane: $R = \text{H}_2\text{N}(\text{CH}_2)_2\text{COOH}$


Nod2 bound to MDP with high affinity (Figure 3). The biologically active MDP, **3-Lane**, bound to Nod2 with a K_D of $51 \text{ nM} \pm 18$. Surprisingly, we found that Nod2 was able to bind to both isomers of MDP (Figure 3), as Nod2 bound to the **4-Lane** with a K_D of $150 \text{ nM} \pm 24$, which is only slightly higher than the K_D observed for the **3-Lane**. Thus, the iso-glutamine stereochemistry is not a key recognition determinant. To demonstrate that binding of Nod2 to the MDP-chip is indeed specific, a competition study was performed. When Nod2 was pre-treated with either **3** or **4**, diminished binding to the chip was observed (Figure 4). Encouragingly, the competition data show the same trend as the on-chip data, i.e. the free D-isomer of MDP is a better competitor than the free L-isomer. These data suggest that Nod2 senses bacterial cell wall fragments by binding directly to them.



resonance signal was recorded, ethanolamine background was subtracted from the experimental signal and plotted. The data was fitted to a standard one site binding model by non-linear regression analysis using Prism 4 (GraphPad).



At low concentrations the MDP isomers **2** and **4** do not activate the NF- κ B response via Nod2 in cellular assays^{18,23}. However, we show that MDP-(L) is able to activate the pathway at higher concentrations using the established cellular NF- κ B luciferase reporter assay and transfected Nod2 (Figure 5). The NF- κ B activation observed in the absence of transfected Nod2 DNA is the result of low levels of endogenous Nod2 in Hek293T cells²⁶. The cellular assay results demonstrate that both isomers of MDP are able to activate the Nod2/NF- κ B pathway, which is consistent with the binding data showing that Nod2 can bind one isomer better than the other. Cellular potencies often do not exactly match their *in vitro* K_D s^{27,28}. We find the cellular activation parallels *in vitro* binding but at slightly lower potency.



The *in vitro* interaction between Nod2 and MDP is dependent on pH (Figure 6), with the pH range from 5.0 to 6.5 giving maximal binding. The data suggest that *in vivo* binding could occur in an acidic cellular compartment, a model supported by cellular assays that show pH-dependent internalization of MDP^{29,30}. Girardin and coworkers show that the internalization of MDP is optimal in the pH range of 5.5 to 6.5, which corresponds with the MDP/Nod2 binding data.

Nod2 is predicted to have ATP binding capabilities, as it has Walker A and Walker B regions³¹. The ATP binding capacities of the protein have been suggested to be important for

oligomerization, protein-protein interactions and subsequent activation³². To test if the nucleotide binding was necessary for Nod2 to bind to MDP, we measured Nod2 binding to MDP $\pm$ ATP/ADP. Nod2 binds with no appreciable change to MDP in the presence and absence of 10 μ M ATP/ADP (Figure 6), suggesting that ATP/ADP is not necessary for Nod2 to bind to MDP.




Figure 6: Nod2/MDPs interact under a variety of conditions. The pH of Nod2 was adjusted before application to the MDP-Chip. Nod2 was pre-incubated with 10 μ M ATP or ADP before application to the MDP-Chip. The relative resonance signal was recorded after ten minutes.

Prior to our investigation, the mechanism of Nod2 activation of NF- κ B by treatment with MDP was unclear. We have taken a biochemical approach to demonstrate that Nod2 binds directly to bacterial cell wall fragments. Recombinant Nod2 and the synthetic MDP tools allowed for the development of *in vitro* assay to detect binding. The assay that we have developed will be a valuable asset in screening for inhibitors/activators of the Nod2 signaling pathway and determining if Nod2 is able to differentiate commensal versus pathogenic bacteria. In addition, the assay will be useful in determining the Nod2 Crohn's mutants are capable of binding to MDP. This is the first biochemical evidence to show an interaction between the two molecules, and establishes that MDP is a high-affinity ligand for Nod2.

ASSOCIATED CONTENT

Supporting Information. Nod2 expression and purification conditions, SPR assay, and NF- κ B assay conditions: this material is available free of charge via the Internet at <http://pubs.acs.org>.

AUTHOR INFORMATION

Corresponding Author

* cgrimes@udel.edu

Author Contributions

The manuscript was written through contributions of all authors. All authors have given approval to the final version of the manuscript.

Funding Sources

This work was supported by start up funds to CLG from the University of Delaware and a NIH-COBRE(P20RR017716-09). EKO is an investigator of the Howard Hughes Medical Institute.

ACKNOWLEDGMENT

We thank Young-Sam Lee for assistance with the development of the SPR assay and John Koh, Colin Thorpe, Neal Zondlo, Brian Bahnson and Vishnu Mohanan, for critical reading of this manuscript.

ABBREVIATIONS

MDP, muramyl dipeptide; SPR, surface Plasmon resonance; SAMs, self assembled monolayers (SAMs)

REFERENCES

- (1) Janeway, C. A., Jr.; Medzhitov, R. *Annu Rev Immunol* **2002**, *20*, 197.
- (2) Akira, S.; Uematsu, S.; Takeuchi, O. *Cell* **2006**, *124*, 783.
- (3) Medzhitov, R. *Nature* **2007**, *449*, 819.
- (4) Jackson, D. J.; Sykes, A.; Mallia, P.; Johnston, S. L. *J Allergy Clin Immunol* **2011**, *128*, 1165.
- (5) Forsythe, P. *Chest* **2011**, *139*, 901.
- (6) Weissmann, G. *Bulletin of the NYU hospital for joint diseases* **2009**, *67*, 251.
- (7) Falgarone, G.; Jaen, O.; Boissier, M. C. *Joint, bone, spine : revue du rhumatisme* **2005**, *72*, 17.
- (8) Chassaing, B.; Darfeuille-Michaud, A. *Gastroenterology* **2011**, *140*, 1720.
- (9) Xavier, R. J.; Podolsky, D. K. *Nature* **2007**, *448*, 427.
- (10) Grivennikov, S. I.; Greten, F. R.; Karin, M. *Cell* **2010**, *140*, 883.
- (11) Chen, G.; Shaw, M. H.; Kim, Y. G.; Nunez, G. *Annu Rev Pathol* **2009**, *4*, 365.
- (12) Magalhaes, J. G.; Sorbara, M. T.; Girardin, S. E.; Philpott, D. J. *Curr Opin Immunol* **2011**, *23*, 29.
- (13) Ogura, Y.; Bonen, D. K.; Inohara, N.; Nicolae, D. L.; Chen, F. F.; Ramos, R.; Britton, H.; Moran, T.; Karaliuskas, R.; Duerr, R. H.; Achkar, J. P.; Brant, S. R.; Bayless, T. M.; Kirschner, B. S.; Hanauer, S. B.; Nunez, G.; Cho, J. H. *Nature* **2001**, *411*, 603.
- (14) Hugot, J. P.; Chamaillard, M.; Zouali, H.; Lesage, S.; Cezard, J. P.; Belaiche, J.; Almer, S.; Tysk, C.; O'Morain, C. A.; Gassull, M.; Binder, V.; Finkel, Y.; Cortot, A.; Modigliani, R.; Laurent-Puig, P.; Gower-Rousseau, C.; Macry, J.; Colombel, J. F.; Sahbatou, M.; Thomas, G. *Nature* **2001**, *411*, 599.
- (15) Cavanaugh, J. *World journal of gastroenterology : WJG* **2006**, *12*, 3673.
- (16) McDonald, C.; Inohara, N.; Nunez, G. *J Biol Chem* **2005**, *280*, 20177.
- (17) Kobayashi, K. S.; Chamaillard, M.; Ogura, Y.; Henegariu, O.; Inohara, N.; Nunez, G.; Flavell, R. A. *Science* **2005**, *307*, 731.
- (18) Girardin, S. E.; Boneca, I. G.; Viala, J.; Chamaillard, M.; Labigne, A.; Thomas, G.; Philpott, D. J.; Sansonetti, P. J. *J Biol Chem* **2003**, *278*, 8869.
- (19) Girardin, S. E.; Travassos, L. H.; Herve, M.; Blanot, D.; Boneca, I. G.; Philpott, D. J.; Sansonetti, P. J.; Mengin-Lecreux, D. *J Biol Chem* **2003**, *278*, 41702.
- (20) Ting, J. P.; Duncan, J. A.; Lei, Y. *Science* **2010**, *327*, 286.
- (21) Fujimoto, Y.; Pradipta, A. R.; Inohara, N.; Fukase, K. *Natural product reports* **2012**, *29*, 568.
- (22) Boudreau, M. A.; Fisher, J. F.; Mobashery, S. *Biochemistry* **2012**, *51*, 2974.
- (23) Grimes, C. L.; Podolsky, D. K.; O'Shea, E. K. *Bioorg Med Chem Lett* **2010**, *20*, 6061.
- (24) Lahiri, J.; Isaacs, L.; Tien, J.; Whitesides, G. M. *Anal Chem* **1999**, *71*, 777.
- (25) Mrksich, M.; Whitesides, G. M. *Annual review of biophysics and biomolecular structure* **1996**, *25*, 55.
- (26) Zhao, L.; Kwon, M. J.; Huang, S.; Lee, J. Y.; Fukase, K.; Inohara, N.; Hwang, D. H. *J Biol Chem* **2007**, *282*, 11618.
- (27) Knight, Z. A.; Shokat, K. M. *Chem Biol* **2005**, *12*, 621.
- (28) Karaman, M. W.; Herrgard, S.; Treiber, D. K.; Gallant, P.; Atteridge, C. E.; Campbell, B. T.; Chan, K. W.; Cicceri, P.; Davis, M. I.; Edeen, P. T.; Faraoni, R.; Floyd, M.; Hunt, J. P.; Lockhart, D. J.; Milanov, Z. V.; Morrison, M. J.; Pallares, G.; Patel, H. K.; Pritchard, S.; Wodicka, L. M.; Zarrinkar, P. P. *Nat Biotechnol* **2008**, *26*, 127.
- (29) Marina-Garcia, N.; Franchi, L.; Kim, Y. G.; Miller, D.; McDonald, C.; Boons, G. J.; Nunez, G. *J Immunol* **2008**, *180*, 4050.

- (30) Lee, J.; Tattoli, I.; Wojtal, K. A.; Vavricka, S. R.; Philpott, D. J.; Girardin, S. E. *J Biol Chem* **2009**, *284*, 23818.
- (31) Ogura, Y.; Inohara, N.; Benito, A.; Chen, F. F.; Yamaoka, S.; Nunez, G. *J Biol Chem* **2001**, *276*, 4812.
- (32) Inohara, N.; Chamaillard, M.; McDonald, C.; Nunez, G. *Annu Rev Biochem* **2005**, *74*, 355.

