



Common genetic variants in the CLDN2 and PRSS1-PRSS2 loci alter risk for alcohol-related and sporadic pancreatitis

Citation

Whitcomb, D. C., J. LaRusch, A. M. Krasinskas, L. Klei, J. P. Smith, R. E. Brand, J. P. Neoptolemos, et al. 2012. "Common genetic variants in the CLDN2 and PRSS1-PRSS2 loci alter risk for alcohol-related and sporadic pancreatitis." *Nature genetics* 44 (12): 1349-1354. doi:10.1038/ng.2466. <http://dx.doi.org/10.1038/ng.2466>.

Published version

<https://doi.org/10.1038/ng.2466>

Link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:11708571>

Terms of use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material (LAA), as set forth at

<https://harvardwiki.atlassian.net/wiki/external/NGY5NDE4ZjgzNTc5NDQzMGIzZWZhMGFIOWI2M2EwYTg>

Accessibility

<https://accessibility.huit.harvard.edu/digital-accessibility-policy>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#)

Published in final edited form as:

Nat Genet. 2012 December ; 44(12): 1349–1354. doi:10.1038/ng.2466.

Common genetic variants in the *CLDN2* and *PRSS1-PRSS2* loci alter risk for alcohol-related and sporadic pancreatitis

David C. Whitcomb^{1,2,3,128}, Jessica LaRusch¹, Alyssa M. Krasinskas⁴, Lambertus Klei⁵, Jill P. Smith⁶, Randall E. Brand¹, John P. Neoptolemos⁷, Markus M. Lerch⁸, Matt Tector⁹, Bimaljit S. Sandhu¹⁰, Nalini M. Guda⁹, Lidiya Orlichenko¹, Alzheimer's Disease Genetics Consortium¹¹, Samer Alkaade¹², Stephen T. Amann¹³, Michelle A. Anderson¹⁴, John Baillie¹⁵, Peter A. Banks¹⁶, Darwin Conwell¹⁶, Gregory A. Coté¹⁷, Peter B. Cotton¹⁸, James DiSario¹⁹, Lindsay A. Farrer^{11,20}, Chris E. Forsmark²¹, Marianne Johnstone⁷, Timothy B. Gardner²², Andres Gelrud¹, William Greenhalf⁷, Jonathan L. Haines^{11,23}, Douglas J. Hartman⁴, Robert A. Hawes¹⁸, Christopher Lawrence¹⁸, Michele Lewis²⁴, Julia Mayerle⁸, Richard Mayeux^{11,25}, Nadine M. Melhem⁵, Mary E. Money²⁶, Thiruvengadam Muniraj²⁷, Georgios I. Papachristou¹, Margaret A. Pericak-Vance^{11,28}, Joseph Romagnuolo¹⁸, Gerard D. Schellenberg^{11,29}, Stuart Sherman¹⁷, Peter Simon⁸, Vijay K. Singh¹, Adam Slivka¹, Donna Stolz², Robert Sutton⁷, Frank Ulrich Weiss⁸, C. Mel Wilcox³⁰, Narcis Octavian Zarnescu¹, Stephen R. Wisniewski³¹, Michael R. O'Connell¹, Michelle L. Kienholz¹, Kathryn Roeder³², M. Michael Barmada³, Dhiraj Yadav¹, Bernie Devlin^{3,5,128}, Marilyn S. Albert³³, Roger L. Albin^{34,35}, Liana G. Apostolova³⁶, Steven E. Arnold³⁷, Clinton T. Baldwin³⁸, Robert Barber³⁹, Lisa L. Barnes^{40,41}, Thomas G. Beach⁴², Gary W. Beecham²⁸, Duane Beekly⁴³, David A. Bennett^{40,44}, Eileen H. Bigio⁴⁵, Thomas D. Bird⁴⁶, Deborah Blacker^{47,48}, Adam Boxer⁴⁹, James R. Burke¹⁵, Joseph D. Buxbaum^{50,51,52}, Nigel J. Cairns⁵³, Laura B. Cantwell²⁹, Chuanhai Cao⁵⁴, Regina M. Carney⁵⁵, Steven L. Carroll⁵⁶, Helena C. Chui⁵⁷, David G. Clark⁵⁸, David H. Cribbs⁵⁹, Elizabeth A. Crocco⁶⁰, Carlos Cruchaga⁶¹, Charles DeCarli⁶², F. Yesim Demirci³, Malcolm Dick⁶³, Dennis W. Dickson⁶⁴, Ranjan Duara⁶⁵, Nilufer Ertekin-Taner^{64,66}, Kelley M. Faber⁶⁷, Kenneth B. Fallon⁵⁶, Martin R. Farlow⁶⁸, Steven Ferris⁶⁹, Tatiana M. Foroud⁶⁷, Matthew P. Frosch⁷⁰, Douglas R. Galasko⁷¹, Mary Ganguli⁵, Marla Gearing^{72,73}, Daniel H. Geschwind⁷⁴, Bernardino Ghetti⁷⁵, John R. Gilbert²⁸, Sid Gilman³⁴, Jonathan D. Glass⁷⁶, Alison M. Goate⁶¹, Neill R. Graff-Radford^{64,66}, Robert C. Green⁷⁷, John H. Growdon⁷⁸, Hakon Hakonarson⁷⁹, Kara L. Hamilton-Nelson²⁸, Ronald L. Hamilton⁸⁰, Lindy E. Harrell⁵⁸, Elizabeth Head⁸¹, Lawrence S. Honig⁸², Christine

¹²⁸Correspondence should be addressed to D.C.W. (whitcomb@pitt.edu) or B.D. (devlinbj@upmc.edu)..

AUTHOR CONTRIBUTIONS

Writing Group: A.M.K., B.D., D.C.W., J.L., L.K., M.L.K.

Project design, management, and coordination: B.D., D.C.W., D.Y., J.P.N., M.M.B., M.M.L., N.M.M., S.R.W.

Sample collection and phenotyping: A.B., A.C.M., A.C.N., A.G., A.G.S., A.I.L., A.J.S., A.K., A.M.G., A.P., A.P.L., A.R., A.S., B.G., B.L.M., B.N.V., B.R., B.S.S., B.T.H., C.A.M., C.B.W., C.Cao, C.Cruchaga, C.D., C.E.F., C.E.Y., C.F.L., C.G.L., C.L., C.M.H., C.M.W., C.R., C.T.B., D.A.B., D.Beekly, D.Blacker, D.C., D.C.Marson, D.C.Mash, D.C.W., D.G.C., D.H.C., D.H.G., D.R.G., D.W.D., D.W.T., D.Y., E.A.C., E.D.R., E.H., E.H.B., E.H.K., E.M., E.M.R., E.P., E.R.M., F.M., F.M.L., F.U.W., F.Y.D., G.A.C., G.A.J., G.I.P., G.J., G.L., G.W.B., H.C.C., H.H., H.J.R., H.P., H.V.V., J.A.K., J.A.Schneider, J.A.Sonnen, J.B., J.B.L., J.C.M., J.C.T., J.D., J.D.B., J.D.G., J.E.P., J.F.Q., J.H.G., J.H.K., J.L.L., J.M., J.M.O., J.M.R., J.P.N., J.P.S., J.P.V., J.Q.T., J.R., J.R.B., J.R.G., J.R.M., J.W., J.W.M., K.A.W.B., K.B.F., K.L.H.N., K.L.L., K.M.F., L.B.C., L.E.H., L.G.A., L.J.V.E., L.L.B., L.S.H., L.S.S., L.S.W., L.W.J., L.Y., M.A.A., M.D., M.E.M., M.Ganguli, M.Gearing, M.I.K., M.J., M.L., M.M., M.M.B., M.M.L., M.P.F., M.R., M.R.F., M.S., M.S.A., M.T., N.E.T., N.J.C., N.M.G., N.R.G.R., N.W.K., O.L.L., O.V., P.A.B., P.B.C., P.K., P.S., R.A.H., R.A.S., R.B., R.C.G., R.C.P., R.D., R.E.B., R.E.T., R.K., R.L.A., R.L.H., R.L.W., R.M.C., R.N.R., R.S., S.A., S.E.A., S.F., S.G., S.G.Y., S.L.C., S.S., S.Spina, S.T.A., S.Weintrub, T.B.G., T.D.B., T.G.B., T.J.M., T.M., T.M.F., V.M.V.D., W.A.K., W.G., W.J.M., W.W.P., W.W.S. Genotyping and expression studies: A.M.K., D.J.H., D.S., G.D.S., J.L., J.L.H., J.P.S., L.A.F., L.O., M.A.P.V., N.O.Z., R.M., V.K.S. Statistical analysis: B.D., D.Y., K.R., L.K., M.M.B., M.R.O

COMPETING FINANCIAL INTEREST

The authors declare no competing financial interests*.

M. Hulette⁸³, Bradley T. Hyman⁷⁸, Gregory A. Jicha⁸⁴, Lee-Way Jin⁸⁵, Gyungah Jun^{38,86,87}, M. Ilyas Kamboh^{3,88}, Anna Karydas⁴⁹, Jeffrey A. Kaye^{89,90}, Ronald Kim⁹¹, Edward H. Koo⁷¹, Neil W. Kowall^{92,93}, Joel H. Kramer⁹⁴, Patricia Kramer^{89,95}, Walter A. Kukull⁹⁶, Frank M. LaFerla⁹⁷, James J. Lah⁷⁶, James B. Leverenz⁹⁸, Allan I. Levey⁷⁶, Ge Li⁹⁹, Chiao-Feng Lin²⁹, Andrew P. Lieberman¹⁰⁰, Oscar L. Lopez⁸⁹, Kathryn L. Lunetta⁸⁷, Constantine G. Lyketsos¹⁰¹, Wendy J. Mack¹⁰², Daniel C. Marson⁵⁸, Eden R. Martin²⁸, Frank Martiniuk¹⁰³, Deborah C. Mash¹⁰⁴, Eliezer Masliah^{71,105}, Ann C. McKee^{92,93}, Marsel Mesulam¹⁰⁶, Bruce L. Miller⁴⁹, Carol A. Miller¹⁰⁷, Joshua W. Miller⁸⁶, Thomas J. Montine⁹⁸, John C. Morris^{53,108}, Jill R. Murrell^{67,75}, Adam C. Naj²⁸, John M. Olichney⁶², Joseph E. Parisi^{109,110}, Elaine Peskind⁹⁹, Ronald C. Petersen⁸⁵, Aimee Pierce⁵⁹, Wayne W. Poon⁶³, Huntington Potter⁵⁴, Joseph F. Quinn⁸⁹, Ashok Raj⁵⁴, Murray Raskind⁹⁹, Eric M. Reiman^{112,113,114}, Barry Reisberg^{69,115}, Christiane Reitz^{25,82}, John M. Ringman³⁶, Erik D. Roberson⁵⁸, Howard J. Rosen⁴⁹, Roger N. Rosenberg¹¹⁶, Mary Sano⁵¹, Andrew J. Saykin^{67,117}, Julie A. Schneider^{40,118}, Lon S. Schneider^{57,119}, William W. Seeley⁴⁹, Amanda G. Smith⁵⁴, Joshua A. Sonnen⁹⁸, Salvatore Spina⁷⁵, Robert A. Stern⁹², Rudolph E. Tanzi⁷⁸, John Q. Trojanowski²⁹, Juan C. Troncoso¹²⁰, Debby W. Tsuang⁹⁹, Otto Valladares²⁹, Vivianna M. Van Deerlin²⁹, Linda J. Van Eldik¹²¹, Badri N. Vardarajan³⁸, Harry V. Vinters^{36,122}, Jean Paul Vonsattel¹²³, Li-San Wang²⁹, Sandra Weintraub¹⁰⁶, Kathleen A. Welsh-Bohmer^{15,124}, Jennifer Williamson⁸², Randall L. Woltjer¹²⁵, Clinton B. Wright¹²⁶, Steven G. Younkin⁶⁴, Chang-En Yu¹²⁷, and Lei Yu⁴⁰

¹Division of Gastroenterology, Hepatology and Nutrition, University of Pittsburgh School of Medicine, Pittsburgh, Pennsylvania, USA.

²Department of Cell Biology and Physiology, University of Pittsburgh School of Medicine, Pittsburgh, Pennsylvania, USA.

³Department of Human Genetics, University of Pittsburgh Graduate School of Public Health, Pittsburgh, Pennsylvania, USA.

⁴Department of Pathology, University of Pittsburgh School of Medicine, Pittsburgh, Pennsylvania, USA.

⁵Department of Psychiatry, University of Pittsburgh School of Medicine, Pittsburgh, Pennsylvania, USA.

⁶Gastroenterology and Liver Diseases, Pennsylvania State University, Hershey, Pennsylvania, USA.

⁷Department of Molecular and Clinical Cancer Medicine, National Institute for Health Research Liverpool Pancreas Biomedical Research Unit, Institute of Translational Medicine, University of Liverpool, Liverpool, UK.

⁸Department of Medicine A, University Medicine Greifswald, Greifswald, Germany.

⁹Aurora Health Care System, St. Luke's Medical Center, Milwaukee, Wisconsin, USA.

¹⁰Division of Gastroenterology, Hepatology and Nutrition, Virginia Commonwealth University Medical Center, Richmond, Virginia, USA.

¹¹A full list of members and affiliations appears at the end of this paper, Alzheimer Disease Genetics Consortium.

¹²Department of Medicine, Saint Louis University, St. Louis, Missouri, USA.

¹³North Mississippi Medical Center, Tupelo, Mississippi, USA.

¹⁴Division of Gastroenterology, University of Michigan, Ann Arbor, Michigan, USA.

¹⁵Department of Medicine, Duke University Medical Center, Durham, North Carolina, USA.

- ¹⁶Division of Gastroenterology, Brigham and Women's Hospital, Boston, Massachusetts, USA.
- ¹⁷Department of Medicine, Indiana University Medical Center, Indianapolis, Indiana, USA.
- ¹⁸Digestive Disease Center, Medical University of South Carolina, Charleston, South Carolina, USA.
- ¹⁹Department of Medicine, University of Utah Health Science Center, Salt Lake City, Utah, USA.
- ²⁰Department of Medicine/Biomedical Genetics Section, Boston University School of Medicine, Boston, Massachusetts, USA.
- ²¹Department of Medicine, University of Florida, Gainesville, Florida, USA.
- ²²Dartmouth-Hitchcock Medical Center, Hanover, New Hampshire, USA.
- ²³Department of Molecular Physiology and Biophysics and Vanderbilt Center for Human Genetics Research, Vanderbilt University, Nashville, Tennessee, USA.
- ²⁴Division of Gastroenterology and Hepatology, Mayo Clinic, Jacksonville, Florida, USA.
- ²⁵Gertrude H. Sergievsky Center and Taub Institute on Alzheimer's Disease and the Aging Brain, Department of Neurology, Columbia University, New York, New York, USA.
- ²⁶Washington Hospital, Hagerstown, Maryland, USA.
- ²⁷Department of Internal Medicine, Yale University and Griffin Hospital, Derby, Connecticut, USA.
- ²⁸The John P. Hussman Institute for Human Genomics and Dr. John T. Macdonald Foundation Department of Human Genetics, University of Miami, Miami, Florida, USA.
- ²⁹Department of Pathology and Laboratory Medicine, University of Pennsylvania School of Medicine, Philadelphia, Pennsylvania, USA.
- ³⁰University of Alabama at Birmingham, Birmingham, Alabama, USA.
- ³¹Department of Epidemiology, Epidemiology Data Center, Graduate School of Public Health, University of Pittsburgh, Pittsburgh, PA USA.
- ³²Department of Statistics, Carnegie Mellon University, Pittsburgh, Pennsylvania, USA.
- ³³Department of Neurology, Johns Hopkins University, Baltimore, Maryland, USA.
- ³⁴Department of Neurology, University of Michigan, Ann Arbor, Michigan, USA.
- ³⁵Geriatric Research, Education and Clinical Center (GRECC), VA Ann Arbor Healthcare System (VAAHS), Ann Arbor, Michigan, USA.
- ³⁶Department of Neurology, University of California Los Angeles, Los Angeles, California, USA.
- ³⁷Department of Psychiatry, University of Pennsylvania Perelman School of Medicine, Philadelphia, Pennsylvania, USA.
- ³⁸Department of Medicine (Genetics Program), Boston University, Boston, Massachusetts, USA.
- ³⁹Department of Pharmacology and Neuroscience, University of North Texas Health Science Center, Fort Worth, Texas, USA.
- ⁴⁰Department of Neurological Sciences, Rush University Medical Center, Chicago, Illinois, USA.
- ⁴¹Department of Behavioral Sciences, Rush University Medical Center, Chicago, Illinois, USA.
- ⁴²Civin Laboratory for Neuropathology, Banner Sun Health Research Institute, Phoenix, Arizona, USA.
- ⁴³National Alzheimer's Coordinating Center, University of Washington, Seattle, Washington, USA.

- ⁴⁴Rush Alzheimer's Disease Center, Rush University Medical Center, Chicago, Illinois, USA.
- ⁴⁵Department of Pathology, Northwestern University, Chicago, Illinois, USA.
- ⁴⁶Department of Neurology, University of Washington, Seattle, Washington, USA.
- ⁴⁷Department of Epidemiology, Harvard School of Public Health, Boston, Massachusetts, USA.
- ⁴⁸Department of Psychiatry, Massachusetts General Hospital/Harvard Medical School, Boston, Massachusetts, USA.
- ⁴⁹Department of Neurology, University of California San Francisco, San Francisco, California, USA.
- ⁵⁰Department of Neuroscience, Mount Sinai School of Medicine, New York, New York, USA.
- ⁵¹Department of Psychiatry, Mount Sinai School of Medicine, New York, New York, USA.
- ⁵²Departments of Genetics and Genomic Sciences, Mount Sinai School of Medicine, New York, New York, USA.
- ⁵³Department of Pathology and Immunology, Washington University, St. Louis, Missouri, USA.
- ⁵⁴USF Health Byrd Alzheimer's Institute, University of South Florida, Tampa, Florida, USA.
- ⁵⁵Department of Psychiatry, Vanderbilt University, Nashville, Tennessee, USA.
- ⁵⁶Department of Pathology, University of Alabama at Birmingham, Birmingham, Alabama, USA.
- ⁵⁷Department of Neurology, University of Southern California, Los Angeles, California, USA.
- ⁵⁸Department of Neurology, University of Alabama at Birmingham, Birmingham, Alabama, USA.
- ⁵⁹Department of Neurology, University of California Irvine, Irvine, California, USA.
- ⁶⁰Department of Psychiatry and Behavioral Sciences, Miller School of Medicine, University of Miami, Miami, Florida, USA.
- ⁶¹Department of Psychiatry and Hope Center Program on Protein Aggregation and Neurodegeneration, Washington University School of Medicine, St. Louis, Missouri, USA.
- ⁶²Department of Neurology, University of California Davis, Sacramento, California, USA.
- ⁶³Institute for Memory Impairments and Neurological Disorders, University of California Irvine, Irvine, California, USA.
- ⁶⁴Department of Neuroscience, Mayo Clinic, Jacksonville, Florida, USA.
- ⁶⁵Wien Center for Alzheimer's Disease and Memory Disorders, Mount Sinai Medical Center, Miami Beach, Florida, USA.
- ⁶⁶Department of Neurology, Mayo Clinic, Jacksonville, Florida, USA.
- ⁶⁷Department of Medical and Molecular Genetics, Indiana University, Indianapolis, Indiana, USA.
- ⁶⁸Department of Neurology, Indiana University, Indianapolis, Indiana, USA.
- ⁶⁹Department of Psychiatry, New York University, New York, New York, USA.
- ⁷⁰C.S. Kubik Laboratory for Neuropathology, Massachusetts General Hospital, Charlestown, Massachusetts, USA.
- ⁷¹Department of Neurosciences, University of California San Diego, La Jolla, California, USA.
- ⁷²Department of Pathology and Laboratory Medicine, Emory University, Atlanta, Georgia, USA.
- ⁷³Emory Alzheimer's Disease Center, Emory University, Atlanta, Georgia, USA.

⁷⁴Neurogenetics Program, University of California Los Angeles, Los Angeles, California, USA.

⁷⁵Department of Pathology and Laboratory Medicine, Indiana University, Indianapolis, Indiana, USA.

⁷⁶Department of Neurology, Emory University, Atlanta, Georgia, USA.

⁷⁷Division of Genetics, Department of Medicine and Partners Center for Personalized Genetic Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, Massachusetts, USA.

⁷⁸Department of Neurology, Massachusetts General Hospital/Harvard Medical School, Boston, Massachusetts, USA.

⁷⁹Center for Applied Genomics, Children's Hospital of Philadelphia, Philadelphia, Pennsylvania, USA.

⁸⁰Department of Pathology (Neuropathology), University of Pittsburgh, Pittsburgh, Pennsylvania, USA.

⁸¹Sanders-Brown Center on Aging, Department of Molecular and Biomedical Pharmacology, University of Kentucky, Lexington, Kentucky, USA.

⁸²Taub Institute on Alzheimer's Disease and the Aging Brain, Department of Neurology, Columbia University, New York, New York, USA.

⁸³Department of Pathology, Duke University, Durham, North Carolina, USA.

⁸⁴Sanders-Brown Center on Aging, Department Neurology, University of Kentucky, Lexington, Kentucky, USA.

⁸⁵Department of Pathology and Laboratory Medicine, University of California Davis, Sacramento, California, USA.

⁸⁶Department of Biostatistics, Boston University, Boston, Massachusetts, USA.

⁸⁷Department of Ophthalmology, Boston University, Boston, Massachusetts, USA.

⁸⁸University of Pittsburgh Alzheimer's Disease Research Center, Pittsburgh, Pennsylvania, USA.

⁸⁹Department of Neurology, Oregon Health & Science University, Portland, Oregon, USA.

⁹⁰Department of Neurology, Portland Veterans Affairs Medical Center, Portland, Oregon, USA.

⁹¹Department of Pathology and Laboratory Medicine, University of California Irvine, Irvine, California, USA.

⁹²Department of Neurology, Boston University, Boston, Massachusetts, USA.

⁹³Department of Pathology, Boston University, Boston, Massachusetts, USA.

⁹⁴Department of Neuropsychology, University of California San Francisco, San Francisco, California, USA.

⁹⁵Department of Molecular & Medical Genetics, Oregon Health & Science University, Portland, Oregon, USA.

⁹⁶Department of Epidemiology, University of Washington, Seattle, Washington, USA.

⁹⁷Department of Neurobiology and Behavior, University of California Irvine, Irvine, California, USA.

⁹⁸Department of Pathology, University of Washington, Seattle, Washington, USA.

- ⁹⁹Department of Psychiatry and Behavioral Sciences, University of Washington, Seattle, Washington, USA.
- ¹⁰⁰Department of Pathology, University of Michigan, Ann Arbor, Michigan, USA.
- ¹⁰¹Department of Psychiatry, Johns Hopkins University, Baltimore, Maryland, USA.
- ¹⁰²Department of Preventive Medicine, University of Southern California, Los Angeles, California, USA.
- ¹⁰³Department of Medicine - Pulmonary, New York University, New York, New York, USA.
- ¹⁰⁴Department of Neurology, University of Miami, Miami, Florida, USA.
- ¹⁰⁵Department of Pathology, University of California San Diego, La Jolla, California, USA.
- ¹⁰⁶Cognitive Neurology and Alzheimer's Disease Center, Northwestern University, Chicago, Illinois, USA.
- ¹⁰⁷Department of Pathology, University of Southern California, Los Angeles, California, USA.
- ¹⁰⁸Department of Neurology, Washington University, St. Louis, Missouri, USA.
- ¹⁰⁹Department of Anatomic Pathology, Mayo Clinic, Rochester, Minnesota, USA.
- ¹¹⁰Department of Laboratory Medicine and Pathology, Mayo Clinic, Rochester, Minnesota, USA.
- ¹¹¹Department of Neurology, Mayo Clinic, Rochester, Minnesota, USA.
- ¹¹²Arizona Alzheimer's Consortium, Phoenix, Arizona, USA.
- ¹¹³Banner Alzheimer's Institute, Phoenix, Arizona, USA.
- ¹¹⁴Neurogenomics Division, Translational Genomics Research Institute, Phoenix, Arizona, USA.
- ¹¹⁵Alzheimer's Disease Center, New York University, New York, New York, USA.
- ¹¹⁶Department of Neurology, University of Texas Southwestern, Dallas, Texas, USA.
- ¹¹⁷Department of Radiology and Imaging Sciences, Indiana University, Indianapolis, Indiana, USA.
- ¹¹⁸Department of Pathology (Neuropathology), Rush University Medical Center, Chicago, Illinois, USA.
- ¹¹⁹Department of Psychiatry, University of Southern California, Los Angeles, California, USA.
- ¹²⁰Department of Pathology, Johns Hopkins University, Baltimore, Maryland, USA.
- ¹²¹Sanders-Brown Center on Aging, Department of Anatomy and Neurobiology, University of Kentucky, Lexington, Kentucky
- ¹²²Department of Pathology & Laboratory Medicine, University of California Los Angeles, Los Angeles, California, USA.
- ¹²³Taub Institute on Alzheimer's Disease and the Aging Brain, Department of Pathology, Columbia University, New York, New York, USA.
- ¹²⁴Department of Psychiatry & Behavioral Sciences, Duke University, Durham, North Carolina, USA.
- ¹²⁵Department of Pathology, Oregon Health & Science University, Portland, Oregon, USA.
- ¹²⁶Evelyn F. McKnight Brain Institute, Department of Neurology, Miller School of Medicine, University of Miami, Miami, Florida, USA.
- ¹²⁷Department of Medicine, University of Washington, Seattle, Washington, USA.

Abstract

Pancreatitis is a complex, progressively destructive inflammatory disorder. Alcohol was long thought to be the primary causative agent, but genetic contributions have been of interest since the discovery that rare *PRSS1*, *CFTR*, and *SPINK1* variants were associated with pancreatitis risk. We now report two significant genome-wide associations identified and replicated at *PRSS1-PRSS2* (1×10^{-12}) and x-linked *CLDN2* ($p < 1 \times 10^{-21}$) through a two-stage genome-wide study (Stage 1, 676 cases and 4507 controls; Stage 2, 910 cases and 4170 controls). The *PRSS1* variant affects susceptibility by altering expression of the primary trypsinogen gene. The *CLDN2* risk allele is associated with atypical localization of claudin-2 in pancreatic acinar cells. The homozygous (or hemizygous male) *CLDN2* genotype confers the greatest risk, and its alleles interact with alcohol consumption to amplify risk. These results could partially explain the high frequency of alcohol-related pancreatitis in men – male hemizygous frequency is 0.26, female homozygote is 0.07.

The exocrine pancreas is a simple digestive gland of only two primary cell types, each with a single function (Supplementary Figure 1). Recurrent acute pancreatic inflammation can, but does not always, progress to irreversible damage of the gland, including fibrosis, atrophy, pain, and exocrine and endocrine insufficiency,¹⁻³ known as chronic pancreatitis. Different genetic and environmental factors produce the same clinical phenotype⁴.

We collected biological samples and phenotypic data from 1000 patients with recurrent acute pancreatitis and chronic pancreatitis plus controls in the North American Pancreatitis Study 2 (NAPS2)⁵. The primary environmental risk factor identified was heavy alcohol drinking when symptoms of pancreatitis began, based on the assessment of the study physician, called here alcohol-related pancreatitis.

To further define genetic risk, we conducted a two-stage (discovery/replication) genomewide association study (GWAS). The final data set for the Stage 1 cohort included 676 chronic pancreatitis cases and 4507 controls of European ancestry (Supplementary Figs. 2-3) genotyped at 625,739 SNPs (Table 1; Supplementary Table 1). Genomewide significant associations ($p\text{-value} < 5 \times 10^{-8}$) were identified at two loci. The most highly associated SNP fell in Xq23.3, dubbed the *CLDN2* locus, the other in 7q34, the *PRSS1-PRSS2* locus (Fig. 1; Table 2; Supplementary Figs. 4-5, Supplementary Table 2). *CLDN2* encodes the protein claudin-2, while *PRSS1* encodes cationic trypsinogen, and *PRSS2* encodes anionic trypsinogen.

The Stage 2 cohort included 910 cases (331 chronic pancreatitis, 579 recurrent acute pancreatitis; Table 1, Supplementary Table 1), again genotyped at 625,739 SNPs, and 4170 controls, most genotyped previously on the Illumina 1M. All subjects were of European ancestry as determined by genetic analyses. Recurrent acute pancreatitis and chronic pancreatitis were modeled as having common susceptibilities, with chronic pancreatitis occurring over time in the presence of additional disease-modifying factors.⁶ It is possible that this assumption reduces power relative to a study comprising solely chronic pancreatitis or recurrent acute pancreatitis cases. Our primary targets in Stage 2 were the *PRSS1-PRSS2* and *CLDN2* loci, although we also conducted a joint analysis⁷ of Stage 1 and Stage 2 data to uncover any new risk loci. After controlling for ancestry, these data demonstrated significant effects for the *CLDN2* and *PRSS1-PRSS2* loci (Figure 1; Supplementary Table 2-3; Supplementary Figs. 6-7). Quality of SNP genotypes supported the association (Supplementary Fig. 8). The frequencies of the putative risk alleles at these 2 loci were 0.57 for the C allele at rs10273639 (*PRSS1-PRSS2* locus), with the minor T allele reducing risk, and 0.26 for the T allele at rs12688220 (*CLDN2* locus). No other locus shows association after accounting for SNP genotype quality (Supplementary Figs. 6-8).

PRSS1 gain-of-function mutations, such as p.R122H, increase risk for recurrent acute pancreatitis and chronic pancreatitis⁸, as do increased copy number^{9,10}. Rare loss-of-function mutations in *PRSS2* are protective¹¹. However, rs10273639 is in the 5' promoter region of *PRSS1*. Because it is the only highly associated SNP in the locus, we validated its genotypes by independent TaqMan genotyping and also genotyped two SNPs in linkage disequilibrium with it (footnote, Supplementary Table 4)^{12,13}. We screened *PRSS1* for rare variants in 1138 subjects: 418 chronic pancreatitis, 350 recurrent acute pancreatitis, and 379 controls. Three known disease-associated variants (A16V, N29I, R122H) were identified in 23 subjects (Supplementary Table 4). These gain-of-function variants occur almost solely in cases (22 out of 23), and two of them, A16V and R122H, likely fall on the C or risk haplotype of this locus (Supplementary Table 4). Nonetheless, with only 19 A16V and R122H events in cases, these rare alleles cannot account for the association observed at this locus.

Sixty-nine control pancreas tissue samples from three sources were genotyped at rs10273639, and cDNA was used to quantify *PRSS1* and control gene expression (Supplementary Table 5). For all three sets of quantitative PCR data, the slope relating count of genotype C allele to *PRSS1* expression level was positive; together, the samples provide evidence ($p = 0.01$) that alleles at rs10273639 affect expression of *PRSS1*: expression levels were highest in patients with two C alleles at rs10273639, intermediate in heterozygotes, and lowest in subjects with two T alleles. Based on this evidence, we posit that reduced trypsinogen production protects the pancreas from injury, as has been observed in genetic mouse models¹⁴.

CLDN2 is considered the primary candidate gene within our *CLDN2* locus. Claudin-2 is attractive because it serves as a highly regulated tight junction protein forming low-resistance, cation-selective ion and water channels between endothelial cells^{15,16} and is normally expressed at low levels between cells of the pancreatic ducts and in pancreatic islets^{17,18}. The *CLDN2* promoter includes an NF κ B binding site¹⁹, and gene expression is enhanced in other cells under conditions associated with injury or stress²⁰⁻²². Claudin-2 can also be expressed by acinar cells when stressed, as reported in porcine models of acute pancreatitis²³. Other genes within the *CLDN2* locus include *MORC4*, *RIPPLY1*, and *TBC1D8B*. *MORC4* is expressed at low levels in most tissues, including the pancreas, with higher levels in the placenta and testis²⁴. The *MORC4* protein contains a CW four-cysteine zinc-finger motif, nuclear localization signal, and nuclear matrix-binding domain, suggesting that it may be a transcription factor²⁴, but its expression does not appear to correlate with pancreatitis (Supplementary Fig. 9). *RIPPLY1* and *TBC1D8B* are not known to be expressed in the pancreas.

To our knowledge, genetic variations in *CLDN2* have never been associated with disease in humans. We assessed DNA sequence variants around *CLDN2*, RNA, and protein expression for claudin-2 in control tissue classified by histology and genotype (Supplementary Table 6, Supplementary Fig. 10). Evaluating 1000 Genomes data, no exonic variation was identified that could explain the association signal. Using materials and methods described previously for *PRSS1* expression, *CLDN2* expression levels in control tissues did not correlate with the *CLDN2* locus risk genotype (p -value = 0.32). Protein was extracted from the tissue, and only one protein band of the appropriate size was observed with anti-claudin-2 antibodies on Western blot, which correlated with tissue inflammation as determined by systematic grading of histology in adjacent tissue (Fig. 2A, Supplementary Fig. 10). Immunohistochemical staining with anti-claudin-2 antibodies was verified in normal tissue (Fig. 2B), with kidney, duodenum, and bile ducts serving as additional positive controls (not shown). Protein localization was assessed in 12 GWAS cases who underwent pancreatic surgery: 6 with the *CLDN2*-containing high-risk genotype and 6 without. Claudin-2

cytoplasmic granular staining was markedly increased in both duct and acinar cells in chronic pancreatitis cases (Fig. 2C-E). Only chronic pancreatitis cases with the high-risk *CLDN2* genotype demonstrated moderate-to-strong claudin-2 staining along the basolateral membrane of acinar cells (Fig. 2D, 2E, Supplementary Table 6). Claudin-2 was also expressed in macrophages, which could contribute to the pathologic inflammatory process²⁵ (Fig. 2C, F).

Most studies report excessive alcohol consumption as the major risk factor for adult-onset chronic pancreatitis²⁶⁻²⁹. However, only 3% of patients who are alcoholics develop chronic pancreatitis³⁰, suggesting a pancreas-targeting risk factor. We compared genotypes based on whether pancreatitis was alcohol-related (yes/no)^{5,31}. Setting control genotypes counts as the baseline category to be compared with case genotypes, the jointly estimated odds ratios for cases with a positive alcohol-related pancreatitis was greater for both rs10273639 (*PRSSI-PRSS2* locus) and rs12688220 (*CLDN2* locus) than those estimated for cases with a negative alcohol-related pancreatitis (Table 3). Thus, the effects of both loci appeared to be amplified by alcohol consumption. In a *case-only analysis*, both loci appear to interact with alcohol-related pancreatitis (Table 3), the *CLDN2* locus most prominently (p-value = 4×10^{-7}).

We conclude that a common allele in the *PRSSI-PRSS2* locus is associated with lower *PRSSI* gene expression and that this effect is independent of the previously reported rare gain-of-function *PRSSI* variants that increase susceptibility to both recurrent acute pancreatitis and chronic pancreatitis⁸. For this reason, and because risk variants at the *PRSSI-PRSS2* locus exert a similar effect in patients with recurrent acute pancreatitis or chronic pancreatitis, it is reasonable to conjecture that variation at rs10273639 or variation in linkage disequilibrium with it directly affects risk for chronic pancreatitis and recurrent acute pancreatitis through its impact on trypsinogen expression. Variation at the *CLDN2* locus, however, is much more strongly associated with chronic pancreatitis than recurrent acute pancreatitis, suggesting that it likely acts as a disease modifier to accelerate transition from recurrent acute pancreatitis to chronic pancreatitis. The significant association of the *CLDN2* locus with alcohol suggests that the high-risk allele in the *CLDN2* locus may modify risk through a non-trypsin-dependent process. Thus, we have characterized two common genetic risk modifiers for sporadic and alcohol-related chronic pancreatitis.

Online Methods

Subject recruitment

Details of recruitment of cases and controls are reported in Supplementary Table 1. All studies were conducted under institutional review board-approved protocols.

Stage 1 samples

All N = 758 Stage 1 case samples were from the North American Acute Pancreatitis Study (NAPS2⁵) were diagnosed with chronic pancreatitis, and were characterized for alcohol-related pancreatitis (Table 1). chronic pancreatitis occurs in less than 0.05% of the population, so a convenience sample provides essentially identical power as a same-sized sample of controls selected for the absence of chronic pancreatitis³². For controls, we used genotypes from 4076 cases and controls from the Alzheimer Disease Genetics Consortium (ADGC) and 493 NAPS2 subjects, all genotyped on the same platform as the chronic pancreatitis samples.

Stage 2 samples

The Stage 2 samples consisted of N=343 chronic pancreatitis and N=627 recurrent acute pancreatitis cases (Table 1, Supplementary Table 1) as well as 4191 control subjects (3986 from the NeuroGenetics Research Consortium, NGRC, and 205 NAPS2 controls).

Genotypes

All cases and NAPS2 controls were genotyped by the University of Pittsburgh Genomics and Proteomics Core Laboratories using the Illumina HumanOmniExpress Beadchip. Samples were processed and scanned using the manufacturer's recommended protocols with no modifications. ADGC samples³³ were also genotyped using Illumina HumanOmniExpress Beadchips, whereas NGRC samples³⁴ were genotyped on the Illumina Human1M-Duo DNA Analysis BeadChip.

Quality Control (QC) for Stage 1

QC was performed for individuals and then SNPs to determine which samples and SNPs should not be included in the analysis ("dropped"). Assessing sex miscalls based on \times chromosome genotypes using Plink³⁵, 7 chronic pancreatitis cases and 20 controls (10 NAPS2; 10 ADGC) were dropped. Based on the requirement for 95% complete genotypes per individual, 40 cases and 27 controls (20 NAPS2 controls and 7 ADGC controls) were dropped. Searching duplicate or highly related samples based on genotype and using GCTA software³⁶ (Genetic Relationship Matrix score GRM > 0.4), 35 cases and 78 controls (2 NAPS2, 76 ADGC) were dropped. After these QC filters, 676 cases and 4507 controls remained for association analysis.

SNP QC was first performed using NAPS2 and ADGC samples separately. Ancestry was estimated using dacGem³⁷ based on 9700 SNPs that had a genotype completion rate of 99.9%, a minor allele frequency MAF \geq 0.05, and were separated by at least 500Kb. Analysis of genotypes from NAPS2 subjects identified 1 significant dimensions of ancestry and clustered subjects into 3 groups (Supplementary Fig. 1). Groups A and B, illustrated in Supplementary Figure 1, delineate 764 and 282 subjects, respectively, of European ancestry (self-identified); SNP QC for MAF and Hardy Weinberg Equilibrium (HWE) were performed on data from these subjects. Of 731,442 SNPs received, 633,790 passed QC filters. SNPs were dropped for the following reasons: 3165 for map location; 11,977 for call rate; 77,300 for MAF < 0.01; and 5219 failed HWE (p-value < 0.005).

ADGC data were received in three waves of 1763, 1110, and 1266 subjects. In the first wave, 659,224 SNPs were received, while in waves two and three, 730,525 SNPs were received. After QC as described for the chronic pancreatitis cohort, including harmonization with SNPs passing QC in the chronic pancreatitis cohort, 604,059, 632,761, and 633,023 SNPs remained, respectively. After merging cohorts, 30 related subjects were dropped, leaving 4046 ADGC subjects. Of the 633,615 unique SNPs in this ADGC, QC filters dropped 5 for low MAF and 5316 for HWE, leaving 628,294 SNPs. Combining ADGC and chronic pancreatitis cohorts and performing another round of QC yielded 625,739 SNPs for analysis.

QC for Stage 2

QC for individuals was performed as described for Stage 1. These individual-specific QC filters removed 60 cases, leaving 331 chronic pancreatitis and 579 recurrent acute pancreatitis cases for analysis; 14 controls were also removed, leaving 4177 controls for analysis. We analyzed all SNPs passing QC at Stage 1.

Association analysis

To control confounding due to ancestry, the first 10 major eigenvectors from the spectral decomposition were used as covariates in Stage 1 and Stage 2 analyses³⁸, although only one was significant. We contrasted the genotypes of case subjects and controls via logistic regression and a log-additive (logit) model using Plink³⁵. Genotypes for any SNPs showing association p-value $< 5 \times 10^{-7}$ were manually inspected for valid genotype clustering. SNPs showing poor-quality clustering were excluded. Following Skol *et al.*⁷ and others, we take an overall significance level of 5×10^{-8} and 5×10^{-7} for strongly suggestive association.

To determine whether alcohol interacts with genetic variation to alter risk of pancreatitis, data from cases were fit to a general linear model in which count of alleles or genotypes predicted alcohol etiology (yes/no). The test statistic was obtained as a likelihood ratio chi-square. Note that in these analyses and any analyses other than genomewide association, we model the male genotypes as 0 and 2^{39,40}. For the genomewide association, Plink encodes the count of minor alleles in males as 0 and 1 and includes a sex effect, but the 0/2 encoding for males is a more powerful approach^{39,40}.

DNA extraction

DNA was obtained using standard methods⁴¹.

Pancreatic tissue processing

Tissue was obtained from two sources [Pitt and Pancreatic Adenocarcinoma Gene-Environment Risk (PAGER) from the University of Pittsburgh and PSU from Pennsylvania State University] and processed in three batches: banked (Pitt) and prospectively collected (PAGER) surgical waste from uninvolved pancreas and normal pancreas specimens from the Gift of Life Program that were not used for transplantation (PSU). PAGER samples were snap-frozen, placed in RNAlater solution (Ambion), and stored at -80°C . PSU pancreas samples were also snap frozen and preserved in formalin or placed in RNAlater solution. RNA was isolated using Trizol reagent (Invitrogen), and its quality examined in 1% agarose gel stained with ethidium bromide. cDNA was transcribed using oligo dt primers and the Superscript II reverse transcriptase kit (Invitrogen).

Gene Expression

Relative expression of *PRSS1*, *PRSS2*, *CTRC*, and *18S* was determined by analyzing cDNA using Taqman®-based rtPCR assays (Applied Biosystems). Raw absolute quantitation results were analyzed and converted to relative expression results by software packages SDS V2.3 and DataAssist V1.0 (Applied Biosystems). Assays were repeated in triplicate or quadruplicate. Three sets of samples were assessed, two from Pitt (N=10 and 22) and one from PSU (N=37). PSU results were normalized against 18S, Pitt against CTCR. From each of these three data sets, mean gene expression per sample was regressed against allele count to obtain an estimated slope, standard error, and z-score. We then calculated an overall z-score as a weighted average of the individual z-scores, with weights determined by sample size.

Antibodies

Antibodies against claudin proteins (Invitrogen) were assessed using Western blot for mouse anti-claudin-2 (Catalog No. 32-5600), mouse anti-claudin-4 (Catalog No. 32-9400), and mouse alpha-tubulin antibody (Catalog no. AA12.1 The Developmental Studies Hybridoma Bank at the University of Iowa, <http://dshb.biology.uiowa.edu/Antibody-list>). Immunohistochemistry was performed using monoclonal antibodies for claudin-2 (Catalog #32-5600, 1:1,000 dilution). Immunofluorescence was performed using mouse anti-claudin-2

(Catalog No. 32-5600) and goat anti-human CD68 (Catalog #sc-7082, Santa Cruz Biotechnology Inc.). The secondary antibodies for Immunofluorescence were goat anti-mouse CY3 and anti goat Cy5 from Jackson Immunoresearch.

SDS-PAGE and WESTERN Blotting

Protein homogenates for Western blotting were obtained from snap-frozen tissue that was homogenized and sonicated in lysis buffer supplemented with protease inhibitors. Protein concentration was determined by the Bradford method using a kit from Bio-Rad. Proteins were separated on 12% SDS-PAGE⁴² followed by transfer to polyvinylidene difluoride (PVDF) membranes⁴³, for Western blotting⁴⁴. Immunodetection of bound antibodies on PVDF membrane was performed using ECL reagents (Amersham Biosciences). All procedures were carried out according to manufacturer instructions.

Immunohistochemistry

Standard automated immunohistochemistry was performed for claudin 2 (antibodies listed above) on formalin-fixed, paraffin-embedded, 5 micron-thick tissue sections. Following deparaffinization in xylene and rehydration in ethanol, antigen retrieval was performed using EDTA pH8 buffer. The Dako Autostainer Plus was used; the slides were incubated for 30 minutes with the primary antibodies, followed by incubation with the secondary reagent (Mach 2 Mouse HRP Polymer from Biocare Medical) for 30 minutes. The chromogen was developed (Dako DAB+) for 10 minutes. The immunohistochemical stains were reviewed by one of the authors (A.M.K.). Cytoplasmic, granular, and membranous staining, predominantly in the lateral cell membranes, were graded on an intensity scale of 0-4 (0, negative; 1, weak; 2, moderate; 3, strong). The staining intensity was very patchy from lobule to lobule in most cases.

Immunofluorescence

Cryostat sections (5 micron) of pancreas were washed 3 times with phosphate-buffered saline (PBS), followed by 3 washes with solution of .5% BSA in PBS. Sections were blocked with 2% BSA solution for 30 minutes. The slides were incubated for 1 hour at room temperature with primary antibody for claudin-2 1:100 and goat anti-human CD68 in 0.5% BSA solution. Slides were washed 3 times with BSA solution and incubated for 1 hour at 20°C with 1:500 dilution anti-goat CY5 and 1:1000 dilution goat anti-mouse CY3 secondary antibodies in BSA solution. Nuclei were stained with Hoeschts dye (bisbenzamide 1mg/100ml water) for 30 seconds. After 3 rinses with PBS, sections were cover slipped with Gelvatol mounting media. Fluorescent images were captured with an Olympus Fluoview 1000 confocal microscope (software version 1.7a). The Cy5 signal (CD68) was pseudocolored as green to show colocalization with the red Claudin signal as yellow.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Technical support was provided by Kimberly Stello, Siddhartha Das, Danielle Dwyer, Alexander Rowland, Paul Alexander Blake, Mark Ross, Liz Kish PhD, and Haq Nawaz MD, Shiela Solomon MS CGC and Sharon Boggiano RN from the University of Pittsburgh; Rachel Ostroff, Meredith Goss and Jan Timm from SomaLogic Inc (Boulder CO). Data management by Laurie Silfies and Darina Protivnak, Epidemiology Data Center, Graduate School of Public Health, University of Pittsburgh. Clinical support: Sharon Boggaino RN, Megan Hendricks RN, Beth Elinoff RN MPH CCRC, University of Pittsburgh, Lee McHenry MD, Glen Lehman MD, James Watkins MD, Evan Fogel MD, Laura Lazzell-Pannell, BSN, RN, CCRC, Indiana University, Indianapolis, IN. Additional samples were provided by Frank Burton MD (deceased) Department of Internal Medicine, St. Louis University School of Medicine, St Louis, MO, Simon Lo MD, Department of Medicine, Cedars-Sinai Medical Center, University of

California, Los Angeles; Mark T. DeMeo MD, Department of Medicine, Rush University Medical Center, Chicago, IL; William M. Steinberg MD, Washington Hospital Center, Washington DC; Michael L. Kochman MD, Department of Medicine, University of Pennsylvania, Philadelphia, PA; Babak Etemad MD, Department of Gastroenterology and Hepatology, Ochsner Medical Center, New Orleans, LA. Herbert Zeh MD, Arthur James Moser MD, K. Kenneth Lee MD, Department of Surgery, University of Pittsburgh, Pittsburgh PA.

This publication was made possible by Grant Numbers DK061451 (DCW), DK054709 (DCW), DK063922 (DCW), MH057881 (BD/KR), CA117926 (JPS), UL1 RR024153 and UL1TR000005. Its contents are solely the responsibility of the authors and do not necessarily represent the official view of the NIH. This project used the University of Pittsburgh Genomics and Proteomics Core Laboratories (UL1 RR024153) and the UPCI Clinical Genomics Immunoproteomics and Sequencing Facility (NIH P30CA047904). Jessica LaRusch was supported by Digestive Disease Training Program T32DK063922 (DCW), Narsis Zarnescu MD was supported by the American Gastroenterology Association John I Isenberg MD International Scholar Award. The Liverpool cohort was supported by NIHR Biomedical Research Unit award. Additional support was provided by National Pancreas Foundation (DCW), the Frieda G. and Saul F. Shapira BRCA Cancer Research Program (DCW) and the Wayne Fusaro Pancreatic Cancer Research Fund (DCW) and Gift of Life Foundation

Support for control cohort 1. Alzheimer's Disease Genetics Consortium: The National Institutes of Health, National Institute on Aging (NIH-NIA) supported this work through the following grants: ADGC, U01 AG032984, RC2 AG036528; NACC, U01 AG016976; NCRAD, U24 AG021886; Banner Sun Health Research Institute P30 AG019610; Boston University, P30 AG013846, U01 AG10483, R01 CA129769, R01 MH080295, R01 AG017173, R01 AG025259, R01AG33193; Columbia University, P50 AG008702, R37 AG015473; Duke University, P30 AG028377, AG05128; Emory University, AG025688; Indiana University, P30 AG10133; Johns Hopkins University, P50 AG005146, R01 AG020688; Massachusetts General Hospital, P50 AG005134; Mayo Clinic, P50 AG016574; Mount Sinai School of Medicine, P50 AG005138, P01 AG002219; New York University, P30 AG08051, MO1RR00096, and UL1 RR029893; Northwestern University, P30 AG013854; Oregon Health & Science University, P30 AG008017, R01 AG026916; Rush University, P30 AG010161, R01 AG019085, R01 AG15819, R01 AG17917, R01 AG30146; University of Alabama at Birmingham, P50 AG016582, UL1RR02777; University of Arizona, R01 AG031581; University of California, Davis, P30 AG010129; University of California, Irvine, P50 AG016573, P50, P50 AG016575, P50 AG016576, P50 AG016577; University of California, Los Angeles, P50 AG016570; University of California, San Diego, P50 AG005131; University of California, San Francisco, P50 AG023501, P01 AG019724; University of Kentucky, P30 AG028383, AG05144; University of Michigan, P50 AG008671; University of Pennsylvania, P30 AG010124; University of Pittsburgh, P50 AG005133, AG030653; University of Southern California, P50 AG005142; University of Texas Southwestern, P30 AG012300; University of Miami, R01 AG027944, AG010491, AG027944, AG021547, AG019757; University of Washington, P50 AG005136; Vanderbilt University, R01 AG019085; and Washington University, P50 AG005681, P01 AG03991. The Kathleen Price Bryan Brain Bank at Duke University Medical Center is funded by NINDS grant # NS39764, NIMH MH60451 and by Glaxo Smith Kline. We thank Drs. D. Stephen Snyder and Marilyn Miller from NIA who are *ex-officio* ADGC members.

Support for control group 2. Data for control group 2 genotypes were obtained from Genome-Wide Association Study of Parkinson Disease: Genes and Environment, dbGaP Study Accession phs000196.v2.p1. This NeuroGenetics Research Consortium (NGRC) is a gene-environment study of Parkinson's disease. Principal Investigator is Haydeh Payami, PhD, New York State Department of Health Wadsworth Center, Albany, NY, USA. Co-investigators are: John Nutt, MD., Oregon Health & Sciences University; Cyrus Zabetian, MS, MD. University of Washington and Puget Sound Veterans Medical Center, Seattle, WA, USA; Stewart Factor, DO. Emory University, Atlanta, GA, USA; Eric Molho, MD. Albany Medical Center, Albany, NY, USA; and Donald Higgins, MD. Albany Medical Center and Albany Veterans Medical Center Albany, NY, USA. Funding Source: 5R01NS36960-10. Genotyping provided by Genotyping Center, Johns Hopkins University Center for Inherited Disease Research (CIDR), Baltimore, MD, USA. Funding Source for Genotyping: HHSN268200782096C. NIH contract "High throughput genotyping for studying the genetic contributions to human disease" from the National Institutes of Health, Bethesda, MD, USA.

References

1. Etemad B, Whitcomb DC. Chronic pancreatitis: Diagnosis, classification, and new genetic developments. *Gastroenterology*. 2001; 120:682–707. [PubMed: 11179244]
2. Chen JM, Ferec C. Chronic pancreatitis: genetics and pathogenesis. *Annu Rev Genomics Hum Genet*. 2009; 10:63–87. [PubMed: 19453252]
3. Witt H, Apte MV, Keim V, Wilson JS. Chronic pancreatitis: challenges and advances in pathogenesis, genetics, diagnosis, and therapy. *Gastroenterology*. 2007; 132:1557–73. [PubMed: 17466744]
4. Whitcomb DC. What is personalized medicine - what does it replace? *Nat Rev Gastroenterol Hepatol*. 2012

5. Whitcomb DC, et al. Multicenter approach to recurrent acute and chronic pancreatitis in the United States: the North American Pancreatitis Study 2 (NAPS2). *Pancreatol.* 2008; 8:520–31. [PubMed: 18765957]
6. Yadav D, Whitcomb DC. The role of alcohol and smoking in pancreatitis. *Nat Rev Gastroenterol Hepatol.* 2010; 7:131–45. [PubMed: 20125091]
7. Skol AD, Scott LJ, Abecasis GR, Boehnke M. Joint analysis is more efficient than replication-based analysis for two-stage genome-wide association studies. *Nature genetics.* 2006; 38:209–13. [PubMed: 16415888]
8. Whitcomb DC, et al. Hereditary pancreatitis is caused by a mutation in the cationic trypsinogen gene. *Nature Genetics.* 1996; 14:141–5. [PubMed: 8841182]
9. Masson E, et al. Trypsinogen copy number mutations in patients with idiopathic chronic pancreatitis. *Clin Gastroenterol Hepatol.* 2008; 6:82–8. [PubMed: 18063422]
10. Larusch J, Barmada MM, Solomon S, Whitcomb DC. Whole exome sequencing identifies multiple, complex etiologies in an idiopathic hereditary pancreatitis kindred. *JOP : Journal of the pancreas.* 2012; 13:258–62. [PubMed: 22572128]
11. Witt H, et al. A degradation-sensitive anionic trypsinogen (PRSS2) variant protects against chronic pancreatitis. *Nat Genet.* 2006; 38:668–73. [PubMed: 16699518]
12. A map of human genome variation from population-scale sequencing. *Nature.* 2010; 467:1061–73. [PubMed: 20981092]
13. Johnson AD, et al. SNAP: a web-based tool for identification and annotation of proxy SNPs using HapMap. *Bioinformatics.* 2008; 24:2938–9. [PubMed: 18974171]
14. Dawra R, et al. Intra-acinar trypsinogen activation mediates early stages of pancreatic injury but not inflammation in mice with acute pancreatitis. *Gastroenterology.* 2011; 141:2210–2217. e2. [PubMed: 21875495]
15. Van Itallie CM, et al. The density of small tight junction pores varies among cell types and is increased by expression of claudin-2. *Journal of cell science.* 2008; 121:298–305. [PubMed: 18198187]
16. Amasheh S, et al. Claudin-2 expression induces cation-selective channels in tight junctions of epithelial cells. *Journal of cell science.* 2002; 115:4969–76. [PubMed: 12432083]
17. Lee JH, et al. Immunohistochemical analysis of claudin expression in pancreatic cystic tumors. *Oncology reports.* 2011; 25:971–8. [PubMed: 21206985]
18. Aung PP, et al. Differential expression of claudin-2 in normal human tissues and gastrointestinal carcinomas. *Virchows Archiv : an international journal of pathology.* 2006; 448:428–34. [PubMed: 16328347]
19. Sakaguchi T, et al. Cloning of the human claudin-2 5'-flanking region revealed a TATA-less promoter with conserved binding sites in mouse and human for caudal-related homeodomain proteins and hepatocyte nuclear factor-1alpha. *The Journal of biological chemistry.* 2002; 277:21361–70. [PubMed: 11934881]
20. Mankertz J, et al. TNFalpha up-regulates claudin-2 expression in epithelial HT-29/B6 cells via phosphatidylinositol-3-kinase signaling. *Cell and tissue research.* 2009; 336:67–77. [PubMed: 19214581]
21. Suzuki T, Yoshinaga N, Tanabe S. Interleukin-6 (IL-6) regulates claudin-2 expression and tight junction permeability in intestinal epithelium. *The Journal of biological chemistry.* 2011; 286:31263–71. [PubMed: 21771795]
22. Mankertz J, et al. Functional crosstalk between Wnt signaling and Cdx-related transcriptional activation in the regulation of the claudin-2 promoter activity. *Biochemical and biophysical research communications.* 2004; 314:1001–7. [PubMed: 14751232]
23. Merilainen S, et al. Acute edematous and necrotic pancreatitis in a porcine model. *Scandinavian journal of gastroenterology.* 2008; 43:1259–68. [PubMed: 18609130]
24. Liggins AP, et al. MORC4, a novel member of the MORC family, is highly expressed in a subset of diffuse large B-cell lymphomas. *British journal of haematology.* 2007; 138:479–86. [PubMed: 17608765]

25. Van den Bossche J, et al. Claudin-1, claudin-2 and claudin-11 genes differentially associate with distinct types of anti-inflammatory macrophages in vitro and with parasite- and tumor-elicited macrophages in vivo. *Scandinavian journal of immunology*. 2012
26. Ammann RW, Akovbiantz A, Largiader F, Schueler G. Course and outcome of chronic pancreatitis. Longitudinal study of a mixed medical-surgical series of 245 patients. *Gastroenterology*. 1984; 86:820–8. [PubMed: 6706066]
27. Marks IN, Bank S, Louw JH. Chronic pancreatitis in the Western Cape. *Digestion*. 1973; 9:447–53. [PubMed: 4784706]
28. Robles-Diaz G, Vargas F, Uscanga L, Fernandez-del Castillo C. Chronic pancreatitis in Mexico City. *Pancreas*. 1990; 5:479–83. [PubMed: 2381902]
29. Irving HM, Samokhvalov AV, Rehm J. Alcohol as a risk factor for pancreatitis. A systematic review and meta-analysis. *JOP*. 2009; 10:387–92. [PubMed: 19581740]
30. Yadav D, Eigenbrodt ML, Briggs MJ, Williams DK, Wiseman EJ. Pancreatitis: prevalence and risk factors among male veterans in a detoxification program. *Pancreas*. 2007; 34:390–8. [PubMed: 17446836]
31. Yadav D, et al. Alcohol consumption, cigarette smoking, and the risk of recurrent acute and chronic pancreatitis. *Arch Intern Med*. 2009; 169:1035–45. [PubMed: 19506173]
32. Bacanu SA, Devlin B, Roeder K. The power of genomic control. *American journal of human genetics*. 2000; 66:1933–44. [PubMed: 10801388]
33. Naj AC, et al. Common variants at MS4A4/MS4A6E, CD2AP, CD33 and EPHA1 are associated with late-onset Alzheimer's disease. *Nature genetics*. 2011; 43:436–41. [PubMed: 21460841]
34. Hamza TH, et al. Genome-wide gene-environment study identifies glutamate receptor gene GRIN2A as a Parkinson's disease modifier gene via interaction with coffee. *PLoS genetics*. 2011; 7:e1002237. [PubMed: 21876681]
35. Purcell S, et al. PLINK: a tool set for whole-genome association and populationbased linkage analyses. *American journal of human genetics*. 2007; 81:559–75. [PubMed: 17701901]
36. Yang J, Lee SH, Goddard ME, Visscher PM. GCTA: a tool for genome-wide complex trait analysis. *American journal of human genetics*. 2011; 88:76–82. [PubMed: 21167468]
37. Klei L, Kent BP, Melhem N, Devlin B, Roeder K. GemTools: a fast and efficient approach to estimating genetic ancestry. 2011
38. Price AL, et al. Principal components analysis corrects for stratification in genome-wide association studies. *Nature genetics*. 2006; 38:904–9. [PubMed: 16862161]
39. Clayton DG. Sex chromosomes and genetic association studies. *Genome medicine*. 2009; 1:110. [PubMed: 19939292]
40. Zheng G, Joo J, Zhang C, Geller NL. Testing association for markers on the X chromosome. *Genetic epidemiology*. 2007; 31:834–43. [PubMed: 17549761]
41. Pfitzer RH, et al. SPINK1/PSTI polymorphisms act as disease modifiers in familial and idiopathic chronic pancreatitis. *Gastroenterology*. 2000; 119:615–23. [PubMed: 10982753]
42. Laemmli UK. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature*. 1970; 227:680–5. [PubMed: 5432063]
43. Towbin H, Staehelin T, Gordon J. Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets: procedure and some applications. *Proceedings of the National Academy of Sciences of the United States of America*. 1979; 76:4350–4. [PubMed: 388439]
44. Burnette WN. “Western blotting”: electrophoretic transfer of proteins from sodium dodecyl sulfate--polyacrylamide gels to unmodified nitrocellulose and radiographic detection with antibody and radioiodinated protein A. *Analytical biochemistry*. 1981; 112:195–203. [PubMed: 6266278]




Figure 1. Manhattan plot showing the negative log (base 10) of the p-value for the association of SNP genotype with affection status for all SNPs passing quality control filters and falling within a selected region of the *PRSS1-PRSS2* and *CLDN2* loci. Regions selected to highlight the most associated SNPs. Squares indicate Stage 1 results, circles for Stage 2, diamonds for combined Stage 1 and 2 data. After accounting for the most highly associated SNP at each locus, no other SNP approached genomewide-significant association.




Figure 2.

Expression and localization of claudin-2 in the human pancreas using mouse anti-claudin-2 antibodies based on rs12688220 genotype. **A.** Western blot of anti-claudin-2 antibody from 3 control samples genotyped at rs12688220 (TT is high risk). The antibody reacts with a protein at ~22-23 kDa, consistent with claudin-2. Samples had inflammation and/or fibrosis on histology of adjacent tissue. α -tubulin, loading control. Blots from all controls are presented in Supplementary Figure 8. **B.** Anti-claudin-2 staining (brown color) of normal-appearing control tissue localizing to ducts but not to acinar cells (scale bar=50 μ m). **C.** Severe chronic pancreatitis from a case with the high-risk (T male or TT female) genotype. Claudin-2 staining localizes to the intralobular duct (Duct), atrophic acini (*), and cells with morphologic appearance of macrophages (arrow)(scale bar=50 μ m). **D.** Chronic pancreatitis tissue from a patient with the low-risk genotype (CC or CT) with staining localizing to the duct and granular staining in acinar cells (scale bar = 100 μ m). **E.** Chronic pancreatitis, high-risk genotype with intense staining of acinar cell basolateral membrane (scale bar=100 μ m, enlarged in inset, scale bar=10 μ m). **F.** Immunofluorescence staining of control human pancreatic tissue claudin-2 staining (red) localizing to the ducts (*) and co-localizing with the macrophage marker CD68 (green, colocalized with red is yellow, arrows. Nuclei stained with Hoechst's dye, blue, scale bar = 100 μ m).

Table 1

Characterization of case subjects used for GWAS.*

	chronic pancreatitis	recurrent acute pancreatitis	chronic pancreatitis + recurrent acute pancreatitis
Stage 1	676	--	676
Alcohol-related pancreatitis †	264	--	264
Yes			
No	411	--	411
Unknown	1	--	1
Stage 2	331	579	910
Alcohol-related pancreatitis †	70	113	183
Yes			
No	256	462	718
Unknown	5	4	9
Combined	930	579	1506
Alcohol-related pancreatitis †	334	113	447
Yes			
No	667	462	1129
Unknown	6	4	10

* Does not include information from controls in Stage 1 (n=4514) or from Stage 2 (n=4053). For more complete characterization of samples, please see Supplementary Table 1 in the Appendix.

† Alcohol-related pancreatitis was assigned by the study physician at enrollment.

Table 2

Results for leading SNPs at the *PRSS1-PRSS2* and *CLDN2* loci from Stage 1, Stage 2, and joint analysis.

CHR	SNP	BP	A1*	A2	Allele Frequency (A1)		Stage 1			Stage 2			Combined		
					cases	controls	OR	se(OR)	P	OR	se(OR)	P	OR	se(OR)	P
7	rs10273639	142456928	T	C	0.350	0.424	0.712	0.044	3.0×10^{-8}	0.748	0.039	7.5×10^{-8}	0.734	0.029	2.0×10^{-14}
X	rs7057398	106144529	C	T	0.374	0.281	1.493	0.075	1.4×10^{-15}	1.210	0.066	1.8×10^{-5}	1.321	0.049	4.6×10^{-17}
X	rs12688220	106244767	T	C	0.367	0.261	1.612	0.081	2.4×10^{-21}	1.238	0.073	2.3×10^{-6}	1.385	0.054	2.3×10^{-22}

* A1 is the allele counted for purposes of computing odds ratio and associated statistics. The model used here includes covariates to control for the two leading eigenvectors for ancestry, as was done in the Pink analyses, but differs in its treatment of the minor allele count for the *CLDN2* locus, which resides on the X chromosome (as described in Online Methods). Alleles given are refSNP alleles according to dbSNP. See Supplementary Table 2 for all SNPs passing quality control and showing p-value $< 5 \times 10^{-7}$ for Stage 1 or Stage 2 or the joint analysis.

Table 3

Allele frequencies for rs10273639 (risk allele C) and rs12688220 (risk allele T) when data are stratified by controls or pancreatitis ± alcohol-related diagnosis.

Status	Alcohol-related	Number of individuals	rs10273639 ¹ (C) frequency	rs12688220 ¹ (T) frequency
Control	--	8029	0.576	0.261
Pancreatitis	No	1129	0.634	0.322
	Yes	447	0.696	0.427

¹Using data from *cases only* and in a joint analysis of both SNPs, rs12688220 predicts alcohol-related pancreatitis as genotypes ($\chi^2=29.57$; DF=2; p-value = 4×10^{-7}) or count of risk alleles ($\chi^2=13.17$; DF=1; p-value = 3×10^{-4}). rs10273639 (*PRSS1-PRSS2* locus) is a significant predictor (count of risk alleles: $\chi^2=5.68$; DF=1; p-value = 0.017; genotypes: $\chi^2=6.05$; DF=2; p-value = 0.049), even after accounting for the effects of rs12688220.