



Feasibility of the Assessment of Cholesterol Crystals in Human Macrophages Using Micro Optical Coherence Tomography

Citation

Kashiwagi, Manabu, Linbo Liu, Kengyeh K. Chu, Chen-Hsin Sun, Atsushi Tanaka, Joseph A. Gardecki, and Guillermo J. Tearney. 2014. "Feasibility of the Assessment of Cholesterol Crystals in Human Macrophages Using Micro Optical Coherence Tomography." PLoS ONE 9 (7): e102669. doi:10.1371/journal.pone.0102669. <http://dx.doi.org/10.1371/journal.pone.0102669>.

Published version

<https://doi.org/10.1371/journal.pone.0102669>

Link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:12717545>

Terms of use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material (LAA), as set forth at

<https://harvardwiki.atlassian.net/wiki/external/NGY5NDE4ZjgzNTc5NDQzMGIzZWZhMGFIOWI2M2EwYTg>

Accessibility

<https://accessibility.huit.harvard.edu/digital-accessibility-policy>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#)



Feasibility of the Assessment of Cholesterol Crystals in Human Macrophages Using Micro Optical Coherence Tomography

Manabu Kashiwagi¹, Linbo Liu², Kengyeh K. Chu¹, Chen-Hsin Sun¹, Atsushi Tanaka¹, Joseph A. Gardecki¹, Guillermo J. Tearney^{1,3,4*}

1 Harvard Medical School and Wellman Center for Photomedicine, Massachusetts General Hospital, Boston, Massachusetts, United States of America, **2** School of Electrical & Electronic Engineering and School of Chemical & Biomedical Engineering, Nanyang Technological University, Singapore, Singapore, **3** Harvard-MIT Division of Health Sciences and Technology, Cambridge, Massachusetts, United States of America, **4** Department of Pathology, Harvard Medical School and Massachusetts General Hospital, Boston, Massachusetts, United States of America

Abstract

The presence of cholesterol crystals is a hallmark of atherosclerosis, but until recently, such crystals have been considered to be passive components of necrotic plaque cores. Recent studies have demonstrated that phagocytosis of cholesterol crystals by macrophages may actively precipitate plaque progression via an inflammatory pathway, emphasizing the need for methods to study the interaction between macrophages and crystalline cholesterol. In this study, we demonstrate the feasibility of detecting cholesterol in macrophages *in situ* using Micro-Optical Coherence Tomography (μ OCT), an imaging modality we have recently developed with 1- μ m resolution. Macrophages containing cholesterol crystals frequently demonstrated highly scattering constituents in their cytoplasm on μ OCT imaging, and μ OCT was able to evaluate cholesterol crystals in cultured macrophage cells. Our results suggest that μ OCT may be useful for the detection and characterization of inflammatory activity associated with cholesterol crystals in the coronary artery.

Citation: Kashiwagi M, Liu L, Chu KK, Sun C-H, Tanaka A, et al. (2014) Feasibility of the Assessment of Cholesterol Crystals in Human Macrophages Using Micro Optical Coherence Tomography. PLoS ONE 9(7): e102669. doi:10.1371/journal.pone.0102669

Editor: Marc van Zandvoort, Maastricht University, Netherlands

Received: January 27, 2014; **Accepted:** June 23, 2014; **Published:** July 21, 2014

Copyright: © 2014 Kashiwagi et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: Research reported in this publication was supported by National Heart, Lung, and Blood Institute of the National Institutes of Health under award numbers R01HL076398 and R01HL093717. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* Email: gtearney@mgh.harvard.edu

Introduction

Cholesterol crystals are generally considered hallmarks of atherosclerosis, though their roles have long been thought to be passive elements of necrotic cores [1,2], imparting mechanical stability and stiffness to atherosclerotic lesions [3]. Recent studies have indicated that macrophage phagocytosis of cholesterol crystals may precipitate plaque progression by stimulating the nucleotide-binding domain and leucine-rich repeat containing proteins 3 (NLRP3) inflammasome pathway [4,5]. After cholesterol crystals are phagocytosed by macrophages, lysosomal destabilization and leakage of cathepsin B into the cytoplasm follow, where the enzyme indirectly activates the NLRP3 inflammasome. These findings have heightened interest in the interaction between macrophages and cholesterol crystals and have created a demand for an imaging modality capable of visualizing macrophage phagocytosis of cholesterol crystals as a tool for evaluating inflammatory activity in atherosclerosis. Cholesterol monohydrate crystals are birefringent and change the polarization of transmitted light. For this reason, polarization microscopy is considered a gold standard optical microscopy technique for visualization of cholesterol crystals in cells *in vitro*. In histologic slides of human atherosclerotic plaque, cholesterol crystals appear as oblong clefts that represent the voids left behind

following histopathologic processing. Techniques for assessment of crystalline cholesterol *in situ* however are lacking. [6,7].

Optical coherence tomography (OCT) is the highest resolution intracoronary imaging modality available currently, which provides an axial resolution of 10–20 μ m and a transverse resolution of 20–40 μ m [8–11]. *In vivo* OCT studies have been shown to elucidate the mechanism of acute coronary syndrome and atherosclerosis [12–14] and reports have suggested that OCT may be capable of identifying macrophage accumulations and large, extracellular cholesterol crystal plates [15]. Recently, we have developed new OCT technology termed Micro-OCT (μ OCT), which exhibits ten-fold improvement in resolution along every spatial direction compared to conventional OCT. μ OCT has shown an improved capability to visualize subcellular features of the human coronary artery compared with conventional OCT, including the visualization of individual macrophages and detailed morphology of extracellular cholesterol crystals [16].

In this study, we aimed to determine the feasibility of using μ OCT to evaluate and quantify macrophages and cholesterol crystals with a particular emphasis on the macrophage-cholesterol crystal interaction by imaging cultured human macrophages and cadaver human coronary arteries.




Figure 1. μ OCT instrumentation schematic. System diagram. RM: reference mirror. OL: objective lens. EC: environmental chamber. AO: analog output board. G: grating. IMAQ: image acquisition board. L: camera lens. LSC: line scan camera. SMF: single mode fiber. PC: personal computer. RAID: redundant array of independent disks. CL: Camera Link cable.
doi:10.1371/journal.pone.0102669.g001

Methods

Ethics Statement

The Institutional Review Board at the Massachusetts General Hospital approved the studies using human blood (IRB #2011P002726) and human arterial tissue (IRB #2004P000578). The written informed consents from donors were obtained for human blood draw and attached into the manuscript. The human arterial tissues have been described in the previous publication [16].

μ OCT system

OCT measures the electric field amplitude of light that is elastically scattered from within tissue in three dimensions. Depth or axial (z) ranging is achieved by interferometric measurement of the optical delay of light returned from the sample. μ OCT, as implemented here, is based on a form of OCT known as spectral-

domain OCT with several key improvements that yield high resolution in both lateral and axial directions (Figure 1) [17]. A super-continuum source (SuperK OCT Extreme, NKT Photonics, Birkerød, Denmark) provides high-bandwidth (600 nm to 1800 nm) short coherence length light, of which the spectral range from 650 nm to 950 nm is utilized by μ OCT, resulting in high axial resolution (1.3 μ m in air). Light is delivered to and collected from the interferometer optics via SM600 fiber (Thorlabs, Newton NJ). To achieve an acceptable balance between high lateral resolution (2 μ m) and sufficient depth of field (0.2 mm), μ OCT replaces the beamsplitter element typically used in OCT with a 45 degree rod mirror, which redirects the center portion of the illumination into the reference arm and introduces a circular obscuration in the center of the sample beam. The rod mirror obscures a region corresponding to a numerical aperture (NA) of approximately 0.06, in comparison to the 0.12 total beam NA. The annular geometry of the sample beam both enhances




Figure 2. Volumetric analysis of macrophage cell and cholesterol crystal by μ OCT. Cross-sectional μ OCT images of macrophage cell in coronary artery (A) and cultured macrophage cell with cholesterol crystal (B). Long and short axes (green dotted lines) of macrophage were measured at the slice with biggest cross-sectional cell area. Each 2-D area of macrophage cell and cholesterol crystals was measured. Scale bars = 50 μ m. doi:10.1371/journal.pone.0102669.g002

lateral resolution and improves depth of field [18], but does not sacrifice as much sensitivity and mitigates the presence of sidelobe artifacts compared to a fully Bessel-beam configuration. The total power incident on the sample is 10 mW. Custom software was created to control the galvanometer scanning motors while acquiring spectral data from the line camera (Sprint spL4096-140k, Basler AG, Ahrensburg, Germany).

Macrophage Cell culture

Macrophage culture was performed as described in a previous report [19]. Human peripheral blood mononuclear cells were isolated from healthy donors by density gradient centrifugation with Histopaque-1077 (Sigma-Aldrich, Saint Louis, Missouri) and washed 3 times in Dulbecco's phosphate-buffered saline (Life Technologies, Carlsbad, California). Cells were suspended in RPMI 1640 (supplemented with 10% human serum, 40 μ g/ml gentamicin, and 2 mM glutamine), and incubated for 4 hours at 37°C with 5% CO₂. Nonadherent cells were discarded, and adherent monocytes were maintained in RPMI 1640 for 7 days.

Foam cell differentiation in resting macrophages occurred after 7 days in culture with human oxidized low-density lipoprotein

(LDL). Human oxidized LDL (Intracel Resources) was added at day 8, and after 7 days, foam cells were incubated with cholesterol monohydrate crystals for 24 hours. Synthetic cholesterol (Sigma-Aldrich) was solubilized in hot acetone and crystallized by cooling [4]. After six cycles of recrystallization, the final crystallization was performed in the presence of 10% endotoxin-free water to obtain hydrated cholesterol crystals, which were confirmed by measuring angles of crystals on μ OCT (79.81 ± 2.98 (acute) and 102.13 ± 3.89 (obtuse) degrees) [20]. Crystal size was varied with a micro-tube tissue grinder (Sigma-Aldrich). At day 16, cultured cells were examined by polarization microscopy and μ OCT. μ OCT imaging was performed on the cells with illumination originating from below.

Human tissue specimens

We examined 45 human coronary arterial specimens with macrophage cells from grossly diseased arterial segments. Coronary arteries were obtained from freshly explanted human hearts provided by Capital Bioscience (Rockville, Maryland; <http://www.capitalbiosciences.com/>). Explanted hearts were harvested from organ donors after the cessation of vital signs, perfused with




Figure 3. Representative image of cultured macrophage cell with cholesterol crystal. A. Representative image of a macrophage with a cholesterol crystal. The macrophage on the right demonstrated highly scattering constituents inside its cytoplasm (yellow arrow). B, C. The Cross sectional images of the macrophage on the right. D, E. Polarization microscopy confirmed that the inclusions are cholesterol crystals (red arrow). Scale bars = 50 μ m. doi:10.1371/journal.pone.0102669.g003




Figure 4. Discrepancy image of cultured macrophage cell with cholesterol crystal. A. Discrepancy image of macrophage with cholesterol crystal. Two macrophages determined to contain cholesterol crystals by polarized light microscopy (B, C red arrow) did not show definitive evidence of crystals on the μ OCT image (yellow arrow). Scale bars = 50 μ m
doi:10.1371/journal.pone.0102669.g004

UW transplant solution and shipped on ice within a 24 hour postmortem interval. The major coronary arteries from the heart were prosected and opened longitudinally. Regions of interest were identified by gross visual inspection and optical frequency domain imaging (OFDI), a form of OCT well established for intravascular imaging [21]. Before μ OCT imaging, specimens were immersed in a thin layer of phosphate buffered saline (PBS; Sigma-Aldrich) at 25°C. μ OCT images were acquired from the luminal surface. The time between death and μ OCT imaging did not exceed 48 hours.

μ OCT image analysis

Figure 2-A shows cross-sectional μ OCT image of a macrophage in coronary artery. In the coronary artery, macrophages are characterized as highly scattering, round or ellipsoidal structures that are clearly delineated from other coronary artery components (Figure 2-A). Cross-sectional measurements of macrophage areas were manually performed by ImageJ. Long and short axes of each macrophage were measured at the slice with biggest cross-sectional cell area. The volume was determined by calculating the sum of all cross-sectional areas multiplied by the slice thickness (Simpson's rule). Cholesterol crystals within cultured macrophage cells were defined as highly scattering inclusions (Figure 2-B). Intracellular crystal volumes were measured using the same technique as macrophage volume.

Statistical analysis

Statistical analysis was performed using SPSS software for Windows version 11.0 (SPSS Inc., Chicago, Illinois). Results are expressed as medians and interquartile ranges. Qualitative data are presented as numbers and percentages satisfying the given criteria. Differences between two groups were tested by Fisher's exact test for categorical variables. Pearson correlation coefficient was used to analyze the correlation between 2 parameters. Values of $p < 0.05$ were considered significant.

Results

Imaging cultured macrophages with intracellular cholesterol crystals

We scanned 339 cultured macrophage cells and obtained matched images using μ OCT, phase contrast, and polarization

microscopy. A representative image set of macrophages containing cholesterol crystals is shown in Figure 3. On μ OCT, the cholesterol crystal-containing macrophages demonstrated highly scattering inclusions within the cytoplasm matching the location of birefringent crystals visualized under polarization microscopy. Cholesterol crystal inclusions were not always evident by μ OCT imaging however. Figure 4 illustrates such a case, in which the cholesterol crystals were detected by polarization microscopy but were not seen on the μ OCT image. Using polarization microscopy as the gold standard, the sensitivity, specificity, positive predictive value (PPV) and negative predictive value (NPV) of μ OCT for detecting cholesterol crystal within macrophages were measured to be 40.4%, 92.9%, 74.2% and 75.5%, respectively. Larger cholesterol crystals ($\geq 100 \text{ mm}^2$, assessed by polarization microscopy) were more accurately detected by μ OCT than smaller crystals ($< 100 \text{ mm}^2$) (52% vs 36%, $p < 0.05$). Among macrophage cells in which μ OCT could detect cholesterol crystals, the crystal volumes assessed by μ OCT were significantly correlated with those by polarization microscopy ($r = 0.63$, $p < 0.01$) (Figure 5).

Tissue sample imaging

We scanned 45 cadaver human coronary artery segments containing macrophage cells by μ OCT. The analysis results are summarized in Table 1. Because the μ OCT image depth depends on sample properties, we applied 2D scale for scanning area.

μ OCT clearly revealed extracellular cholesterol crystals within arterial tissue, which were characterized by intense reflections from its top and bottom surfaces (Figure 6). Transverse section of the μ OCT images showed that these crystals were cholesterol monohydrate as typified by the average crystal angles of 76.68 ± 5.83 (acute) and 105.58 ± 6.31 (obtuse) degrees of 15 randomly selected samples, which were consistent with a previous report (79.15 and 100.85 degrees) [20].

Macrophage volumes were readily quantified by μ OCT. Macrophages in human tissue specimens were visualized as highly scattering, round or ellipsoidal structures within the coronary artery that were clearly delineated from other coronary artery components (Figure 7) [16]. We found 741 total macrophages, of which 49% contained highly scattering inclusions within their cytoplasm. Figure 8 shows one example of a macrophage containing a highly scattering inclusion that is also clearly visualized after 3D rendering of the cell. This figure also shows that




Figure 5. Correlation of cholesterol crystal size between μOCT and polarization microscopy. Cholesterol crystal sizes assessed by μOCT demonstrated significant correlations with those by polarization microscopy ($r=0.63$ $p<0.01$).
doi:10.1371/journal.pone.0102669.g005

the pseudopods of the macrophage are oriented in the direction of the crystal, suggesting that this macrophage is “polarized” in the direction of the putative foreign body.

Discussion and Conclusion

OCT imaging has proven to be a unique tool for high-resolution evaluation of the coronary artery [9,12]. Crystalline cholesterol is one of the major coronary artery components that show the highest intensity signals on OCT image in human coronary artery [15,22]. In our results, macrophages that contain cholesterol crystals demonstrated highly scattering inclusions in their cytoplasm, and the feasibility of μOCT to detect cholesterol crystals in macrophage cells quantitatively with high specificity *in*

vitro was shown. Furthermore, we found some macrophages in fresh human cadaver coronary arteries demonstrated μOCT evidence of high intensity inclusions within their cytoplasm, consistent with the appearance of cholesterol crystals observed *in vitro*.

Macrophages play a key role in all phases of atherosclerosis. In particular, macrophages accumulate in vulnerable plaques prone to rupture, which causes an acute coronary event [23]. Macrophages and other plaque-related cells produce proteolytic enzymes that digest extracellular matrix and compromise the integrity of the fibrous cap. Therefore, in addition to assessment of cholesterol phagocytosis, quantitative assessment of macrophage cell distributions and sizes are also important for evaluating arterial inflammation. We have shown that μOCT is capable of

Table 1. μ OCT measured parameters from human coronary artery ex vivo.

Coronary lesion, n	45
Scanning area, μm^2	1.80 [1.04–2.98]
Macrophage number, n	741
Macrophage density, $/\mu\text{m}^2$	6.58 [1.41–16.9]
Long length, μm	35.57 [27.82–44.72]
Short length, μm	22.2 [17.26–27.16]
Volume, μm^3	7543 [4117–13679]
With high scattering constituent, n	360 (49%)
Without high scattering constituent, n	381 (51%)

Data presented are median [interquartile range] or number count (%).
doi:10.1371/journal.pone.0102669.t001




Figure 6. Representative images of cholesterol crystal in human tissue. The upper figure is a representative image of cholesterol crystal in human coronary artery, characterized by multiple intense reflections from its top and bottom surfaces (A). An en face image of a cholesterol crystal (B) is shown in panel (C). This transverse cut shows the typical angles associated with cholesterol monohydrate crystals of (83.7 (acute) and 104.2 (obtuse) degrees). Scale bars = 100 μm
doi:10.1371/journal.pone.0102669.g006




Figure 7. Macrophage cells in coronary artery. A. Macrophages (green arrow) seen by μ OCT appear as highly scattering ellipsoidal structures that can be clearly distinguished from other coronary artery cellular and subcellular components. The macrophage on the right contains a highly scattering inclusion that is consistent with a cholesterol crystal (red arrow). B. Necrotic core fibroatheroma with macrophages (yellow arrow) infiltrating the cap. C. Some macrophages attenuated the OCT signal deep to the cells. Scale bars = 100 μ m (A, B), 50 μ m (C).
doi:10.1371/journal.pone.0102669.g007

morphological quantification of macrophage cells. This detailed information that is derived from μ OCT could enable the further exploration for atherosclerosis beyond that possible with other existing imaging devices.

The present study has several limitations. We used synthetic cholesterol, which was artificially crystallized for macrophage cell cultures. These crystals might differ subtly from those crystallized in macrophages and human coronary artery *in vivo*. The detection sensitivity was low. Since light scattered from cholesterol




Figure 8. Macrophage cells with highly scattering constituents. A. Representative image of macrophage cells in human coronary artery contained highly scattering inclusions within their cytoplasm. B. Three-dimensional image of a macrophage showing a highly scattering inclusion (red-yellow color) within its cytoplasm (red arrow).
doi:10.1371/journal.pone.0102669.g008

crystals competes with scattering organelles also present in macrophages, intracellular contrast of the crystals was inconsistent. However, because cholesterol crystals are strongly birefringent, the addition of polarization sensitivity to the μ OCT system is likely to improve detection of cholesterol crystals [24]. μ OCT, as currently implemented using our bench top setup, is only appropriate for *ex vivo* or *in vitro* samples. When μ OCT is conducted *in vivo*, issues such as motion artifacts will need to be ameliorated in order to obtain clear imaging of tissue microstructure. It can sometimes be difficult to confirm whether or not macrophages in plaque contain cholesterol crystals within their cytoplasm because the cholesterol crystal is soluble in the organic agents used in histology. Therefore, we don't have the golden standard for macrophage cells with cholesterol crystals in human plaques at current time and the diagnostic accuracy of μ OCT for detecting those cells.

References

- Small DM (1998) George Lyman Duff memorial lecture. Progression and regression of atherosclerotic lesions. Insights from lipid physical biochemistry. *Arteriosclerosis* 8: 103–29.
- Schaar JA, Muller JE, Falk E, Virmani R, Fuster V, et al. (2004) Terminology for high-risk and vulnerable coronary artery plaques. Report of a meeting on the vulnerable plaque. *Eur Heart J* 25: 1077–82.
- Lorce HM, Tobias BJ, Gibson LJ, Kamm RD, Small DM, et al. (1994) Mechanical properties of model atherosclerotic lesion lipid pools. *Arterioscler Thromb.* 114: 230–4.
- Duwell P, Kono H, Rayner KJ, Sirois CM, Vladimer G, et al. (2010) NLRP3 inflammasomes are required for atherogenesis and activated by cholesterol crystals. *Nature* 464: 1357–61.
- Rajamäki K, Lappalainen J, Oörni K, Välimäki E, Matikainen S, et al. (2010) Cholesterol crystals activate the NLRP3 inflammasome in human macrophages: a novel link between cholesterol metabolism and inflammation. *PLoS One* 5: e11765.
- Klinkner AM, Waites CR, Kerns WD, Bugelski PJ (1995) Evidence of foam cell and cholesterol crystal formation in macrophages incubated with oxidized LDL by fluorescence and electron microscopy. *J Histochem Cytochem.* 43: 1071–8.
- Enomoto M, Nakagami K, Ohkuma S, Takano T (1987) Transformation of macrophages into foam cells *in vitro* induced by cholesteryl oleate liquid crystals. *J Biochem.* 101: 933–8.
- Tearney GJ, Brezinski ME, Bouma BE, Boppart SA, Pitris C, et al. (1997) *In vivo* endoscopic optical biopsy with optical coherence tomography. *Science* 276: 2037–9.
- Jang IK, Bouma BE, Kang DH, Park SJ, Park SW, et al. (2002) Visualization of coronary atherosclerotic plaques in patients using optical coherence tomography: comparison with intravascular ultrasound. *J Am Coll Cardiol* 39: 604–9.
- Yabushita H, Bouma BE, Houser SL, Aretz HT, Jang IK, et al. (2002) Characterization of human atherosclerosis by optical coherence tomography. *Circulation* 106: 1640–5.
- Tearney GJ, Yabushita H, Houser SL, Aretz HT, Jang IK, et al. (2003) Quantification of macrophage content in atherosclerotic plaques by optical coherence tomography. *Circulation* 107: 113–9.
- Jang IK, Tearney GJ, MacNeill B, Takano M, Moselewski F, et al. (2005) *In vivo* characterization of coronary atherosclerotic plaque by use of optical coherence tomography. *Circulation* 111: 1551–5.
- Tanaka A, Imanishi T, Kitabata H, Kubo T, Takarada S, et al. (2008) Morphology of exertion-triggered plaque rupture in patients with acute coronary syndrome: an optical coherence tomography study. *Circulation* 118: 2368–73.
- Raffel OC, Merchant FM, Tearney GJ, Chia S, Gauthier DD, et al. (2008) *In vivo* association between positive coronary artery remodelling and coronary plaque characteristics assessed by intravascular optical coherence tomography. *Eur Heart J* 29: 1721–8.
- Tearney GJ, Regar E, Akasaka T, Adriaenssens T, Barlis P, et al; International Working Group for Intravascular Optical Coherence Tomography (IWG-IVOC). (2012) Consensus standards for acquisition, measurement, and reporting of intravascular optical coherence tomography studies: a report from the International Working Group for Intravascular Optical Coherence Tomography Standardization and Validation. *J Am Coll Cardiol* 59: 1058–72.
- Liu L, Gardecki JA, Nadkarni SK, Toussaint JD, Yagi Y, et al. (2011) Imaging the subcellular structure of human coronary atherosclerosis using micro-optical coherence tomography. *Nat Med* 17: 1010–4.
- Liu L, Chu KK, Houser GH, Diephuis BJ, Li Y, et al. (2013) Method for quantitative study of airway functional microanatomy using micro-optical coherence tomography. *PLoS One* 8: e54473.
- Liu L, Liu C, Howe WC, Sheppard CJ, Chen N (2007) Binary-reflection polygon mirror for high-speed optical coherence microscopy. *Opt Lett.* 32: 2375–7.
- Bouhrel MA, Derudas B, Rigamonti E, Diévert R, Brozek J, et al. (2007) PPAR γ activation primes human monocytes into alternative M2 macrophages with anti-inflammatory properties. *Cell Metab* 6: 137–43.
- Loomis CR, Shipley GG, Small DM (1979) The phase behavior of hydrated cholesterol. *J Lipid Res.* 20: 525–35.
- Yun S, Tearney G, de Boer J, Itimtia N, Bouma B (2003) High-speed optical frequency-domain imaging. *Opt Express* 11: 2953–63.
- Tearney GJ, Jang IK, Bouma BE (2003) Evidence of Cholesterol Crystals in Atherosclerotic Plaque by Optical Coherence Tomographic (OCT) Imaging. *Eur. Heart J* 24: 1462.
- Virmani R, Burke AP, Farb A, Kolodgie FD (2006) Pathology of the vulnerable plaque. *J Am Coll Cardiol* 47: C13–8.
- Nadkarni SK, Pierce MC, Park BH, de Boer JF, Whittaker P, et al. (2007) Measurement of collagen and smooth muscle cell content in atherosclerotic plaques using polarization-sensitive optical coherence tomography. *J Am Coll Cardiol* 49: 1474–81.

Author Contributions

Conceived and designed the experiments: GJT MK LL. Performed the experiments: MK LL KKC CS JG. Analyzed the data: MK LL CS AT. Contributed reagents/materials/analysis tools: GJT MK LL KKC CS JG. Wrote the paper: GJT MK KK.