



Evidence that mutation accumulation does not cause aging in *Saccharomyces cerevisiae*

Citation

Kaya, Alaattin, Alexei V Lobanov, and Vadim N Gladyshev. 2014. "Evidence that mutation accumulation does not cause aging in *Saccharomyces cerevisiae*." *Aging Cell* 14 (3): 366-371. doi:10.1111/accel.12290. <http://dx.doi.org/10.1111/accel.12290>.

Published version

<https://doi.org/10.1111/accel.12290>

Link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:17295764>

Terms of use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material (LAA), as set forth at

<https://harvardwiki.atlassian.net/wiki/external/NGY5NDE4ZjgzNTc5NDQzMGIzZWZhMGFIOWI2M2EwYTg>

Accessibility

<https://accessibility.huit.harvard.edu/digital-accessibility-policy>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#)



Evidence that mutation accumulation does not cause aging in *Saccharomyces cerevisiae*

Alaattin Kaya, Alexei V. Lobanov and Vadim N. Gladyshev

Division of Genetics, Department of Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, MA 02115, USA

Summary

The concept that mutations cause aging phenotypes could not be directly tested previously due to inability to identify age-related mutations in somatic cells and determine their impact on organismal aging. Here, we subjected *Saccharomyces cerevisiae* to multiple rounds of replicative aging and assessed *de novo* mutations in daughters of mothers of different age. Mutations did increase with age, but their low numbers, < 1 per lifespan, excluded their causal role in aging. Structural genome changes also had no role. A mutant lacking thiol peroxidases had the mutation rate well above that of wild-type cells, but this did not correspond to the aging pattern, as old wild-type cells with few or no mutations were dying, whereas young mutant cells with many more mutations continued dividing. In addition, wild-type cells lost mitochondrial DNA during aging, whereas shorter-lived mutant cells preserved it, excluding a causal role of mitochondrial mutations in aging. Thus, DNA mutations do not cause aging in yeast. These findings may apply to other damage types, suggesting a causal role of cumulative damage, as opposed to individual damage types, in organismal aging.

Key words: aging; DNA damage; lifespan; mitochondria; mutations; thiol peroxidase; yeast.

Introduction

The nature of aging remains one of the grand mysteries of biology (Kirkwood & Austad, 2000). In the 1950s, Failla and Szilard proposed that random somatic mutations cause cell and tissue dysfunction, leading to aging (Failla, 1958; Szilard, 1959). Maynard Smith questioned this hypothesis (Maynard, 1959), initiating the debate that is not resolved to this day. Previous studies revealed accumulation of age-related mutations in human cells (Oller *et al.*, 1989; Jones *et al.*, 1995), mice (Dolle *et al.*, 2000), and flies (Garcia *et al.*, 2010). However, the molecular mechanisms involved remain unknown. One possibility is that decreased fidelity (e.g., due to protein damage) of enzymes responsible for preventing or repairing DNA damage leads to mutation accumulation and aging (Andressoo *et al.*, 2006; Brosh & Bohr, 2007; Gorbunova *et al.*, 2007), but this could also be due to critical mutations in these genes accumulated through the lifespan. Deficiency in these enzymes

can lead to premature aging, but whether they limit lifespan under conditions of their sufficiency is unclear. Another possibility is the accumulation of oxidative damage, which decreases or abolishes the activities of repair enzymes. Oxidative stress is considered as one of the main factors in the development and progression of cancer and other age-related diseases (Ames *et al.*, 1993).

While previous data are consistent with increased mutation load as a function of age, it remains unclear whether mutations in the DNA, the only theoretically nonrenewable molecular species within cells (the species that can be modified irreversibly), cause aging. This is because the concept that mutations cause functional decline and other aging phenotypes could not be directly tested due to inability to fully assess age-related mutations in individual somatic cells and determine their impact on organismal aging. Recently, single-cell sequencing technology has been applied to individual normal human clonal crypt, showing that chromosomal rearrangements occur in the last decades of normal human lifespan (Hsieh *et al.*, 2013). However, it has been hard to determine if these mutations contribute to the aging process.

In this regard, the unicellular yeast, *Saccharomyces cerevisiae*, could be very useful in addressing the half-century-old debate. Yeast cells produce a certain number of daughter cells before they die, and the total number of daughter cells produced by the mother defines her replicative lifespan (Kaeberlein *et al.*, 2007). Significant conservation of proteins in the DNA repair pathway between yeast and higher eukaryotes further elevates the yeast as a model to study age-related genomic instability (Bitterman *et al.*, 2003). Most importantly, the yeast genome is well characterized, and individual cells can give rise to colonies that can be sequenced, thereby establishing the genotype of the daughter. Moreover, the yeast genome is small, so the impact of its mutations can be quantified, which is more difficult to do in the case of model organisms with large genomes. Here, we subjected *S. cerevisiae* to multiple rounds of replicative aging and assessed *de novo* mutations in daughters of mothers of different age. The data show that DNA mutations do not cause aging in yeast.

Results and discussion

To address the role of DNA mutations in aging, we subjected *Saccharomyces cerevisiae* to replicative lifespan analyses and sequenced the genomes of clones derived from individual daughter cells of mothers of different age. First, we collected 5th (designated young) and last alive (~30th, designated old) daughter cells from each of the 4 wild-type (WT) mother cells and generated colonies from them (Figs 1 and 2A). We then used virgin cells from these colonies as mothers and again collected 5th (from the young lineage) and last (from the old lineage) daughters and made colonies from them. This procedure was repeated six times by selecting young cells from young lineages and old cells from old lineages (Figs 1 and S1), followed by high-coverage sequencing of the genomes (>99.99% completeness) of single cell-derived colonies after 2, 4, and 6 cycles of the procedure (a total of 24 genomes).

Analyses of these nuclear genomes revealed a steady increase in *de novo* mutations (base substitutions and indels) (Table 1) with each cycle of the procedure (Fig. 2B). After six cycles, young cells accumulated ~0.5 mutations/cell and mutations increased threefold in old cells ($P < 0.03$),

Correspondence

Vadim N. Gladyshev, Genetics Division, Department of Medicine, Brigham and Women's Hospital and Harvard Medical School, 77 Ave Louis Pasteur, Boston, MA 02115, USA. Tel.: 617-525-5140; fax: 617-525-5147; e-mail: vgladyshev@rics.bwh.harvard.edu

Accepted for publication 18 October 2014




Fig. 1 Experimental design of the pedigree assay. Virgin mothers (shown as initial mother in the figure) were subjected individually to replicative lifespan, and their 5th and last daughters isolated to give rise to individual clones. Virgin mothers from these clones were again subjected to the replicative lifespan procedure, isolating either 5th (from the 'young' lineage) or last (from the 'old' lineage) daughters. This procedure was repeated six times for four independent WT and four independent $\Delta 8$ lines, followed by genome sequencing of the respective colonies following the 2nd, 4th, and 6th cycles (shown by red squares). In total, 48 genomes of individual cell-derived lineages were sequenced and analyzed.

consistent with the previous data in flies and mice, representing both dividing (mouse) and postmitotic (fruit fly) cells (Dolle *et al.*, 2000; Garcia *et al.*, 2010). However, the low number of mutations observed in old yeast cells, ~ 1.5 mutations per six cycles or ~ 0.4 mutations per lifespan of an average cell, was inconsistent with the causal role of these mutations in aging. This finding also agrees with observations of yeast lifespan. For example, although daughters of very old mother cells have a shorter lifespan, the granddaughters restored lifespan to that observed in the daughters of young mothers (Kennedy *et al.*, 1994). We also found that the lifespan of progeny of young and old cells at the end of six cycles was similar and matched that of original cells (Fig. S2). In addition, a recent report demonstrated that spores generated by aging diploid cells had the same replicative potential as those from young cells (Unal *et al.*, 2011). Thus, while nuclear mutations do accumulate with age, they do not cause aging.

Our further analysis of the observed mutations revealed that many of them occurred on chromosome XII, where the ribosomal DNA (rDNA) repeats are located. rDNA has been identified as a factor leading to ERCs, which accumulate during replicative aging (Sinclair & Guarente, 1997). It was also shown that rDNA instability is a contributing factor to the aging process (Kobayashi, 2008) and that aging is accompanied by a significant increase in loss of heterozygosity (Lindstrom *et al.*, 2011). In this regard, our observation of a higher mutation rate in this region is consistent with previous findings. An increased number of rDNA reads was recently shown in aged mother cells (Hu *et al.*, 2014). We also analyzed the reads aligned to the rDNA locus in both young and old cells and found no large deletions/insertions or an increase in the number of rDNA reads during replicative aging (Fig. S3).

To assess the contribution of oxidative stress to mutation accumulation and compare it with age-dependent mutation load, we similarly analyzed a mutant strain (designated $\Delta 8$) lacking eight thiol peroxidases (Fomenko *et al.*, 2011; Kaya *et al.*, 2014), which are the enzymes (peroxiredoxins and glutathione peroxidases) that detoxify hydroperoxides. We further sequenced the 24 genomes corresponding to young and old $\Delta 8$ cells after 2, 4, and 6 cycles, similarly to the procedure described above for WT cells (Table 2; Figs S1 and S4). This mutant showed a 12-fold higher mutation rate than WT cells, ~ 6 mutations per




Fig. 2 Mutation accumulation does not cause aging in yeast. (A) Overall experimental design of the study. Individual yeast cells (mothers) were subjected to replicative aging, and their daughters were collected at different ages of mother cells. These cells gave rise to colonies, from which new mother cells were taken and subjected to a second round of the aging procedure. The genomes of young and old lines after 2, 4, and 6 cycles were sequenced. See Fig. 1 for details. (B) Average number of mutations per aging cycle of young and old WT cells. (C) Average number of mutations per aging cycle of young and old $\Delta 8$ cells. Error bars represent the standard error of mean. (D) Mean mutation rate (number of mutations per line divided by the number of cell divisions and the length of the genome). The graph shows the mean of four lines in each experiment, and error bars are standard errors. *P*-values for each comparison were calculated using the Wilcoxon rank-sum test.

six cycles (Fig. 2C). However, even young $\Delta 8$ cells had twice as many mutations as the old WT cells, and this situation did not correspond to the aging patterns of these cells, that is, old WT cells had few mutations, yet stopped dividing and died, whereas young $\Delta 8$ cells had many more mutations, yet they continued dividing. The overall substitution mutation rate of WT cells was in line with the known mutation rate in yeast (Lynch

Table 1 Mutations observed in young and old WT clones following pedigree analysis

	WT							
	Y1	Y2	Y3	Y4	O1	O2	O3	O4
Substitutions Position								
Noncoding	0	0	1	0	1	0	0	1
Nonsense	0	0	0	0	0	0	0	0
Missense	0	0	0	0	0	0	1	0
Silent	0	0	0	0	0	0	0	0
Indels Position								
Noncoding	1	0	0	0	0	1	0	2
Coding	0	0	0	0	0	0	0	0
All Mutations	1	0	1	0	1	1	1	3
Mean ($\times 10^{-10}$)	2.65×10^{-10}				4.1×10^{-10}			

Mutation distributions for each pedigree lines are shown after completion of six cycles of aging. Mean mutation rates are calculated based on the total number of substitutions and small indels divided by the total number of generations for four lines. The whole list of mutations together with positions and nucleotide changes is shown in Table S1. Y is young, and O is old.

Table 2 Mutations observed in young and old $\Delta 8$ clones following pedigree analysis

	$\Delta 8$							
	Y1	Y2	Y3	Y4	O1	O2	O3	O4
Substitutions Position								
Noncoding	0	0	0	1	4	3	1	0
Nonsense	0	0	0	0	0	0	0	0
Missense	1	1	1	0	2	0	1	0
Silent	1	3	1	0	1	1	1	1
Indels Position								
Noncoding	1	1	2	1	0	1	1	1
Coding	0	0	0	0	1	0	3	1
All Mutations	3	5	4	2	8	5	7	3
Mean ($\times 10^{-10}$)	41.5×10^{-10}				47.1×10^{-10}			

Mutation distributions for each pedigree lines are shown after completion of six cycles of aging. Mean mutation rates are calculated based on the total number of substitutions and small indels divided by the total number of generations for four lines. The whole list of mutations together with positions and nucleotide changes is shown in Table S2. Y is young, and O is old.

et al., 2008) (Fig. 2D). Although larger sample sizes would have allowed measuring the mutation rates more precisely, our goal was to determine the actual number of mutations in individual yeast cells per lifespan. It was clear from our analysis that both young and old WT cells had a much lower mutation rate than either young or old $\Delta 8$ cells (Fig. 2D). We also examined mutations in repetitive regions, which are thought to exhibit higher mutation rates (Lynch *et al.*, 2008) and impact protein quality control (Nyström & Liu, 2013), but detected no such mutations during aging. Together, these findings, and the low number of mutations in old cells (< 1 per lifespan), exclude a causal role of mutation accumulation in aging in yeast.

Do mutations in the yeast mitochondrial genome cause aging? As mitochondria are the major source of reactive oxygen species, mtDNA may be vulnerable to oxidative damage (Harman, 1956). The mitochon-

drial genome has a higher mutation rate than the nuclear genome (Lynch *et al.*, 2008), and the biology of yeast mitochondria has been implicated in aging (Breitenbach *et al.*, 2012). We found that mtDNA was consistently lost in WT cells after they generated 10–20 daughter cells, and this effect was also seen in young cells after several aging cycles, making many of these cells respiration deficient (Fig. 3; Figs S5 and S7). The observation that cells lose mtDNA during the replicative aging process leading to nonrespiring cells was also reported previously (Veatch *et al.*, 2009; Lindstrom *et al.*, 2011). An interesting observation was that some of these cells showed mitochondrial mutations just prior to completely losing the mitochondrial genome, explaining the loss of the mitochondrial genome during the aging progress. Although yeast cells grown on glucose do not rely on respiration and do not utilize mtDNA, mutations may disrupt integrity of mtDNA, so it may be lost. Interestingly, among the cells that we analyzed, the two old WT lines that still had functional mitochondria possessed 10 and 6 distinct noncoding substitution mutations.

In contrast, $\Delta 8$ cells had a higher mtDNA content in both old and young cells, did not lose mtDNA during aging and progression through aging cycles, and did not accumulate mutations in the mitochondrial genome (Fig. 3; Figs S6 and S7). Although in some samples we observed heterogeneous mutations (i.e., both original and mutant sequences were present) none of them persisted, that is, mutated mtDNA was lost in subsequent cycles, whereas the original mtDNA remained. Throughout the procedure, $\Delta 8$ cells were fully dependent on respiration, yet their lifespan was shorter than that of WT cells. Thus, not only mtDNA




Fig. 3 Mitochondrial content is higher in $\Delta 8$ cells and does not decrease during aging. Genomes of individual clones (four young and four old at each cycle) were sequenced following 2, 4, and 6 cycles of aging. X_{mt} coverage (mitochondrial genome) divided by X_{nuc} coverage (nuclear genome) is shown. (A) Mitochondrial genome abundance of young WT and $\Delta 8$ cells following 2 and 6 cycles. (B) Mitochondrial genome abundance of old WT and $\Delta 8$ cells following 2 and 6 cycles.

mutations, but even the presence of the functional mitochondrial genome, did not correlate with the aging process. A possibility remains that transient heterogeneous mitochondrial mutations in $\Delta 8$ cells contribute to the aging process at the level of individual mitochondria, but deleterious mutations are selected against during colony expansion. It should also be noted that although an increase in oxidative stress levels leads to increased mutation load, oxidative stress also targets numerous other molecules. It is unlikely that these conditions affect only DNA in $\Delta 8$ cells. In future studies, it would be useful to use a strain, which has a mutated proofreading domain of a nuclear DNA polymerase, thus leading specifically to increased mutation load without affecting other cellular systems.

It would also be interesting to understand why $\Delta 8$ cells rely on respiration. It is known that age asymmetry between the mother and daughter cells is affected by the end of life of mother cells (McMurray & Gottschling, 2004). Old cells pass some of the damaged molecules including aggregated proteins and dysfunctional mitochondria to their daughter cells (Henderson & Gottschling, 2008; Steinkraus *et al.*, 2008; Kaerberlein, 2010), and daughters of old mothers live a shorter life (Kennedy *et al.*, 1994). However, granddaughters of old mother cells are apparently able to clear the damage (McMurray & Gottschling, 2004; Kaerberlein, 2010). Previously, age-dependent deterioration in mitochondrial genome integrity was shown in mice; however, mitochondrial mutations did not limit the lifespan of these animals (Vermulst *et al.*, 2007). In yeast, it was also shown that the absence of the mitochondrial genome does not affect the replicative lifespan in both BY4742 and YDW2 backgrounds (Kaerberlein *et al.*, 2005; Woo & Poyton, 2009), and the replicative lifespan is increased in JM43 and D273-10B backgrounds (Woo & Poyton, 2009).

Aging is often viewed in terms of gradual degeneration and accumulation of damage, leading to decline in fitness. We agree with this notion of aging as a gradual process driven by damage (in the broad sense of it, that is, damage not limited to by-products and errors) (Gladyshev, 2013). Universality of the aging process and the conservation of aging pathways and genetic and dietary interventions that affect lifespan point to a common basis for aging in diverse organisms. Point mutations represent an important form of cellular damage, but not the only one. For example, genome rearrangements have been shown to increase during replicative (McMurray & Gottschling, 2004) and chronological (Longo & Fabrizio, 2012) aging in yeast. However, we found no structural genome rearrangements during aging in the cells that we examined. A possibility remains that point mutations and structural rearrangements are eliminated during cell growth, that is, they accumulate in large numbers during aging, but selection acts during colony expansion of daughters of old cells eliminating them. However, we think it is highly unlikely that all mutations and rearrangements can be eliminated, returning the genome to the original state, especially considering that most age-related mutations are expected to be neutral.

A recent study analyzed populations of yeast old mother cells, in which more than ~85% of cells ceased to divide, and observed many genomic rearrangements accompanying aging, including mitochondrial DNA transfer to the nuclear genome, translocations, and retrotranspositions (Hu *et al.*, 2014). This study cannot be directly related to our conclusion that DNA mutations do not cause aging (but together the two studies hold significant information about progression of replicative aging and the processes after terminally old mother cells stop dividing). This is because that other study analyzed enriched, purified aged mother cells at the population level, whereas we did this at the individual cell level, which was required to assess the causal role of mutations. It is

possible that a significant fraction of the genomic rearrangements observed (Hu *et al.*, 2014) occurred after the last daughter was formed, or these mutations occurred in a fraction of dying cells in the population. Thus, we cannot exclude a possibility that mutations contribute to demise of the mother cell (which is synonymous with her inability to bud off one more daughter). But, this would be equivalent to the contribution of mutations to death of an organism, as opposed to having a causal role in the aging process.

Another recent study analyzed Ty1 retrotransposon expression and mobility during chronological aging in *S. paradoxus* (VanHoute & Maxwell, 2014). Increased expression and mobility of retrotransposons with age have been found in nearly every aging model and thought to promote genome instability during aging (Moskalev *et al.*, 2013). However, this study showed that, while the strain expressing Ty1 element had 40-fold increased mobility to the rDNA region, neither Ty1 mobility nor increased mutation rate of a representative Ty1 gene led to chronological lifespan decrease. Furthermore, increased expression of a Ty1 could extend chronological lifespan under certain conditions (VanHoute & Maxwell, 2014).

Replicative aging in yeast can be defined as the progression of mitotic division until an ultimate phenotype that prevents further daughter cell formation. During progression of aging, accumulation of different types of damage has been reported. If a single damage form limits lifespan, selection on other damage forms should be relaxed, leading to a certain degree of synchronization with regard to their deleterious impact. Thus, we speculate that application of our findings to other individual damage types and to other organisms suggests that there is no individual damage form that represents, when considered in isolation, a main causal factor in aging. If a damage form decreases fitness, organism can develop protective strategies to downgrade it to a milder form or modify its metabolism to decrease production of this damage. But the mild damage forms are too numerous to be protected against (Gladyshev, 2013). Thus, even if mutations and other individual damage forms, when taken in isolation, do not cause aging, aging may still result from cumulative damage, to which these damage forms contribute. From this perspective, mutations in nuclear and mitochondrial genomes, together with the myriad of other damage forms, contribute to normal aging, but can cause aging only at the level of cumulative damage.

Experimental procedures

Yeast strains

All yeast strains were in BY4741 background. A mutant strain, lacking all eight thiol peroxidases, was reported previously (Fomenko *et al.*, 2011). Cells were grown on yeast extract–peptone–dextrose (YPD) medium.

Pedigree analyses and mutation detection

WT and $\Delta 8$ cells were grown at 30°C for 2 days. Several virgin mothers were placed in predetermined positions with a micromanipulator. We isolated 5th and last daughters from these mothers and allowed them to form colonies. For the 2nd round, single virgin cells from these colonies were subjected to replicative lifespan analysis, and 5th (from each young line) and last (from each old line) daughters were collected again, to obtain additional colonies. This procedure was repeated six times. At the end of the 6th cycle, DNA was isolated from each colony and paired-end sequenced on an Illumina platform (400 bp insert size, 75 bp nucleotide reads, paired end). The depth of coverage for each line was between 50X and 150X. Burrows–Wheeler Aligner (BWA) was used for alignment

(Li & Durbin, 2009), revealing more than 99.99% genome completeness (regions covered by more than three reads). Pairwise comparisons of sequenced genomes were performed using the Genome Analysis Toolkit (GATK) (McKenna *et al.*, 2010) to identify mutations covered by at least three reads and with the GATK quality score higher than 10 ($P = 0.001$ for misidentification). The data were further analyzed using the SHORE toolkit (Schneeberger *et al.*, 2009), the nucleotide variants were cross-checked with the GATK output, and mutations were manually verified with Tablet (Milne *et al.*, 2010). We also adjusted parameters to include cases in which the fraction of reads supporting an alternative allele was lowered to 0.5; however, this did not result in an increase in the number of candidate mutations. In addition, variable tandem repeats were extracted from Vences *et al.*, 2009 (intergenic repeats) and Verstrepen *et al.*, 2005 (intragenic repeats) and further subjected to our analysis.

Calculation of mutation rates

Mutation rate was calculated as the number of mutations per line divided by the number of cell divisions and the length of the genome. The figures show the mean of four lines in each experiment, and error bars are standard errors. P -values for each comparison were calculated using the Wilcoxon rank-sum test.

Analyses of mitochondrial mutations and the rDNA locus

For normalization, X_{mt} coverage (mitochondrial genome) was divided by X_{nuc} coverage (nuclear genome). The same procedure was also applied to analyze the rDNA locus (X_{rDNA}). The total number of reads mapped to the rDNA locus was divided by X_{nuc} coverage and plotted. YPEG plates containing ethanol and glycerol were used to assess growth on respiratory substrates. Petite colony formation was performed as previously described (Ferguson & Von Borstel, 1992). Antimycin A sensitivity was tested on YPD plates containing 2 $\mu\text{g}/\text{mL}$ of antimycin A.

Acknowledgments

We thank Amnon Koren for the help with mutation rate analyses, and Daniel Promislow and Matt Kaeberlein for suggestions.

Author contributions

Conceived and designed the experiments: AK, VNG. Performed the experiments: AK. Analyzed the data: AK, AVL, VNG. Wrote the paper: AK, VNG.

Funding

This research was funded by NIH AG038004 and GM065204.

Conflict of interest

The authors declare no conflict of interest.

References

Ames BN, Shigenaga MK, Hagen TM (1993) Oxidants, antioxidants, and the degenerative diseases of aging. *Proc. Natl Acad. Sci. USA* **90**, 7915–7922.
Andressoo JO, Hoeijmakers JH, Mitchell JR (2006) Nucleotide excision repair disorders and the balance between cancer and aging. *Cell Cycle* **5**, 2886–2888.

Bitterman KJ, Medvedik O, Sinclair DA (2003) Longevity regulation in *Saccharomyces cerevisiae*: linking metabolism, genome stability, and heterochromatin. *Microbiol. Mol. Biol. Rev.* **67**, 376–399.
Breitenbach M, *et al.* (2012) The role of mitochondria in the aging processes of yeast. *Subcell. Biochem.* **57**, 55–78.
Brosh RM Jr, Bohr VA (2007) Human premature aging, DNA repair and RecQ helicases. *Nucleic Acids Res.* **35**, 7527–7544.
Breitenbach M, Laun P, Dickinson JR, Klocker A, Rinnerthaler M, Dawes IW, Aung-Htut MT, Breitenbach-Koller L, Caballero A, Nyström T, Büttner S, Eisenberg T, Madeo F, Ralser M (2012) The role of mitochondria in the aging processes of yeast. *Subcell. Biochem.* **57**, 55–78.
Dolle ME, Snyder WK, Gossen JA, Lohman PH, Vijg J (2000) Distinct spectra of somatic mutations accumulated with age in mouse heart and small intestine. *Proc. Natl Acad. Sci. USA* **97**, 8403–8408.
Failla G (1958) The aging process and cancerogenesis. *Ann. N. Y. Acad. Sci.* **71**, 1124.
Ferguson LR, Von Borstel RC (1992) Induction of the cytoplasmic ‘petite’ mutation by chemical and physical agents in *Saccharomyces cerevisiae*. *Mutat. Res.* **265**, 103–148.
Fomenko DE, *et al.* (2011) Thiol peroxidases mediate specific genome-wide regulation of gene expression in response to hydrogen peroxide. *Proc. Natl Acad. Sci. USA* **108**, 2729–2734.
Garcia AM, *et al.* (2010) Age and temperature-dependent somatic mutation accumulation in *Drosophila melanogaster*. *PLoS Genet.* **6**, e1000950.
Gladyshev VN (2013) The origin of aging: imperfectness-driven non-random damage defines the aging process and control of lifespan. *Trends Genet.* **29**, 506–512.
Gorbunova V, Seluanov A, Mao Z, Hine C (2007) Changes in DNA repair during aging. *Nucleic Acids Res.* **35**, 7466–7474.
Harman D (1956) Aging: a theory based on free radical and radiation chemistry. *J. Gerontol.* **11**, 298–300.
Henderson KA, Gottschling DE (2008) A mother’s sacrifice: what is she keeping for herself? *Curr. Opin. Cell Biol.* **20**, 723–728.
Hsieh JC, Van Den Berg D, Kang H, Hsieh CL, Lieber MR (2013) Large chromosome deletions, duplications, and gene conversion events accumulate with age in normal human colon crypts. *Aging Cell* **12**, 269–279.
Hu Z, Chen K, Xia Z, Chavez M, Pal S, Seol JH, Chen CC, Li W, Tyler JK (2014) Nucleosome loss leads to global transcriptional up-regulation and genomic instability during yeast aging. *Genes Dev.* **28**, 396–408.
Jones IM, *et al.* (1995) Impact of age and environment on somatic mutation at the hprt gene of T lymphocytes in humans. *Mutat. Res.* **338**, 129–139.
Kaeberlein M (2010) Lessons on longevity from budding yeast. *Nature* **464**, 513–519.
Kaeberlein M, Hu D, Kerr EO, Tsuchiya M, Westman EA, Dang N, Fields S, Kennedy BK (2005) Increased life span due to calorie restriction in respiratory-deficient yeast. *PLoS Genet.* **1**, e69.
Kaeberlein M, Burtner CR, Kennedy BK (2007) Recent developments in yeast aging. *PLoS Genet.* **3**, e84.
Kaya A, Lobanov AV, Gerashchenko MV, Koren A, Fomenko DE, Koc A, Gladyshev VN (2014) Thiol peroxidase deficiency leads to increased mutation load and decreased fitness in *Saccharomyces cerevisiae*. *Genetics*. doi:10.1534/genetics.114.169243. Epub ahead of print.
Kennedy BK, Austriaco NR Jr, Guarente L (1994) Daughter cells of *Saccharomyces cerevisiae* from old mothers display a reduced life span. *J. Cell Biol.* **127**, 1985–1993.
Kirkwood TB, Austad SN (2000) Why do we age? *Nature* **408**, 233–238.
Kobayashi T (2008) A new role of the rDNA and nucleolus in the nucleus–rDNA instability maintains genome integrity. *BioEssays* **30**, 267–272.
Li H, Durbin R (2009) Fast and accurate short read alignment with Burrows–Wheeler transform. *Bioinformatics* **25**, 1754–1760.
Lindstrom DL, Leverich CK, Henderson KA, Gottschling DE (2011) Replicative age induces mitotic recombination in the ribosomal RNA gene cluster of *Saccharomyces cerevisiae*. *PLoS Genet.* **7**, e1002015.
Longo VD, Fabrizio P (2012) Chronological aging in *Saccharomyces cerevisiae*. *Subcell. Biochem.* **57**, 101–121.
Lynch M, *et al.* (2008) A genome-wide view of the spectrum of spontaneous mutations in yeast. *Proc. Natl Acad. Sci. USA* **105**, 9272–9277.
Maynard SJ (1959) A theory of aging. *Nature* **184**, 956–957.
McKenna A, *et al.* (2010) The genome analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Res.* **20**, 1297–1303.
McMurray MA, Gottschling DE (2004) An age-induced switch to a hyper-recombinational state. *Science* **301**, 1908–1911.

- Milne I, et al. (2010) Tablet-next generation sequence assembly visualization. *Bioinformatics* **26**, 401–402.
- Moskalev AA, Shaposhnikov MV, Plyusnina EN, Zhavoronkov A, Budovsky A, Hagai Y, Faraifeld VE (2013) The role of DNA damage and repair in aging through the prism of Koch-like criteria. *Ageing Res. Rev.* **12**, 661–684.
- Nyström T, Liu B (2013) Protein quality control in time and space - links to cellular aging. *FEMS Yeast Res.* doi:10.1111/1567-1364. Epub ahead of print.
- Oller AR, Rastogi P, Morgenthaler S, Thilly WG (1989) A statistical model to estimate variance in long term-low dose mutation assays: testing of the model in a human lymphoblastoid mutation assay. *Mutat. Res.* **216**, 149–161.
- Schneeberger K, et al. (2009) Simultaneous alignment of short reads against multiple genomes. *Genome Biol.* **10**, R98.
- Sinclair DA, Guarente L (1997) Extrachromosomal rDNA circles—a cause of aging in yeast. *Cell* **91**, 1033–1042.
- Steinkraus KA, Kaeberlein M, Kennedy BK (2008) Replicative aging in yeast: the means to the end. *Annu. Rev. Cell Dev. Biol.* **24**, 29–54.
- Szilard L (1959) On the nature of the aging process. *Proc. Natl Acad. Sci. USA* **45**, 30–45.
- Unal E, Kinde B, Amon A (2011) Gametogenesis eliminates age-induced cellular damage and resets life span in yeast. *Science* **332**, 1554–1557.
- VanHoute B, Maxwell PH (2014) Extension of *Saccharomyces paradoxus* chronological lifespan by retrotransposons in certain media conditions is associated with changes in reactive oxygen species. *Genetics*. doi:10.1534/genetics.114.168799. Epub ahead of print.
- Veatch JR, McMurray MA, Nelson ZW, Gottschling DE (2009) Mitochondrial dysfunction leads to nuclear genome instability via an iron-sulfur cluster defect. *Cell* **137**, 1247–1258.
- Vermulst M, et al. (2007) Mitochondrial point mutations do not limit the natural lifespan of mice. *Nat. Genet.* **39**, 540–543.
- Verstrepen KJ, Jansen A, Lewitter F, Fink GR (2005) Intragenic tandem repeats generate functional variability. *Nat. Genet.* **37**, 986–990.
- Vinces MD, Legendre M, Caldara M, Hagihara M, Verstrepen KJ (2009) Unstable tandem repeats in promoters confer transcriptional evolvability. *Science* **324**, 1213–1216.
- Woo DK, Poyton RO (2009) The absence of a mitochondrial genome in rho0 yeast cells extends lifespan independently of retrograde regulation. *Exp. Gerontol.* **44**, 390–397.

Supporting Information

Additional Supporting Information may be found in the online version of this article at the publisher's web-site.

Fig. S1 Age at which daughter cells of young and old WT mothers were taken for further analysis.

Fig. S2 Lifespan of clones derived from young and old mother cells is not affected by the aging process.

Fig. S3 Comparison of the reads mapped to the rDNA region of WT young and old lines.

Fig. S4 Age at which daughter cells of young and old $\Delta 8$ mothers were taken for further analysis.

Fig. S5 Depletion of mitochondrial DNA and respiratory function in WT lines during aging.

Fig. S6 No depletion of mitochondrial DNA in $\Delta 8$ lines during aging.

Fig. S7 Induction of petite colony formation.

Table S1 Mutations identified in WT cells during aging.

Table S2 Mutations identified in $\Delta 8$ cells during aging.