



Scaling dimensions of monopole operators in the CPN_b!1 theory in 2 + 1 dimensions

Citation

Dyer, Ethan, Márk Mezei, Silviu S. Pufu, and Subir Sachdev. 2015. "Scaling dimensions of monopole operators in the CPN_b!1 theory in 2 + 1 dimensions." J. High Energ. Phys. 2015 (6) (June). doi:10.1007/jhep06(2015)037.

Published version

[https://doi.org/10.1007/JHEP06\(2015\)037](https://doi.org/10.1007/JHEP06(2015)037)

Link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:34331446>

Terms of use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Open Access Policy Articles (OAP), as set forth at

<https://harvardwiki.atlassian.net/wiki/external/NGY5NDE4ZjgzNTc5NDQzMGIzZWZhMGFIOWI2M2EwYTg>

Accessibility

<https://accessibility.huit.harvard.edu/digital-accessibility-policy>

Share Your Story

The Harvard community has made this article openly available. Please share how this access benefits you. [Submit a story](#)

Scaling dimensions of monopole operators in the \mathbb{CP}^{N_b-1} theory in $2 + 1$ dimensions

Ethan Dyer,¹ Márk Mezei,² Silviu S. Pufu,³ and Subir Sachdev^{4,5}

¹Stanford Institute for Theoretical Physics, Stanford University, Stanford, CA 94305, USA

²Princeton Center for Theoretical Science, Princeton University, Princeton, NJ 08544, USA

³Joseph Henry Laboratories, Princeton University, Princeton, NJ 08544, USA

⁴Department of Physics, Harvard University, Cambridge, MA 02138, USA

⁵Perimeter Institute for Theoretical Physics, Waterloo, Ontario N2L 2Y5, Canada

Abstract

We study monopole operators at the conformal critical point of the \mathbb{CP}^{N_b-1} theory in $2 + 1$ spacetime dimensions. Using the state-operator correspondence and a saddle point approximation, we compute the scaling dimensions of the operators that insert one or two units of magnetic flux to next-to-leading order in $1/N_b$. We compare our results to numerical studies of quantum antiferromagnets on two-dimensional lattices with $SU(N_b)$ global symmetry, using the mapping of the monopole operators to valence bond solid order parameters of the lattice antiferromagnet. For the monopole operators that insert three or more units of magnetic flux, we find that the rotationally-symmetric saddle point is unstable; in order to obtain the scaling dimensions of these operators, even at leading order in $1/N_b$, one should consider non-spherically-symmetric saddles.

Contents

1	Introduction	2
2	Setup	6
3	$N_b = \infty$ theory	8
4	$1/N_b$ corrections	10
4.1	Effective action at quadratic level	10
4.2	Eigenvalues of the integration kernels	11
4.3	Kernels at $q = 0$	16
4.3.1	The D kernel at $q = 0$	16
4.3.2	The K kernels at $q = 0$	17
4.4	Kernels for general q	18
4.4.1	The D kernel	19
4.4.2	The K and F kernels	20
4.5	Numerical Results	24
5	Stability	25
5.1	Numerical results	25
5.2	Analytic understanding in the large q limit	27
6	Conclusions	30
A	Derivation of integration kernels	31
A.1	Evaluation of Fourier transforms	31
A.2	Check of gauge invariance	37
B	Formulas for \mathcal{I}_B and \mathcal{I}_F	37
B.1	Results for $q = 0, 1/2$	37
B.2	Results for arbitrary q	38
C	Asymptotic expansions	44
C.1	Small distance expansion of the Green's function	44
C.2	UV asymptotic of the scalar kernel	47
C.3	The mixed kernel	49
C.4	Gauge field kernel	50

C.5 Ultraviolet expansion in the $\mathbb{C}\mathbb{P}^{N-1}$ model	54
C.6 Fourier transforms on $S^2 \times \mathbb{R}$	55

1 Introduction

In 2 + 1 dimensions, pure U(1) gauge theory confines [1]. One can prevent confinement by introducing a sufficiently large number N of massless matter fields, in which case the infrared dynamics is believed to be governed by a non-trivial interacting conformal field theory (CFT). Such CFTs arise quite frequently in the description of quantum critical points of condensed matter systems in two spatial dimensions [2–15]. They also serve as useful toy-models for more intricate four-dimensional dynamics, as they can be studied perturbatively in the $1/N$ expansion, where the gauge interactions are suppressed [16–19].

Our goal in this paper is to study monopole operators in one such CFT, namely the $\mathbb{C}\mathbb{P}^{N_b-1}$ theory tuned to criticality. This theory is a nonlinear sigma-model with $\mathbb{C}\mathbb{P}^{N_b-1}$ target space, and can be equivalently described as a U(1) gauge theory coupled to N_b complex scalars of unit charge that satisfy a length constraint; see [20] for a textbook treatment. The action is

$$\mathcal{S} = \frac{N_b}{\mathfrak{g}} \int d^3x \left[|(\partial_\mu - iA_\mu)\phi_\alpha|^2 + i\lambda(|\phi_\alpha|^2 - 1) \right], \quad (1.1)$$

where $\alpha = 1 \dots N_b$, λ is a Lagrange multiplier imposing the length constraint, and \mathfrak{g} is a coupling constant. This theory becomes critical provided that one tunes the coupling to $\mathfrak{g} = \mathfrak{g}_c$ for some \mathfrak{g}_c .

The interest in monopole operators in this theory is motivated by their interpretation as order parameters for the valence bond solid (VBS) order of quantum antiferromagnets [21–23]. The quantum antiferromagnets are defined on bipartite lattices in two spatial dimensions, and have a global $SU(N_b)$ symmetry. Each site of the first (second) sublattice has states transforming under the fundamental (anti-fundamental) of $SU(N_b)$. The sites interact via short-range exchange interactions with $SU(N_b)$ symmetry. There is no explicit reference to a gauge field in the lattice Hamiltonian. Nevertheless, when the spin states on each site are represented in terms of ‘parton’ degrees of freedom, a U(1) gauge field, A_μ , emerges in the path integral formulation in the $1/N_b$ expansion. The partons become the ϕ_α matter fields in this gauge theory. As their exchange constants are varied, such antiferromagnets can exhibit ground states with two distinct broken symmetries. First, there is the state with antiferromagnetic order, in which the $SU(N_b)$ symmetry is broken by the condensation of

ϕ_α : this is the Higgs phase of the U(1) gauge theory. Second, we have the state with VBS order in which $SU(N_b)$ symmetry is preserved but a lattice rotation symmetry is broken. In the U(1) gauge theory, this state appears initially as a Coulomb phase; however, the confinement of the U(1) gauge theory by the proliferation of monopoles leads to the appearance of VBS order in the lattice antiferromagnet. This is a consequence of subtle Berry phases associated with the monopole tunneling events in the lattice antiferromagnet, which endow the monopole operators with non-trivial transformations under lattice symmetry operations, identical to those of the VBS order. The quantum phase transition between these two states of the lattice antiferromagnet has been argued to be continuous [8, 9], and described by the $\mathbb{C}\mathbb{P}^{N_b-1}$ theory in (1.1), with monopoles suppressed at the quantum critical point; the critical point is therefore ‘deconfined’.

From the perspective of the $\mathbb{C}\mathbb{P}^{N_b-1}$ theory, this connection to the VBS order of the antiferromagnet is powerful because it allows the monopole operators to be expressed as simple, local, gauge-invariant operators of the lattice model. Moreover, the couplings of the lattice model can be chosen to avoid the ‘sign’ problem of quantum Monte Carlo, and this allows efficient studies on large lattices of the $\mathbb{C}\mathbb{P}^{N_b-1}$ CFT at the deconfined critical point between the Higgs and VBS phases [24–28]. Block *et al.* [27, 28] have obtained the scaling dimensions of VBS operators on a number of lattice antiferromagnets, and here we will compare their results with the $1/N_b$ expansion for the scaling dimensions of the monopole operators in the $\mathbb{C}\mathbb{P}^{N_b-1}$ CFT.

Unlike the lattice antiferromagnet, the monopole operators of the $\mathbb{C}\mathbb{P}^{N_b-1}$ field theory are not defined simply as products of fields that appear in the Lagrangian (1.1). Instead, the monopole operators appear as singular boundary conditions that these fields must obey at the point where the monopole operator is inserted [29, 30]. One way of defining monopole operators is the following. The $\mathbb{C}\mathbb{P}^{N_b-1}$ model has $SU(N_b) \times U(1)_{\text{top}}$ global symmetry. Under the action of $SU(N_b)$, the charged scalar fields ϕ_α transform in the fundamental representation. The $U(1)_{\text{top}}$ factor in the global symmetry group is a topological symmetry whose conserved current is

$$j_\mu = \frac{1}{4\pi} \epsilon_{\mu\nu\rho} F^{\nu\rho}. \quad (1.2)$$

The Dirac quantization condition implies that the conserved charge $q = \int d^2x j_0$ satisfies $q \in \mathbb{Z}/2$. It should be noted that our definition of q here differs by a factor of 2 from earlier work [23, 27, 31].

One can define the monopole operators as operators that have non-vanishing $U(1)_{\text{top}}$

charge q . For each q , we will focus on the monopole operator \mathcal{M}_q with the lowest scaling dimension. The other operators in the same topological charge sector can be thought intuitively as products between the monopole operator with lowest scaling dimension and more conventional operators from the $q = 0$ sector. Throughout this paper, we assume without loss of generality that $q \geq 0$. The physical quantities that we compute depend only on $|q|$.

We aim to determine the scaling dimension of the monopole operator \mathcal{M}_q with $U(1)_{\text{top}}$ charge q to next-to-leading order in $1/N_b$. The most convenient way of performing this computation is to use the state-operator correspondence, under which a local operator inserted at the origin of \mathbb{R}^3 is mapped to a state of the CFT on the conformally-flat background $S^2 \times \mathbb{R}$. The scaling dimension of the monopole operator \mathcal{M}_q is therefore mapped to the ground state energy on S^2 in the sector with magnetic flux $\int F = 4\pi q$ through the S^2 [29–31]. (See also [23] for a different approach to computing scaling dimensions of monopole operators.) As is standard in thermodynamics, this ground state energy Δ_q can be related to the partition function on $S^2 \times \mathbb{R}$ via

$$\Delta_q = - \lim_{\beta \rightarrow \infty} \frac{1}{\beta} \log Z_q(\beta) \equiv - \log Z_q^{S^2 \times \mathbb{R}} \equiv \mathcal{F}_q, \quad (1.3)$$

where in the middle equality we regularized the $S^2 \times \mathbb{R}$ partition function by compactifying the \mathbb{R} direction into a large circle of circumference β .

In the case at hand, the S^2 ground state energy in the presence of $4\pi q$ magnetic flux is easily computed at leading order in N_b , where the Lagrange multiplier field λ and the gauge field don't fluctuate and assume a saddle point configuration that minimizes this energy. It is reasonable to assume that the large N_b saddle point (for both the magnetic flux through S^2 and the value assumed by the Lagrange multiplier field) is rotationally-symmetric. At leading order in N_b , the ground state energy on S^2 comes from performing the Gaussian integral over the matter fields [23]. The $1/N_b$ correction to this result takes into account the Gaussian fluctuations of the gauge field as well as of those of λ . We perform an analysis of these fluctuations around the rotationally-invariant saddle point; by computing their determinant, we extract the $1/N_b$ correction to the scaling dimension of \mathcal{M}_q . Our results for $\mathcal{F}_{1/2}$ are compared with numerical studies of the lattice antiferromagnet in Fig. 1: notice that the agreement at the largest value of $N_b = 10$ is quite encouraging, although the uncertainties in the quantum Monte Carlo values are significant.

Similar calculations were performed in theories with fermionic matter in [32, 33].¹ (See

¹The calculation in the case with scalar matter performed in this paper is technically more challenging than that for fermionic matter due to additional UV divergences. We renormalize these divergences using




Figure 1: The scaling dimension of the $q = 1/2$ monopole operator, $\mathcal{F}_{1/2}$. The full line is the $N_b = \infty$ result (Ref. [23]), and the dashed line is the leading $1/N_b$ correction computed in the present paper (see Table 2). The quantum Monte Carlo results are for lattice antiferromagnets with global $SU(N_b)$ symmetry on the square (Refs. [24,25]), honeycomb (Ref. [27]), and rectangular (Ref. [27]) lattices.

also [35] where only the fluctuations of the Lagrange multiplier field were calculated with the purpose of studying monopole insertions in theories with global $U(1)$ symmetry.) In [36,37], monopole operators were studied holographically. For studies of monopole operators in supersymmetric theories, see, for instance, [38–43].

Our interest in the fluctuations around the spherically-symmetric saddle point is not purely to calculate the $1/N_b$ correction to the scaling dimension of \mathcal{M}_q . It turns out that some of these fluctuations correspond to negative modes, in which case we conclude that the spherically-symmetric configuration for the gauge field and the Lagrange multiplier field λ does not represent the dominant saddle point configuration. In fact, as we will show, we encounter negative modes for $q > 1$. In other words, in this case it is only the lowest two monopole operators, with $q = 1/2$ and $q = 1$, for which a spherically-symmetric saddle is the dominant one. When $q > 1$, the leading order computation of the S^2 ground state energy that we described above represents only an upper bound on the scaling dimension of the monopole operators. To obtain the correct leading order results at large N_b , one would

zeta-function techniques and devote a separate paper to their thorough study in other renormalization schemes [34].

have to first find the dominant (non-spherically symmetric) saddle point configuration for the gauge field and the Lagrange multiplier field λ . We leave this task to future work.

The rest of this paper is organized as follows. In Section 2, we set up our computation. In Section 3, we review the leading order analysis at large N_b . In Section 4, we examine the $1/N_b$ corrections around the spherically-symmetric saddle points of the effective action for the gauge field and Lagrange multiplier. In Section 5 we perform a perturbative stability analysis of our spherically-symmetric saddle points and find that, as mentioned above, the rotationally-symmetric saddles with $q > 1$ are unstable. We end with concluding remarks in Section 6. Several technical details of our computation are included in the Appendices.

2 Setup

In order to study the large N_b limit of the \mathbb{CP}^{N_b-1} theory, it is convenient to rescale the fields such that the action (1.1), appropriately generalized to that on an arbitrary conformally-flat space with metric tensor $g_{\mu\nu}$, takes the form

$$\mathcal{S} = \frac{1}{\mathfrak{g}} \int d^3x \sqrt{g(x)} \left[g^{\mu\nu} [(\nabla_\mu + iA_\mu)\phi_\alpha^*][(\nabla_\nu - iA_\nu)\phi_\alpha] + \frac{\mathcal{R}}{8} |\phi_\alpha|^2 + i\lambda(|\phi_\alpha|^2 - N_b) \right], \quad (2.1)$$

where \mathcal{R} is the Ricci scalar.² In this paper we will work on $S^2 \times \mathbb{R}$, which we parameterize by coordinates $x \equiv (\theta, \phi, \tau)$.

The monopole scaling dimension is equal to the ground state energy \mathcal{F}_q on $S^2 \times \mathbb{R}$ in the presence of a magnetic flux $\int F = 4\pi q$ through the S^2 . Our main task is to determine the $1/N_b$ expansion of this ground state energy, which we write as

$$\mathcal{F}_q = N_b \left(\mathcal{F}_q^\infty + \frac{1}{N_b} \delta\mathcal{F}_q + \mathcal{O}(1/N_b^2) \right). \quad (2.2)$$

When $q = 0$, the corresponding ground state energy \mathcal{F}_0 is nothing but the scaling dimension of the unit operator. We therefore must have $\mathcal{F}_0 = 0$.

It is not hard to see that at large N_b , the fluctuations of λ and of A_μ around any saddle point configuration are suppressed. Indeed, upon integrating out the scalars ϕ_α in the action (2.1), one obtains an effective action for the gauge field and Lagrange multiplier given by

$$\mathcal{S}_{\text{eff}}[A_\mu, i\lambda] = N_b \left[\text{tr} \log \left(-(\nabla_\mu - iA_\mu)^2 + \frac{1}{4} + i\lambda \right) - \frac{i}{\mathfrak{g}} \int d^3x \sqrt{g} \lambda \right]. \quad (2.3)$$

²We could have absorbed the conformal mass term $\frac{\mathcal{R}}{8} |\phi_\alpha|^2$ by shifting $i\lambda$, but we chose not to.

Let's expand A_μ and λ around a saddle point by writing³

$$\begin{aligned} A_\mu &= \mathcal{A}_\mu^q + a_\mu, \\ i\lambda &= \mu_q^2 + i\sigma, \end{aligned} \tag{2.4}$$

where a_μ and σ are fluctuations around the saddle point configuration $A_\mu = \mathcal{A}_\mu$ and $i\lambda = \mu_q^2$. As can be easily seen from (2.3), the effective action for these fluctuations is proportional to N_b , so their typical size is of order $1/\sqrt{N_b}$ and are therefore suppressed at large N_b . To leading order in $1/N_b$, it is therefore correct to set $a_\mu = \sigma = 0$, provided that the background values \mathcal{A}_μ^q and μ_q^2 are such that the saddle point conditions

$$\left. \frac{\delta \mathcal{S}_{\text{eff}}[A_\mu, i\lambda]}{\delta A_\mu} \right|_{\sigma=a_\mu=0} = \left. \frac{\delta \mathcal{S}_{\text{eff}}[A_\mu, i\lambda]}{\delta \lambda} \right|_{\sigma=a_\mu=0} = 0 \tag{2.5}$$

are obeyed. One can then develop the $1/N_b$ expansion to higher orders by integrating over the fluctuations a_μ and σ using the effective action (2.3).

In this paper we will focus only on saddles that are rotationally-invariant on S^2 and translationally-invariant along \mathbb{R} . These conditions imply that μ_q^2 is a constant and that, in the sector of monopole flux $\int F = 4\pi q$, the background magnetic field $\mathcal{F}^q = d\mathcal{A}^q$ is uniformly distributed over S^2 :

$$\mathcal{F}^q = q \sin \theta d\theta \wedge d\phi. \tag{2.6}$$

One can choose a gauge where the background gauge potential \mathcal{A}^q can be written as

$$\mathcal{A}^q = q(1 - \cos \theta)d\phi. \tag{2.7}$$

(This expression is well-defined everywhere away from the South pole at $\theta = \pi$.) The saddle point condition (2.5) is satisfied provided that the constant μ_q^2 is chosen such that it minimizes the value of the effective action evaluated when $a_\mu = \sigma = 0$. In other words, the equation that determines μ_q^2 is

$$\frac{\partial \mathcal{S}_{\text{eff}}[\mathcal{A}_\mu^q, \mu_q^2]}{\partial \mu_q^2} = 0. \tag{2.8}$$

This equation depends non-trivially on q , and hence so does μ_q^2 .

³In terms of the quantity a_q^2 introduced in [31], we have $a_q^2 = \mu_q^2 - q^2$.

In the next section, we calculate the coefficient \mathcal{F}_q^∞ by simply evaluating \mathcal{S}_{eff} at the saddle point, while in Section 4, we compute the correction $\delta\mathcal{F}_q$ from the functional determinant of the fluctuations around this saddle point. As explained in Section 4, our computations are valid only for $q = 1/2$ and $q = 1$. For $q \geq 3/2$, some of the fluctuations have negative directions, rendering the saddle (2.4) not a local minimum of the effective action. In these cases, the leading order computation from the next section should be thought of as an upper bound on the scaling dimension of the corresponding monopole operators.⁴

3 $N_b = \infty$ theory

At leading order in N_b , one can identify

$$N_b \mathcal{F}_q^\infty = \mathcal{S}_{\text{eff}}[\mathcal{A}_\mu^q, \mu_q^2], \quad (3.1)$$

evaluated for the value of μ_q^2 that solves (2.8), or equivalently at large N_b

$$\frac{\partial \mathcal{F}_q^\infty}{\partial \mu_q^2} = 0, \quad (3.2)$$

and with the coupling \mathfrak{g} tuned to the critical value $\mathfrak{g} = \mathfrak{g}_c$. In other words,

$$\mathcal{F}_q^\infty = \text{tr} \log \left(-(\nabla_\mu - i\mathcal{A}_\mu^q)^2 + \mu_q^2 + \frac{1}{4} \right) - \frac{4\pi}{\mathfrak{g}_c} \mu_q^2. \quad (3.3)$$

Using the fact that the eigenvalues of the gauge-covariant Laplacian on S^2 in the presence of magnetic flux $4\pi q$ are $j(j+1) - q^2$ [44, 45], we obtain

$$\mathcal{F}_q^\infty = \int \frac{d\omega}{2\pi} \sum_{j=q}^{\infty} (2j+1) \log [\omega^2 + (j+1/2)^2 + \mu_q^2 - q^2] - \frac{4\pi}{\mathfrak{g}_c} \mu_q^2. \quad (3.4)$$

The first term in this expression is divergent and requires regularization. The second term is also divergent because, as we explain shortly, the inverse critical coupling $1/\mathfrak{g}_c$ diverges linearly, and so the second term in (3.4) cancels part of the divergence in the first term. To be explicit, let us deduce an expression for $1/\mathfrak{g}_c$. The saddle point condition (3.2) at $q = 0$

⁴The leading order computation provides an upper bound, as the dominant saddle corresponding to the non-spherically symmetric configuration mentioned in the introduction necessarily has lower free energy.

can be written as

$$\frac{4\pi}{\mathfrak{g}} = \int \frac{d\omega}{2\pi} \sum_{j=0}^{\infty} (2j+1) \frac{1}{\omega^2 + (j+1/2)^2 + \mu_0^2}. \quad (3.5)$$

The theory is critical when the correlators on $S^2 \times \mathbb{R}$ are those obtained by conformally mapping the power-law correlators on \mathbb{R}^3 . For a scalar field ϕ_α , this prescription yields a conformally coupled scalar on $S^2 \times \mathbb{R}$, for which $\mu_0 = 0$. Hence criticality is achieved when, to leading order in N_b , we have $\mathfrak{g} = \mathfrak{g}_c$ and $\mu_0^2 = 0$, with

$$\frac{4\pi}{\mathfrak{g}_c} = \int \frac{d\omega}{2\pi} \sum_{j=0}^{\infty} (2j+1) \frac{1}{\omega^2 + (j+1/2)^2}. \quad (3.6)$$

After substituting (3.6) into (3.4), the resulting expression is still divergent, but can be rendered finite using, for instance, zeta-function regularization as in [35], or Pauli-Villars regularization as in [23,31]. We will not repeat that calculation here. The regularized ground state energy coefficient \mathcal{F}_q^∞ is

$$\begin{aligned} \mathcal{F}_q^\infty = 2 \sum_{j=q}^{\infty} & \left[(j+1/2) [(j+1/2)^2 + \mu_q^2 - q^2]^{1/2} - (j+1/2)^2 - \frac{1}{2}(\mu_q^2 - q^2) \right] \\ & - q\mu_q^2 + \frac{q(1+2q^2)}{6}. \end{aligned} \quad (3.7)$$

This expression can easily be evaluated numerically for any μ_q^2 . Note that the same procedure gives $\mathcal{F}_0^\infty = 0$, as required by conformal symmetry.

As mentioned above, the value of μ_q^2 can be obtained from the saddle-point equation (2.8), which yields

$$\sum_{j=q}^{\infty} \left(\frac{j+1/2}{\sqrt{(j+1/2)^2 + \mu_q^2 - q^2}} - 1 \right) - q = 0. \quad (3.8)$$

Note that one can obtain this equation directly by differentiating (3.4) with respect to μ_q^2 without the need of zeta-function regularization. Upon substituting the solution of (3.8) into (3.7), one obtains the values of \mathcal{F}_q^∞ given in Table 1. These values agree precisely with those obtained in [23] by a very different method.⁵

⁵The quantity \mathcal{F}_q^∞ should be identified with $2\rho_{2q}$ in [23], while our μ_q^2 should be identified with $-C_{2q}$ in [23].

q	μ_q^2	\mathcal{F}_q^∞
0	0	0
1/2	-0.199806	0.1245922
1	-0.397830	0.3110952
3/2	-0.595457	0.5440693
2	-0.792936	0.8157878
5/2	-0.990344	1.1214167

Table 1: A few values for the parameters μ_q^2 and \mathcal{F}_q^∞ .

4 $1/N_b$ corrections

In this section we compute the next to leading order correction to the dimensions of monopole operators. The systematics of the calculation are presented in the first four subsections, with the numerical results presented in Section 4.5.

4.1 Effective action at quadratic level

To obtain the leading $1/N_b$ correction $\delta\mathcal{F}_q$ to the result of the previous section, one should consider the quadratic fluctuations of the gauge field and of the Lagrange multiplier around the saddle (2.4). Expanding (2.3) at small a_μ and σ , one can write the quadratic term in the effective action as

$$\begin{aligned}
\mathcal{S}_{\text{eff}}^{(2)} &= \mathcal{S}_{\sigma\sigma}^{(2)} + \mathcal{S}_{aa}^{(2)} + \mathcal{S}_{a\sigma}^{(2)}, \\
\mathcal{S}_{\sigma\sigma}^{(2)} &= \frac{N_b}{2} \int d^3x d^3x' \sqrt{g(x)} \sqrt{g(x')} \sigma(x) D^q(x, x') \sigma(x'), \\
\mathcal{S}_{aa}^{(2)} &= \frac{N_b}{2} \int d^3x d^3x' \sqrt{g(x)} \sqrt{g(x')} a_\mu(x) K^{q, \mu\mu'}(x, x') a_{\mu'}(x'), \\
\mathcal{S}_{a\sigma}^{(2)} &= N_b \int d^3x d^3x' \sqrt{g(x)} \sqrt{g(x')} \sigma(x) F^{q, \mu'}(x, x') a_{\mu'}(x'),
\end{aligned} \tag{4.1}$$

where D^q , K^q , and F^q are integration kernels whose expressions will be given shortly. This effective action is non-local because it was obtained after integrating out the fields ϕ_α , which are massless. The kernels appearing in (4.1) can be written in terms of correlators of $|\phi_\alpha|^2$ and of the current

$$J_\mu = i [\phi_\alpha^* (\nabla_\mu - i\mathcal{A}_\mu^q) \phi_\alpha - \phi_\alpha (\nabla_\mu + i\mathcal{A}_\mu^q) \phi_\alpha^*], \tag{4.2}$$

as

$$\begin{aligned}
N_b D^q(x, x') &= \frac{1}{\mathfrak{g}^2} \langle |\phi_\alpha(x)|^2 |\phi_\alpha(x')|^2 \rangle_q, \\
N_b K^{q, \mu\mu'}(x, x') &= -\frac{1}{\mathfrak{g}^2} \langle J^\mu(x) J^{\mu'}(x') \rangle_q + \frac{2}{\mathfrak{g}} g^{\mu\mu'} \delta(x - x') \langle |\phi_\alpha(x)|^2 \rangle_q, \\
N_b F^{q, \mu'}(x, x') &= -\frac{i}{\mathfrak{g}^2} \langle |\phi_\alpha(x)|^2 J^{\mu'}(x') \rangle_q,
\end{aligned} \tag{4.3}$$

where the delta-function contains a factor of $1/\sqrt{g(x)}$ in its definition. The correlators in (4.3) are evaluated under the assumption that the gauge field and Lagrange multiplier are non-dynamical and fixed at their background values $A_\mu = \mathcal{A}_\mu^q$ and $i\lambda = \mu_q^2$. The subscript q on the angle brackets in the expressions above serves as a reminder of these assumptions.

Performing the Gaussian integral over a_μ and σ , we can write the coefficient $\delta\mathcal{F}_q$ in (2.2) as

$$\delta\mathcal{F}_q = \frac{1}{2} \log \det' M^q, \tag{4.4}$$

where we defined the matrix of kernels

$$M^q(x, x') \equiv \begin{pmatrix} D^q(x, x') & F_\tau^q(x, x') & F_{i'}^q(x, x') \\ F^{q\tau}(x, x') & K^{q\tau}{}_{\tau'}(x, x') & K^{q\tau}{}_{i'}(x, x') \\ F^{qi}(x, x') & K^{qi}{}_{i'}(x, x') & K^{qi}{}_{i'}(x, x') \end{pmatrix}, \tag{4.5}$$

with $i = \theta, \phi$ and the primed indices contracting with the index of the field at x' . The prime on the determinant in (4.4) means that when computing the functional determinant we should ignore the zero eigenvalues that are required to be present due to gauge invariance.⁶ Our goal in the rest of this section is to calculate a regularized version of this determinant, thus obtaining $\delta\mathcal{F}_q$.

4.2 Eigenvalues of the integration kernels

One can start evaluating the expressions in (4.3) in terms of the $N_b = \infty$ limit of the Green's function $G^q(x, x')$ for the complex scalars, which is defined by

$$\langle \phi_\alpha(x) \phi_\beta^*(x') \rangle = \mathfrak{g} \delta_{\alpha\beta} G^q(x, x'). \tag{4.6}$$

⁶For a more detailed treatment of gauge fixing, see [33].

Performing the required Wick contractions in (4.3), we obtain

$$\begin{aligned}
D^q(x, x') &= G^q(x, x')G^{q*}(x, x'), \\
K_{\mu\mu'}^q(x, x') &= D_\mu G^q(x, x')D_{\mu'} G^{q*}(x, x') - G^{q*}(x, x')D_\mu D_{\mu'} G^q(x, x') \\
&\quad + D_\mu G^{q*}(x, x')D_{\mu'} G^q(x, x') - G^q(x, x')D_\mu D_{\mu'} G^{q*}(x, x') \\
&\quad + 2g_{\mu\nu}\delta(x - x')G^q(x, x), \\
F_{\mu'}^q(x, x') &= G^q(x, x')D_{\mu'} G^{q*}(x, x') - G^{q*}(x, x')D_{\mu'} G^q(x, x'),
\end{aligned} \tag{4.7}$$

where $D_\mu = \partial_\mu - i\mathcal{A}_\mu^q(x)$ and $D_{\mu'} = \partial_{\mu'} + i\mathcal{A}_\mu^q(x')$ denote the gauge-covariant derivatives in the presence of the background gauge field.

In order to calculate the eigenvalues of the matrix of kernels (4.5) required for (4.4), we make use of the S^2 rotational symmetry and the translational symmetry along \mathbb{R} . These symmetries imply that the eigenvectors of this matrix are of the form $e^{-i\omega\tau}$ times an appropriate (scalar or vector) spherical harmonic on S^2 . We will need the usual spherical harmonics $Y_{jm}(\theta, \phi)$, as well as the vector harmonics

$$\begin{aligned}
\mathcal{X}_{i,jm} &= \frac{1}{\sqrt{j(j+1)}}\partial_i Y_{jm}, \\
\mathcal{Y}_{jm}^i &= \frac{1}{\sqrt{j(j+1)}}\frac{\epsilon^{ik}}{\sqrt{g}}\partial_k Y_{jm},
\end{aligned} \tag{4.8}$$

where $i, k = \theta, \phi$, and $\epsilon^{\theta\phi} = -\epsilon^{\phi\theta} = 1$. We can decompose the Lagrange multiplier fluctuation σ and the gauge field fluctuation a_μ in terms of these modes. Because, σ is a scalar field, we only need the usual spherical harmonics for its mode expansion. a_τ is also decomposed in terms of Y_{jm} , while a_i is decomposed using the vector harmonics $\mathcal{X}_{i,jm}$ and \mathcal{Y}_{jm}^i . We will refer to the former vector harmonics modes as E modes and to the latter as B modes, as they are the S^2 analogs of E and B modes familiar from other contexts: E and B modes have vanishing curl and divergence, respectively, and they transform like the E and B field under parity.

We can expand each of the kernels in Fourier modes as

$$D^q(x, x') = \int \frac{d\omega}{2\pi} \sum_{jm} D_j^q(\omega) Y_{jm}(\theta, \phi) Y_{jm}^*(\theta', \phi') e^{-i\omega(\tau - \tau')} \tag{4.9}$$

$$\begin{aligned}
K_{\tau\tau}^q(x, x') &= \int \frac{d\omega}{2\pi} \sum_{jm} K_j^{q,\tau\tau}(\omega) Y_{jm}(\theta, \phi) Y_{jm}^*(\theta', \phi') e^{-i\omega(\tau-\tau')} \\
K_{ii'}^q(x, x') &= \int \frac{d\omega}{2\pi} \sum_{jm} \left[K_j^{q,EE}(\omega) \mathcal{X}_{i,jm}(\theta, \phi) \mathcal{X}_{i',jm}^*(\theta', \phi') + K_j^{q,BB}(\omega) \mathcal{Y}_{i,jm}(\theta, \phi) \mathcal{Y}_{i',jm}^*(\theta', \phi') \right. \\
&\quad \left. + K_j^{q,EB}(\omega) \mathcal{X}_{i,jm}(\theta, \phi) \mathcal{Y}_{i',jm}^*(\theta', \phi') + K_j^{q,EB*}(\omega) \mathcal{Y}_{i,jm}(\theta, \phi) \mathcal{X}_{i',jm}^*(\theta', \phi') \right] e^{-i\omega(\tau-\tau')} \\
K_{\tau i'}^q(x, x') &= \int \frac{d\omega}{2\pi} \sum_{jm} \left[K_j^{q,\tau E}(\omega) Y_{jm}(\theta, \phi) \mathcal{X}_{i',jm}^*(\theta', \phi') + K_j^{q,\tau B}(\omega) Y_{jm}(\theta, \phi) \mathcal{Y}_{i',jm}^*(\theta', \phi') \right] e^{-i\omega(\tau-\tau')}
\end{aligned} \tag{4.10}$$

$$\begin{aligned}
F_{\tau}^q(x, x') &= \int \frac{d\omega}{2\pi} \sum_{jm} F_j^{q,\tau}(\omega) Y_{jm}(\theta, \phi) Y_{jm}^*(\theta', \phi') e^{-i\omega(\tau-\tau')} \\
F_{i'}^q(x, x') &= \int \frac{d\omega}{2\pi} \sum_{jm} \left[F_j^{q,E}(\omega) Y_{jm}(\theta, \phi) \mathcal{X}_{i',jm}^*(\theta', \phi') + F_j^{q,B}(\omega) Y_{jm}(\theta, \phi) \mathcal{Y}_{i',jm}^*(\theta', \phi') \right] e^{-i\omega(\tau-\tau')}
\end{aligned} \tag{4.11}$$

and form the matrix of coefficients

$$\mathbf{M}_j^q(\omega) = \left(\begin{array}{c|ccc} D_j^q(\omega) & F_j^{q,B}(\omega) & F_j^{q,\tau}(\omega) & F_j^{q,E}(\omega) \\ \hline F_j^{q,B*}(\omega) & K_j^{q,BB}(\omega) & K_j^{q,\tau B}(\omega) & K_j^{q,EB}(\omega) \\ F_j^{q,\tau*}(\omega) & K_j^{q,\tau B*}(\omega) & K_j^{q,\tau\tau}(\omega) & K_j^{q,\tau E}(\omega) \\ \hline F_j^{q,E*}(\omega) & K_j^{q,EB*}(\omega) & K_j^{q,\tau E*}(\omega) & K_j^{q,EE}(\omega) \end{array} \right). \tag{4.12}$$

Note that S^2 rotational symmetry implies that these coefficients do not depend on the quantum number m .

The entries of this matrix are related by gauge invariance and CP symmetry. Gauge invariance of the kernels in position space imply that at separated points⁷

$$\begin{aligned}
\nabla^\mu K_{\mu\mu'}^q(x, x') &= 0, & \nabla^{\mu'} K_{\mu\mu'}^q(x, x') &= 0, \\
\nabla^{\mu'} F_{\mu'}^q(x, x') &= 0.
\end{aligned} \tag{4.13}$$

Plugging in the decompositions (4.10) and (4.11) in these conservation equations, we obtain that the Fourier space kernel $\mathbf{M}_j^q(\omega)$ should have the following eigenvectors with zero

⁷There are multiple equivalent ways to see that these equations are true. Using the definitions of the kernels in terms of correlators (4.3) they are the consequences of the Ward identity $\nabla^\mu J_\mu(x) = 0$. Alternatively, (4.1) should be zero for a pure gauge configuration $a_\mu = \nabla_\mu \alpha(x)$, for arbitrary $\alpha(x)$. Partial integration readily gives (4.13).

eigenvalue:

$$\left(0 \mid 0 \quad i\omega \quad -\sqrt{j(j+1)}\right) \mathbf{M}_j^q(\omega) = 0, \quad \mathbf{M}_j^q(\omega) \left(0 \mid 0 \quad -i\omega \quad -\sqrt{j(j+1)}\right)^T = 0, \quad (4.14)$$

where the pure gauge eigenvector is written in $(\sigma \mid B, \tau, E)$ components, just like $\mathbf{M}_j^q(\omega)$ in (4.12). From (4.14) we can express $K_j^{q,\tau E}(\omega)$ and $K_j^{q,EE}(\omega)$ in terms of $K_j^{q,\tau\tau}(\omega)$.

The second restriction on the entries of $\mathbf{M}_j^q(\omega)$ comes from the CP invariance of the theory and the monopole background around which we are working. Under CP the modes of σ and the B modes of a_μ transform in the same way, while the τ and E modes acquire a relative minus sign. Because the effective action (4.1) is invariant under CP , we conclude that there is no mixing between σ, B and τ, E modes.

These constraints imply that $\mathbf{M}_j^q(\omega)$ takes a block diagonal form. For $j > 0$,

$$\mathbf{M}_j^q(\omega) = \left(\begin{array}{c|cc} D_j^q(\omega) & F_j^{q,B}(\omega) & 0 & 0 \\ F_j^{q,B^*}(\omega) & K_j^{q,BB}(\omega) & 0 & 0 \\ \hline 0 & 0 & K_j^{q,\tau\tau}(\omega) & \frac{-i\omega}{\sqrt{j(j+1)}} K_j^{q,\tau\tau}(\omega) \\ 0 & 0 & \frac{i\omega}{\sqrt{j(j+1)}} K_j^{q,\tau\tau}(\omega) & \frac{\omega^2}{j(j+1)} K_j^{q,\tau\tau}(\omega) \end{array} \right). \quad (4.15)$$

This matrix has eigenvalues:

$$\lambda_{\pm}^q = \frac{(D_j^q(\omega) + K_j^{q,BB}(\omega)) \pm \sqrt{4|F_j^{q,B}(\omega)|^2 + (D_j^q(\omega) - K_j^{q,BB}(\omega))^2}}{2}, \quad (4.16)$$

$$\lambda_E^q = \frac{j(j+1) + \omega^2}{j(j+1)} K_j^{q,\tau\tau},$$

as well as a zero eigenvalue corresponding to a pure gauge mode.

When $j = 0$, the harmonics \mathcal{X}_{jm} and \mathcal{Y}_{jm} are not defined, so the matrix $\mathbf{M}_j^q(\omega)$ reduces to the 2×2 matrix

$$\mathbf{M}_0^q(\omega) = \left(\begin{array}{c|c} D_0^q(\omega) & 0 \\ \hline 0 & K_0^{q,\tau\tau}(\omega) \end{array} \right). \quad (4.17)$$

In addition, the only remaining vector harmonic Y_{00} is a constant on S^2 , and can be gauged away. Thus gauge invariance imposes $K_0^{q,\tau\tau}(\omega) = 0$, and the only non-vanishing eigenvalue is $D_0^q(\omega)$.

We will derive expressions for the entries of the matrices (4.15)–(4.17) shortly. After doing so, we can calculate $\delta\mathcal{F}_q$ from (4.4). It is convenient to subtract $\delta\mathcal{F}_0 = 0$ from $\delta\mathcal{F}_q$.⁸ The expression we would like to calculate becomes:

$$\delta\mathcal{F}_q = \frac{1}{2} \int \frac{d\omega}{2\pi} \sum_{j=0}^{\infty} (2j+1) \log \frac{\det' M^q}{\det' M^0}. \quad (4.18)$$

Using the expression for the eigenvalues from (4.16) and that $F_j^{0,B}(\omega) = 0$ by parity symmetry, then (4.18) becomes:

$$\delta\mathcal{F}_q = \frac{1}{2} \int \frac{d\omega}{2\pi} \left[\log \frac{D_0^q(\omega)}{D_0^0(\omega)} + \sum_{j=1}^{\infty} (2j+1) \log \frac{K_j^{q,\tau\tau}(\omega) \left[D_j^q(\omega) K_j^{q,BB}(\omega) - |F_j^{q,B}(\omega)|^2 \right]}{D_j^0(\omega) K_j^{0,\tau\tau}(\omega) K_j^{0,BB}(\omega)} \right]. \quad (4.19)$$

Explicit expressions for the coefficients in (4.15) can be obtained by inverting (4.9)–(4.11). Let us explain how to do so for $D_j^q(\omega)$ first, and leave the details of how to perform analogous computations for the K and F kernels to Appendix A. For $D_j^q(\omega)$ we obtain:

$$D_j^q(\omega) 2\pi \delta(\omega - \omega') = \int d^3x d^3x' \sqrt{g(x)} \sqrt{g(x')} Y_{jm}^*(\theta, \phi) D^q(x, x') Y_{jm}(\theta', \phi') e^{i(\omega\tau - \omega'\tau')}. \quad (4.20)$$

Since the LHS is independent of m , we can average the RHS over all possible values of m . After performing the average, the RHS becomes invariant under performing a combined rotation in (θ, ϕ) and (θ', ϕ') , so we can take the limit $\theta' \rightarrow 0$. We can also use that $D^q(x, x')$ depends only on $\tau - \tau'$ to set $\omega' = \omega$ and remove the τ' integral. The simplified expression is

$$D_j^q(\omega) = \frac{4\pi}{2j+1} \int d^3x \sqrt{g(x)} \lim_{\substack{\theta' \rightarrow 0 \\ \tau' \rightarrow 0}} \sum_{m=-j}^j Y_{jm}^*(\theta, \phi) D^q(x, x') Y_{jm}(\theta', \phi') e^{i\omega\tau}. \quad (4.21)$$

It is only the $m = 0$ term that contributes to the sum. Analogous formulas for the K and F kernels are given in (A.3). Using explicit formulas for the spherical harmonics, (4.21) can

⁸Note that as the gauge fixing condition is independent of the monopole background, any possible contribution from the Faddeev-Popov ghosts cancels after subtracting the vacuum contribution.

be simplified further to

$$D_j^q(\omega) = \int d^3x \sqrt{g(x)} P_j(\cos \theta) D^q(x, 0) e^{i\omega\tau}, \quad (4.22)$$

where by $x' = 0$ we mean the limit $\tau', \theta' \rightarrow 0$. Similar expressions (albeit more complicated) can be obtained for the other coefficients appearing in (4.15)–(4.17).

4.3 Kernels at $q = 0$

When $q = 0$, one can obtain closed form formulas for the entries of the matrix (4.15)–(4.17). In this case, the Green's function $G^0(x, x')$ can be obtained by conformally mapping the \mathbb{R}^3 one, namely $1/(4\pi|x - x'|)$, and from G^0 one can construct position-space expressions for all the kernels in (4.7). The conformal mapping from flat space gives

$$G^0(x, x') = \frac{1}{4\pi\sqrt{2}(\cosh(\tau - \tau') - \cos \gamma)}, \quad (4.23)$$

where γ is the angle between the 2 points on S^2 :

$$\cos \gamma = \cos \theta \cos \theta' + \sin \theta \sin \theta' \cos(\phi - \phi'). \quad (4.24)$$

Since $G^0(x, x')$ is real, eq. (4.7) implies that $F^0(x, x') = 0$, and consequently $F_j^{0,B}(\omega) = 0$.

4.3.1 The D kernel at $q = 0$

Plugging (4.7) and (4.23) into (4.22), we obtain

$$D_j^0(\omega) = \frac{1}{8\pi} \int_{-\infty}^{\infty} d\tau \int_0^\pi \sin \theta d\theta \frac{e^{i\omega\tau} P_j(\cos \theta)}{2(\cosh \tau - \cos \theta)}. \quad (4.25)$$

This expression can be evaluated by first performing the θ integral, which yields

$$D_j^0(\omega) = \frac{1}{8\pi} \int_{-\infty}^{\infty} d\tau e^{i\omega\tau} Q_j(\cosh \tau), \quad (4.26)$$

where $Q_j(x)$ is the Legendre function of the second kind. We can then expand the remaining integrand at large τ ,

$$D_j^0(\omega) = \frac{1}{8\pi} \int_{-\infty}^{\infty} d\tau \sum_{n=0}^{\infty} \frac{(n+j)! \Gamma(n+1/2)}{n! \Gamma(n+j+3/2)} e^{-(2n+j+1)|\tau|} e^{i\omega\tau}, \quad (4.27)$$

and perform the τ integral term by term. The result can be written as [35]

$$D_j^0(\omega) = \left| \frac{\Gamma((j+1+i\omega)/2)}{4\Gamma((j+2+i\omega)/2)} \right|^2. \quad (4.28)$$

Note that in deriving (4.28) we encountered no divergences in the sums and integrals we performed.

4.3.2 The K kernels at $q = 0$

Next, we aim to find an expression for $K_j^{0,\tau\tau}(\omega)$. While the expression for $K_j^{0,\tau\tau}(\omega)$ that follows from (4.10) is UV divergent (as would be its flat space analog), the following difference is finite:

$$K_j^{0,\tau\tau}(\omega) - K_0^{0,\tau\tau}(0) = \frac{1}{8\pi} \int_{-\infty}^{\infty} d\tau \int_0^\pi \sin\theta d\theta \frac{1 - \cos\theta \cosh\tau}{2(\cosh\tau - \cos\theta)^3} [e^{i\omega\tau} P_j(\cos\theta) - 1]. \quad (4.29)$$

It can be checked that

$$\frac{1 - \cos\theta \cosh\tau}{2(\cosh\tau - \cos\theta)^3} = \nabla_{S^2}^2 \frac{-1}{4(\cosh\tau - \cos\theta)}. \quad (4.30)$$

Substituting (4.30) into (4.29), integrating by parts twice in the sphere directions, and using $\nabla_{S^2}^2 P_j(\cos\theta) = -j(j+1)P_j(\cos\theta)$, one can easily show that

$$K_j^{0,\tau\tau}(\omega) - K_0^{0,\tau\tau}(0) = \frac{j(j+1)}{2} D_j^0(\omega). \quad (4.31)$$

Similarly, it can be shown that

$$\begin{aligned} K_j^{0,BB}(\omega) - K_1^{0,BB}(0) &= \frac{(\omega^2 + j^2)}{2} D_{j-1}^0(\omega) - \frac{1}{2} D_0^0(0), \\ K_j^{0,EE}(\omega) - K_1^{0,EE}(0) &= \frac{\omega^2}{2} D_j^0(\omega), \\ K_j^{0,\tau E}(\omega) &= -i\omega \frac{\sqrt{j(j+1)}}{2} D_j^0(\omega). \end{aligned} \quad (4.32)$$

These expressions are consistent with the requirements of gauge invariance in (4.15). They also agree with the flat-space limit expected at large ω and j , which was obtained in [14]. Also, it follows from (4.17) that at $j = 0$ gauge invariance requires $K_0^{0,\tau\tau}(0) = 0$. From (4.15) we also know that $K_1^{0,EE}(\omega) = \frac{\omega^2}{2} K_1^{0,\tau\tau}(\omega)$. Taking the $\omega \rightarrow 0$ limit implies $K_1^{0,EE}(0) = 0$,

as $K_1^{0,\tau\tau}(0)$ is finite. Assuming these relations, we have

$$\begin{aligned}
K_j^{0,\tau\tau}(\omega) &= \frac{j(j+1)}{2} D_j^0(\omega), \\
K_j^{0,BB}(\omega) &= \frac{(\omega^2 + j^2)}{2} D_{j-1}^0(\omega) + C_0, \\
K_j^{0,EE}(\omega) &= \frac{\omega^2}{2} D_j^0(\omega), \\
K_j^{0,\tau E}(\omega) &= -i\omega \frac{\sqrt{j(j+1)}}{2} D_j^0(\omega),
\end{aligned} \tag{4.33}$$

where the constant C_0 remains to be determined; we will see later around (4.57) that $C_0 = 0$.

4.4 Kernels for general q

For general q , there is no simple closed form expression for the ϕ_α Green's function. One can determine an integral expression for it by first expanding the fields ϕ_α in Fourier modes:

$$\phi_\alpha = \int \frac{d\omega}{2\pi} \sum_{j=q}^{\infty} \sum_{m=-j}^j \Phi_{\alpha,jm}(\omega) Y_{q,jm}(\theta, \phi) e^{-i\omega\tau}, \tag{4.34}$$

where $Y_{q,jm}$ are the monopole spherical harmonics introduced in [44, 45]. At leading order in N_b , the action for the fields ϕ_α becomes

$$\mathcal{S}_\phi = \frac{1}{\mathfrak{g}} \sum_{j=q}^{\infty} \sum_{m=-j}^j \int \frac{d\omega}{2\pi} [\omega^2 + (j + 1/2)^2 + \mu_q^2 - q^2] |\Phi_{\alpha,jm}(\omega)|^2, \tag{4.35}$$

from which we can read off

$$\langle \Phi_{\alpha,jm}(\omega) \Phi_{\beta,j'm'}(\omega')^* \rangle = 2\pi \mathfrak{g} \delta(\omega - \omega') \delta_{\alpha\beta} \delta_{jj'} \delta_{mm'} G_j(\omega), \tag{4.36}$$

with

$$G_j(\omega) \equiv \frac{1}{\omega^2 + (j + 1/2)^2 + \mu_q^2 - q^2}. \tag{4.37}$$

From (4.6), (4.36), (4.34), and (4.37) we can write $G^q(x, x')$ as

$$\begin{aligned} G^q(x, x') &= \sum_{j=q}^{\infty} \int \frac{d\omega}{2\pi} e^{-i\omega(\tau-\tau')} \left[\sum_{m=-j}^j Y_{q,jm}(\theta, \phi) Y_{q,jm}^*(\theta', \phi') \right] G_j(\omega) \\ &= \sum_{j=q}^{\infty} e^{-2iq\Theta} F_{q,j}(\gamma) \frac{e^{-E_{qj}|\tau-\tau'|}}{2E_{qj}}, \end{aligned} \quad (4.38)$$

where in the second line we defined the polynomial in $\cos \gamma$

$$F_{q,j}(\gamma) \equiv \sqrt{\frac{2j+1}{4\pi}} Y_{q,j(-q)}(\gamma, 0), \quad (4.39)$$

and Θ is a phase factor discussed in [45] that can be defined through

$$e^{i\Theta} \cos(\gamma/2) = \cos(\theta/2) \cos(\theta'/2) + e^{-i(\phi-\phi')} \sin(\theta/2) \sin(\theta'/2). \quad (4.40)$$

The angle γ was defined in (4.24), and the energy E_{qj} is

$$E_{qj} \equiv \sqrt{(j+1/2)^2 + \mu_q^2 - q^2}. \quad (4.41)$$

4.4.1 The D kernel

Let us first determine $D_j^q(\omega)$. Using (4.7), (4.22), and (4.38), we have

$$D_j^q(\omega) = \frac{4\pi}{2j+1} \sum_{j',j''=q}^{\infty} \int d^3x \sqrt{g(x)} F_{0,j}(\theta) F_{q,j'}(\theta) F_{q,j''}(\theta) \frac{e^{-(E_{qj'}+E_{qj''})|\tau|+i\omega\tau}}{4E_{qj}E_{qj'}}. \quad (4.42)$$

Performing the τ integral, we can simplify this expression to

$$D_j^q(\omega) = \frac{8\pi^2}{2j+1} \sum_{j',j''=q}^{\infty} \left[\frac{E_{qj'} + E_{qj''}}{2E_{qj'}E_{qj''}(\omega^2 + (E_{qj'} + E_{qj''})^2)} \right] \mathcal{I}_D(j, j', j''), \quad (4.43)$$

where

$$\mathcal{I}_D(j, j', j'') = \int_0^\pi \sin \theta d\theta F_{0,j}(\theta) F_{q,j'}(\theta) F_{q,j''}(\theta). \quad (4.44)$$

The θ integral can be performed analytically, and we have

$$\mathcal{I}_D(j, j', j'') = \left[\frac{(2j+1)(2j'+1)(2j''+1)}{32\pi^3} \right] \begin{pmatrix} j & j' & j'' \\ 0 & -q & q \end{pmatrix}^2. \quad (4.45)$$

We can check that this result equals (4.28) for $q = 0$ and, for instance, for $j = 0$

$$D_0^0(\omega) = \frac{1}{2\pi} \sum_{j'=0}^{\infty} \frac{1}{\omega^2 + (2j'+1)^2} = \frac{\tanh(\pi\omega/2)}{8\omega}. \quad (4.46)$$

Note that the summation in (4.42) is absolutely convergent.

4.4.2 The K and F kernels

Similarly, for the other kernels we can use (4.7), (4.10), (4.11), and (4.38) to obtain

$$\begin{aligned} F_j^{q,\tau}(\omega) &= F_j^{q,E}(\omega) = K_j^{q,\tau B}(\omega) = K_j^{q,EB}(\omega) = 0, \\ F_j^{q,B}(\omega) &= \frac{16q\pi^2}{(2j+1)\sqrt{j(j+1)}} \sum_{j',j''=q}^{\infty} \left[\frac{E_{qj'} + E_{qj''}}{2E_{qj'}E_{qj''}(\omega^2 + (E_{qj'} + E_{qj''})^2)} \right] \mathcal{I}_F(j, j', j''), \\ K_j^{q,\tau\tau}(\omega) &= \frac{8\pi^2}{2j+1} \sum_{j',j''=q}^{\infty} \left[\frac{-(E_{qj'} + E_{qj''})(\omega^2 + 4E_{qj'}E_{qj''})}{2E_{qj'}E_{qj''}(\omega^2 + (E_{qj'} + E_{qj''})^2)} \right] \mathcal{I}_D(j, j', j'') + \sum_{j'=q}^{\infty} \frac{(2j'+1)}{4\pi E_{qj'}}, \\ K_j^{q,\tau E}(\omega) &= \frac{8\pi^2}{(2j+1)\sqrt{j(j+1)}} \sum_{j',j''=q}^{\infty} \left[\frac{i\omega(E_{qj'} - E_{qj''})}{2E_{qj'}E_{qj''}(\omega^2 + (E_{qj'} + E_{qj''})^2)} \right] \mathcal{I}_\phi(j, j', j''), \\ K_j^{q,EE}(\omega) &= \frac{8\pi^2}{(2j+1)j(j+1)} \sum_{j',j''=q}^{\infty} \left[\frac{(E_{qj'} + E_{qj''})}{2E_{qj'}E_{qj''}(\omega^2 + (E_{qj'} + E_{qj''})^2)} \right] \mathcal{I}_E(j, j', j'') + \sum_{j'=q}^{\infty} \frac{(2j'+1)}{4\pi E_{qj'}}, \\ K_j^{q,BB}(\omega) &= \frac{8\pi^2}{(2j+1)j(j+1)} \sum_{j',j''=q}^{\infty} \left[\frac{(E_{qj'} + E_{qj''})}{2E_{qj'}E_{qj''}(\omega^2 + (E_{qj'} + E_{qj''})^2)} \right] \mathcal{I}_B(j, j', j'') + \sum_{j'=q}^{\infty} \frac{(2j'+1)}{4\pi E_{qj'}}, \end{aligned} \quad (4.47)$$

where

$$\begin{aligned}
\mathcal{I}_F(j, j', j'') &= \int_0^\pi d\theta \sin \theta \tan \frac{\theta}{2} F'_{0,j}(\theta) F_{q,j'}(\theta) F_{q,j''}(\theta), \\
\mathcal{I}_\phi(j, j', j'') &= \left[j''(j'' + 1) - j'(j' + 1) \right] \mathcal{I}_D(j, j', j''), \\
\mathcal{I}_E(j, j', j'') &= - \left[j'(j' + 1) - j''(j'' + 1) \right]^2 \mathcal{I}_D(j, j', j''), \\
\mathcal{I}_B(j, j', j'') &= \int_0^\pi d\theta \sin \theta \left[\frac{4}{\sin \theta} F'_{0,j}(\theta) F'_{q,j'}(\theta) F'_{q,j''}(\theta) - 4q^2 \tan^2(\theta/2) F''_{0,j}(\theta) F_{q,j'}(\theta) F_{q,j''}(\theta) \right].
\end{aligned} \tag{4.48}$$

A detailed derivation of these formulas is contained in Appendix A. The quantity $\mathcal{I}_D(j, j', j'')$ appearing in (4.48) was given explicitly in (4.44). Similar explicit expressions for $\mathcal{I}_F(j, j', j'')$ and $\mathcal{I}_B(j, j', j'')$ are given in Appendix B.

Note that while the expressions for $F_j^{q,\tau}(\omega)$, $F_j^{q,B}(\omega)$, and $K_j^{q,\tau E}(\omega)$ above are absolutely convergent, those for $K_j^{q,\tau\tau}(\omega)$, $K_j^{q,EE}(\omega)$, and $K_j^{q,BB}(\omega)$ are not and require regularization. In order to regularize the latter, it is convenient to first compute the quantities

$$\begin{aligned}
&K_j^{q,\tau\tau}(\omega) - K_0^{0,\tau\tau}(0), \\
&K_j^{q,EE}(\omega) - K_1^{0,EE}(0), \\
&K_j^{q,BB}(\omega) - K_1^{0,BB}(0),
\end{aligned} \tag{4.49}$$

which are free of divergences, and then add back the appropriately regularized values for $K_0^{0,\tau\tau}(0)$, $K_1^{0,EE}(0)$, and $K_1^{0,BB}(0)$. As argued in the previous subsection, gauge invariance implies $K_0^{0,\tau\tau}(\omega) = K_1^{0,EE}(0) = 0$, but does not immediately determine $K_1^{0,BB}(0)$ denoted by C_0 in (4.33). We can return now to that issue. Let us first examine $K_0^{0,\tau\tau}(\omega) = 0$. From (4.47), we have

$$K_0^{0,\tau\tau}(0) = \sum_{j'=0}^{\infty} \left[\frac{-(2j'+1)}{4\pi E_{0j'}} \right] + \sum_{j'=0}^{\infty} \frac{(2j'+1)}{4\pi E_{0j'}} = 0. \tag{4.50}$$

(Each sum is divergent individually, but the combined summation is convergent.) Next, we can examine $K_1^{0,EE}(0)$. Using (4.47) and doing a bit of algebra, we have

$$K_1^{0,EE}(0) = \frac{1}{32\pi} \sum_{|j'-j''|=1} \left[-2 - \frac{j''+1/2}{j'+1/2} - \frac{j'+1/2}{j''+1/2} \right] + \sum_{j'=0}^{\infty} \frac{(2j'+1)}{4\pi(j'+1/2)}. \tag{4.51}$$

By using the symmetry between the summation in j' and j'' , this expression can be written further as

$$K_1^{0,EE}(0) = \frac{1}{16\pi} \sum_{|j'-j''|=1} \left[-1 - \frac{j'' + 1/2}{j' + 1/2} \right] + \sum_{j'=0}^{\infty} \frac{(2j' + 1)}{4\pi(j' + 1/2)}. \quad (4.52)$$

Summing over j'' we can write this expression as

$$K_1^{0,EE}(0) = -\frac{1}{4\pi} \sum_{j'=0}^{\infty} \frac{(j' + 1/2)}{j' + 1/2} + \sum_{j'=0}^{\infty} \frac{(2j' + 1)}{4\pi(j' + 1/2)}. \quad (4.53)$$

Both of the individual sums in this expression as well as the combined summation are divergent, but gauge invariance dictates that $K_1^{0,EE}(0) = 0$. This result should be thought of as a prescription. It can also be justified in zeta-function regularization, in which (4.53) gives $K_1^{0,EE}(0) = \zeta(0, 1/2)/(4\pi) = 0$.

Next, from (4.47), we can also write an expression for $K_1^{0,BB}(0)$:

$$K_1^{0,BB}(0) = \frac{1}{8\pi} \sum_{j'=0}^{\infty} \left[\frac{-j'(j' + 1)(2j' + 1)}{(j' + 1/2)^3} \right] + \sum_{j'=0}^{\infty} \frac{(2j' + 1)}{4\pi(j' + 1/2)}. \quad (4.54)$$

Both terms are again divergent, but, using (4.53), we can calculate

$$K_1^{0,BB}(0) - K_1^{0,EE}(0) = \frac{1}{4\pi} \sum_{j'=0}^{\infty} \frac{1}{(2j' + 1)^2} = \frac{\pi}{32} = \frac{1}{2} D_0^0(0). \quad (4.55)$$

Since $K_1^{0,EE}(0) = 0$, this equation proves that $C_0 = 0$ in (4.33).

We can now provide alternate, appropriately regularized formulas for $K_j^{q,\tau\tau}(\omega)$, $K_j^{q,EE}(\omega)$,

and $K_j^{q,BB}(\omega)$ that are manifestly convergent. They are:

$$\begin{aligned}
K_j^{q,\tau\tau}(\omega) &= \sum_{j'=q}^{\infty} \left[\frac{8\pi^2}{2j+1} \sum_{j''=q}^{\infty} \frac{-(E_{qj'} + E_{qj''})(\omega^2 + 4E_{qj'}E_{qj''})}{2E_{qj'}E_{qj''}(\omega^2 + (E_{qj'} + E_{qj''})^2)} \mathcal{I}_D(j, j', j'') \right. \\
&\quad \left. + \frac{(2j'+1)}{4\pi E_{qj'}} \right], \\
K_j^{q,EE}(\omega) &= \sum_{j'=q}^{\infty} \left[\frac{8\pi^2}{(2j+1)j(j+1)} \sum_{j''=q}^{\infty} \frac{(E_{qj'} + E_{qj''})}{2E_{qj'}E_{qj''}(\omega^2 + (E_{qj'} + E_{qj''})^2)} \mathcal{I}_E(j, j', j'') \right. \\
&\quad \left. + \frac{(2j'+1)}{8\pi E_{qj'}} \right] + C_q, \\
K_j^{q,BB}(\omega) &= \sum_{j'=q}^{\infty} \left[\frac{8\pi^2}{(2j+1)j(j+1)} \sum_{j''=q}^{\infty} \frac{(E_{qj'} + E_{qj''})}{2E_{qj'}E_{qj''}(\omega^2 + (E_{qj'} + E_{qj''})^2)} \mathcal{I}_B(j, j', j'') \right. \\
&\quad \left. + \frac{(2j'+1)}{8\pi E_{qj'}} \right] + C_q,
\end{aligned} \tag{4.56}$$

where

$$C_q \equiv \sum_{j'=q}^{\infty} \frac{2j'+1}{8\pi E_{qj'}} - \sum_{j'=0}^{\infty} \frac{2j'+1}{8\pi E_{0j'}} = \frac{1}{4\pi} \left[\sum_{j'=q}^{\infty} \left(\frac{j'+1/2}{\sqrt{(j'+1/2)^2 + \mu_q^2 - q^2}} - 1 \right) - q \right]. \tag{4.57}$$

The expression for $K_j^{q,\tau\tau}(\omega)$ was obtained by simply combining the two summations in the expression in (4.47). The expressions for $K_j^{q,EE}(\omega)$ and $K_j^{q,BB}(\omega)$ were obtained by subtracting $K_1^{q,EE}(0) = 0$ from the expressions in (4.47). In (4.57) we discover the saddle point equation for μ_q^2 (3.8) (obtained after tuning the coupling \mathfrak{g} to the critical value \mathfrak{g}_c), thus $C_q = 0$.

One can check from (4.47) and (4.48) that the gauge-invariance relations (4.15) are obeyed. Such a check is most simply performed by matching the residues of the functions of ω at their poles at $\omega = \pm i(E_{qj'} + E_{qj''})$ for each j' and j'' , after symmetrization between j' and j'' —see Appendix A.2.

Now that we have expressions (4.56) we can begin evaluating the order $1/N_b$ corrections to the free energy, which is the subject of the next subsection.

4.5 Numerical Results

With the regularized formulas for the kernels in hand, we are almost ready to calculate the subleading correction $\delta\mathcal{F}_q$ to the free energy using (4.19). Let us write this expression as

$$\delta\mathcal{F}_q = \frac{1}{2} \int \frac{d\omega}{2\pi} \sum_{j=0}^{\infty} (2j+1) L_j^q(\omega), \quad (4.58)$$

where $L_j^q(\omega)$ can be read off from (4.19). (See also (C.47)–(C.48).) As shown in Appendix C, at large ω and j the integrand in this expression behaves as

$$L_j^q(\omega) = \frac{8\mu_q^2}{\omega^2 + (j+1/2)^2} + \dots, \quad (4.59)$$

thus rendering the integral (4.58) linearly divergent. There are several ways of understanding how to regularize this divergence. One way is to use zeta-function regularization to write

$$\frac{1}{2} \int \frac{d\omega}{2\pi} \sum_{j=0}^{\infty} (2j+1) \frac{8\mu_q^2}{\omega^2 + (j+1/2)^2} = \frac{1}{2} \sum_{j=0}^{\infty} (2j+1) \frac{8\mu_q^2}{2j+1} = 4\mu_q^2 \zeta(0, 1/2) = 0. \quad (4.60)$$

Then one can subtract (4.60) from (4.58), and evaluate

$$\delta\mathcal{F}_q = \frac{1}{2} \int \frac{d\omega}{2\pi} \sum_{j=0}^{\infty} (2j+1) \left[L_j^q(\omega) - \frac{8\mu_q^2}{\omega^2 + (j+1/2)^2} \right] \quad (4.61)$$

instead of (4.58). This expression is no longer linearly divergent.

Another way of understanding the subtraction in (4.61) is that the critical coupling \mathfrak{g}_c , which was obtained in (3.6) at leading order in N_b , receives $1/N_b$ corrections. A similar phenomenon was encountered in [14] when computing the thermal free energy at subleading order in $1/N_b$. Just as in [14], it can be argued that

$$\frac{4\pi}{\mathfrak{g}_c} = \int \frac{d\omega}{2\pi} \sum_{j=0}^{\infty} (2j+1) \frac{1}{\omega^2 + (j+1/2)^2} \left[1 + \frac{4}{N_b} \right] + O(1/N_b^2). \quad (4.62)$$

The $1/N_b$ term in this expression contributes to $\delta\mathcal{F}_q^\infty$ through the last term in (2.3) precisely as the subtraction implemented in (4.61). This expression will be derived rigorously in [34].

Even after the linear divergence in $\delta\mathcal{F}_q^\infty$ has been taken care of, this quantity is still potentially logarithmically divergent. This logarithmic divergence cancels when using a regularization prescription consistent with conformal symmetry. In practice, we evaluate the

integral in (4.61) with a symmetric cutoff:

$$(j + 1/2)^2 + \omega^2 < \Lambda^2. \quad (4.63)$$

This can be thought of as preserving rotational invariance on \mathbb{R}^3 , as the high energy modes are insensitive to the curvature of the sphere. Given the kernels, (4.56), and the regularization described above, we are able to evaluate $\delta\mathcal{F}_q$ numerically. To obtain good precision, we first evaluate (4.61) numerically in a region $(j + 1/2)^2 + \omega^2 < (\Lambda')^2$; then in the region $(\Lambda')^2 < (j + 1/2)^2 + \omega^2 < \Lambda^2$ we replace $L_j^q(\omega)$ in (4.61) with the asymptotic expansion derived in Appendix C (accurate up to terms of order $O(1/[(j + \frac{1}{2})^2 + \omega^2]^{7/2})$) and evaluate the integral analytically as $\Lambda \rightarrow \infty$. We notice that the result converges very rapidly as we increase Λ' . (In practice, $\Lambda' = 10$ is already sufficiently large.) Our results for $\delta\mathcal{F}_q$ when $q = 1/2$ and $q = 1$ are given in Table 2. The results at $q = 1/2$ were compared with quantum Monte Carlo studies on $SU(N_b)$ antiferromagnets in Fig. 1.


q	\mathcal{F}_q^∞	$\delta\mathcal{F}_q$
0	0	0
1/2	0.1245922	0.05992
1	0.3110952	-0.2338

Table 2: The coefficients in the large N_b expansion (2.2) of the ground state energy in the presence of $2q$ units of magnetic flux through S^2 .


5 Stability

5.1 Numerical results


In the previous sections we studied properties of monopole operators by looking at fluctuations around a rotationally-invariant saddle-point configuration. In the case when this saddle point is a minimum, that is when the quadratic effective action is positive definite, calculating the Euclidean action around this configuration gives the dimension of the corresponding monopole operator. However, when this saddle is unstable, we cannot identify it with any primary operator in the low energy theory. Although the saddle given by (2.7) exists for any $q \in \mathbb{Z}/2$, it need not be a stable saddle. We find that the instability occurs for $q \geq 3/2$, when λ_-^q , the smallest eigenvalue of (4.15), becomes negative as shown in Figure 2.



(a) The smallest eigenvalue, λ_-^q , plotted as a function of ω for various values of the monopole charge q and angular momentum $j = 1$. Note that λ_-^q is only positive for $q < 3/2$.



(b) The eigenvalue λ_-^q for different j . The eigenvalue is negative only for small j and ω .



(c) The other two eigenvalues, λ_E^q and λ_+^q , remain positive even for small j and ω .

Figure 2: Plots of the eigenvalues λ_E^q and λ_{\pm}^q as functions of ω for various values of the monopole charge q and angular momentum j .

Similar instabilities occur when studying quantum fluctuations about classical saddles in other theories [33]. What is particularly surprising about the present case is that these classical backgrounds carry charge under the topological symmetry, $U(1)_{\text{top}}$, and are natural candidates for being the dominant saddles in their respective topological sectors. Despite this, they possess negative modes for $q \geq 3/2$. As there must exist a lowest-dimension primary operator in the low energy theory for any value of the $U(1)_{\text{top}}$ charge, and the only assumption that went into our saddle point configuration was spherical symmetry, the operators carrying charge must not come from spherically symmetric saddles. This is further motivated by the fact that the instability is mediated by modes of the scalar and gauge field carrying angular momentum, $j \geq 1$.⁹

5.2 Analytic understanding in the large q limit

Whereas the analytic computation of $\mathbf{M}_j^q(\omega)$ is a hopeless endeavor for finite q , we found that the $q \rightarrow \infty$ limit is tractable. The reason for the simplification is that this is essentially a flat space limit: reintroducing the radius R of S^2 we have a strong magnetic field $B = q/R^2$ on the sphere at large q . ϕ quanta move on Landau levels, which are localized on $1/\sqrt{B} = R/\sqrt{q}$ distances, hence they don't feel the effect of the curvature of the sphere. To leading order in $1/q$, the problem becomes the analysis of the $\mathbb{C}\mathbb{P}^{N_b-1}$ in a constant magnetic field in flat space.

First, we want to calculate \mathcal{F}_q^∞ . We will be more cavalier about divergences than in the rest of the paper, and write, following (3.7),

$$\begin{aligned} \mathcal{F}_q^\infty &= \sum_{j=q}^{\infty} (2j+1) [(j+1/2)^2 + \mu_q^2 - q^2]^{1/2} \\ &= \sum_{n=0}^{\infty} (2q+2n+1) [2q(n+1/2) + \mu_q^2 + (n+1/2)^2]^{1/2}, \end{aligned} \tag{5.1}$$

where we introduced $n \equiv j - q$, and assumed zeta-function regularization as implicit. The resulting saddle point equation for μ_q^2 is (3.8):

$$\sum_{n=0}^{\infty} \frac{2q+2n+1}{\sqrt{2q(n+1/2) + \mu_q^2 + (n+1/2)^2}} = 0. \tag{5.2}$$

⁹Spherically asymmetric saddles are not entirely unfamiliar, for instance [46] construct such asymmetric saddles in various supersymmetric Chern-Simons matter theories including $U(1)_k$ Chern-Simons theory.

This equation has a solution only provided that we scale μ_q^2 correctly with q , namely

$$\mu_q^2 = 2q\chi_0 + \chi_1 + O\left(\frac{1}{q}\right). \quad (5.3)$$

Plugging this Ansatz into (5.2) and only keeping the leading terms, we obtain

$$0 = \sum_{n=0}^{\infty} \frac{1}{\sqrt{(n+1/2) + \chi_0}} = \zeta\left(\frac{1}{2}, \frac{1}{2} + \chi_0\right). \quad (5.4)$$

Note that in obtaining this equation we assumed that $n \ll q$ even though we are summing over all positive n . This assumption is justified because the contribution of $n \gtrsim q$ is higher order in $1/q$. The constant χ_0 can therefore be obtained as the root of the transcendental equation (5.4). Going to one higher order we can determine χ_1 :

$$\chi_1 = -\chi_0^2 + \frac{3\zeta\left(-\frac{1}{2}, \frac{1}{2} + \chi_0\right)}{\zeta\left(\frac{3}{2}, \frac{1}{2} + \chi_0\right)}, \quad (5.5)$$

which gives the large q expansion for μ_q^2 :

$$\mu_q^2 \approx -0.39456q - 0.00456 + O\left(\frac{1}{q}\right). \quad (5.6)$$

As seen from Figure 3, (5.6) is in excellent agreement with the values of μ_q^2 obtained from solving (3.8) at large q .¹⁰




Figure 3: A comparison between the values of μ_q^2 found by solving (3.8) (blue points) and the large- q analytical approximation (5.6) (solid black line).

¹⁰In [23] it was noticed that $\mu_q^2 \approx -2q/5$ at large q .

Plugging back into (5.1), we have that at leading order in q

$$\begin{aligned}\mathcal{F}_q^\infty &= 2q \sum_{n=0}^{\infty} \sqrt{2q(n+1/2) + 2q\chi_0} = (2q)^{3/2} \zeta\left(-\frac{1}{2}, \frac{1}{2} + \chi_0\right) \\ &= 0.26408 q^{3/2}.\end{aligned}\tag{5.7}$$

Quite nicely, this equation can be understood in flat space terms: the Landau levels of a massive scalar field are given by

$$E_n = \sqrt{2B(n+1/2) + m^2} = \frac{1}{R} \sqrt{2q(n+1/2) + 2q\chi_0},\tag{5.8}$$

and have degeneracy $\mathcal{N} = \frac{B \text{Vol}(S^2)}{2\pi} = 2q$, giving exactly (5.1) for the free energy (if we set $R = 1$). Note that the $q^{3/2}$ scaling of (5.7) follows from flat space dimensional analysis: the free energy density is an intensive quantity of mass dimension 3, hence it has to be independent of R , and we get $\mathcal{F} \sim B^{3/2}$.

To get analytic insight into how the instability of the monopole background is connected to the flat space limit we examine $\mathbf{M}_j^q(\omega)$ at $\omega = 0$ for $j \ll q$. This corresponds in the flat space limit to taking the momentum $p \ll BR$. At $\omega = 0$, we only have to determine D_j^q , $F_j^{q,B}$, $K_j^{q,BB}$, and $K_j^{q,\tau\tau}$, as the rest of the matrix elements vanish—see (4.15). We can obtain a closed-form formula for these kernels by taking the explicit expressions for them, and expanding for large q . This is quite a tedious task, especially for $K_j^{q,BB}$, where we have to expand (B.18) for fixed j, j', j'' and large q . The resulting expressions can be summed over j', j'' analytically using zeta-function regularization. The results are given by the simple expression¹¹

$$\mathbf{M}_j^q(0) = \frac{\zeta\left(\frac{3}{2}, \frac{1}{2} + \chi_0\right)}{8\pi\sqrt{2q}} \left(\begin{array}{c|ccc} \frac{1}{2} & \sqrt{j(j+1)}\chi_0 & 0 & 0 \\ \hline \sqrt{j(j+1)}\chi_0 & 2j(j+1)\chi_1 & 0 & 0 \\ 0 & 0 & 4j(j+1)(\chi_0^2 + \chi_1) & 0 \\ 0 & 0 & 0 & 0 \end{array} \right).\tag{5.9}$$

One curious subtlety is that while the individual terms in $K_j^{q,BB}$, $K_j^{q,\tau\tau}$ are $O(\sqrt{q})$, this $O(\sqrt{q})$ contribution vanishes upon summation over j', j'' . The subleading terms give the result in (5.9). This implies that we have to know the saddle point value of μ_q^2 to first subleading order (5.3) and χ_1 makes appearance in the final result.

¹¹For $j = 0$ the matrix is 2×2 , and the only nonzero element is $D_0^q(0)$, as in (4.17).

We obtain the flat space kernel by the replacement $j(j+1) \rightarrow p^2$ in (5.9). Our expression is valid for $p \ll BR$. It would be an interesting exercise to obtain the full quadratic effective action of the \mathbb{CP}^{N_b-1} model in flat space in a constant magnetic field. Such an analysis could also connect the instability to ‘magnetic catalysis’ [47].

The matrix in (5.9) has one zero eigenvalue corresponding to the pure gauge mode. The nonzero eigenvalues for $j = 1$ are (4.16):

$$\lambda_-^q \approx -\frac{0.03209}{\sqrt{q}}, \quad \lambda_+^q \approx \frac{0.142589}{\sqrt{q}}, \quad \lambda_E^q \approx \frac{0.063044}{\sqrt{q}}. \quad (5.10)$$

Note that λ_-^q is negative, giving an analytic understanding of the instability found numerically in the previous subsection. We can check that (5.10) agrees with the numerical results—see Figure 4.




Figure 4: The numerical results for the three eigenvalues, λ_E^q , λ_+^q , and λ_-^q are plotted against the analytic large q value in black.

6 Conclusions

We have presented here the leading correction to the large N_b result for the scaling dimension of the monopole operator in the \mathbb{CP}^{N_b-1} CFT. This correction was obtained by computing the Gaussian fluctuation determinant of the $U(1)$ gauge field A_μ , and the Lagrange multiplier λ , on $S^2 \times \mathbb{R}$. Computation of higher order terms in the $1/N_b$ expansion appears to be possible by the present methods, but will involve considerable effort.

Our computation now opens the possibility of quantitatively testing the most subtle and

novel aspects of the theory of deconfined criticality [8, 9] in two-dimensional lattice antiferromagnets. An important feature of this theory is the connection between the monopole operator and the VBS operator of the antiferromagnet [21–23]. This connection allows a Monte Carlo computation of the monopole scaling dimension by measuring correlators of the VBS order in lattice models. We compared our present result with the Monte Carlo studies in Fig. 1, and found a promising agreement for the largest value, $N_b = 10$, and for the trends as $N_b \rightarrow \infty$. We hope that future Monte Carlo studies will increase the precision of the exponents, and allow for a conclusive comparison between theory and numerics.

We conclude by noting that our present study is among the most complex theoretical computations of critical exponents which have been compared to Monte Carlo simulations. Our calculation assumed conformal invariance to realize a framework in which the exponents could be determined, and it did not reduce to identifying poles in a Feynman graph expansion [48]. It would be of great interest to apply the recent progress in bootstrap methods [49, 50] to also determine such exponents.

Acknowledgements

We thank Ribhu Kaul for valuable discussions and for providing the numerical results in Fig. 1. We also thank Max Metlitski for useful discussions. The work of ED was supported by the NSF under grant PHY-0756174. The work of MM was supported by the Princeton Center for Theoretical Science. SSP was supported in part by the US NSF under grant No. PHY-1418069. The work of SS was supported by the NSF under grant DMR-1360789, the Templeton foundation, and MURI grant W911NF-14-1-0003 from ARO. Research at Perimeter Institute is supported by the Government of Canada through Industry Canada and by the Province of Ontario through the Ministry of Research and Innovation.

A Derivation of integration kernels

A.1 Evaluation of Fourier transforms

In this Appendix we explain how to obtain the explicit formulas appearing in (4.47). When dealing with vectors on S^2 it is convenient to use the frame e_a^i , with $i = \theta, \phi$ and $a = 1, 2$, defined by

$$e_1^i = (1, 0), \quad e_2^i = \left(0, \frac{1}{\sin \theta}\right). \quad (\text{A.1})$$

One can then convert the coordinate index of a quantity v_i to a frame index by writing

$$v_a = e_a^i v_i . \quad (\text{A.2})$$

The frame index can then be raised and lowered with δ^{ab} and δ_{ab} , respectively; in other words, it makes no difference whether it is upper or lower.

Our starting point are equations (4.9)–(4.11). In the main text, we explained how (4.9) can be inverted to obtain (4.20). The analogous formulas for the other kernels are

$$\begin{aligned} K_j^{q,\tau\tau}(\omega) &= \frac{4\pi}{2j+1} \int d^3x \lim_{x' \rightarrow 0} \sum_{m=-j}^j Y_{jm}^*(\theta, \phi) K_{\tau\tau'}^q(x, x') Y_{jm}(\theta', \phi') e^{i\omega\tau} , \\ K_j^{q,\tau E}(\omega) &= \frac{4\pi}{2j+1} \int d^3x \lim_{x' \rightarrow 0} \sum_{m=-j}^j Y_{jm}^*(\theta, \phi) K_{\tau a'}^q(x, x') \mathcal{X}_{jm}^{a'}(\theta', \phi') e^{i\omega\tau} , \\ K_j^{q,EE}(\omega) &= \frac{4\pi}{2j+1} \int d^3x \lim_{x' \rightarrow 0} \sum_{m=-j}^j \mathcal{X}_{jm}^{a*}(\theta, \phi) K_{aa'}^q(x, x') \mathcal{X}_{jm}^{a'}(\theta', \phi') e^{i\omega\tau} , \\ K_j^{q,BB}(\omega) &= \frac{4\pi}{2j+1} \int d^3x \lim_{x' \rightarrow 0} \sum_{m=-j}^j \mathcal{Y}_{jm}^{a*}(\theta, \phi) K_{aa'}^q(x, x') \mathcal{Y}_{jm}^{a'}(\theta', \phi') e^{i\omega\tau} , \\ F_j^{q,B}(\omega) &= \frac{4\pi}{2j+1} \int d^3x \lim_{x' \rightarrow 0} \sum_{m=-j}^j Y_{jm}^*(\theta, \phi) F_{a'}^q(x, x') \mathcal{Y}_{jm}^{a'}(\theta', \phi') e^{i\omega\tau} , \\ D_j^q(\omega) &= \frac{4\pi}{2j+1} \int d^3x \lim_{x' \rightarrow 0} \sum_{m=-j}^j Y_{jm}^*(\theta, \phi) D^q(x, x') Y_{jm}(\theta', \phi') e^{i\omega\tau} . \end{aligned} \quad (\text{A.3})$$

Using the spectral decomposition (4.38), the position-space kernels appearing in this expres-

sion can be written as

$$\begin{aligned}
D^q(x, x') &= \sum_{j', j''} F_{q, j'}(\gamma) F_{q, j''}(\gamma) \frac{e^{-(E_{qj'} + E_{qj''})|\tau - \tau'|}}{4E_{qj'} E_{qj''}}, \\
K_{\tau\tau'}^q(x, x') &= \sum_{j', j''} F_{q, j'}(\gamma) F_{q, j''}(\gamma) \frac{e^{-(E_{qj'} + E_{qj''})|\tau - \tau'|}}{4E_{qj'} E_{qj''}} [(E_{qj'} - E_{qj''})^2 - 2(E_{qj'} + E_{qj''})\delta(\tau)] \\
&\quad + \sum_{j'} \frac{2j' + 1}{4\pi E_{qj'}} \delta(x - x'), \\
K_{\tau a'}^q(x, x') &= \sum_{j', j''} \frac{e^{-(E_{qj'} + E_{qj''})|\tau - \tau'|}}{4E_{qj'} E_{qj''}} (E_{qj'} - E_{qj''}) \operatorname{sgn}(\tau - \tau') \\
&\quad \times \left([D_{a'} e^{-2iq\Theta} F_{q, j'}(\gamma)] [e^{-2iq\Theta} F_{q, j''}(\gamma)]^* - [e^{-2iq\Theta} F_{q, j'}(\gamma)] [D_{a'} e^{-2iq\Theta} F_{q, j''}(\gamma)]^* \right), \\
K_{a a'}^q(x, x') &= \sum_{j'} \frac{2j' + 1}{4\pi E_{qj'}} \delta^{a a'} \delta(x - x') + \sum_{j', j''} \frac{e^{-(E_{qj'} + E_{qj''})|\tau - \tau'|}}{4E_{qj'} E_{qj''}} \\
&\quad \times \left([D_a e^{-2iq\Theta} F_{q, j'}(\gamma)] [D_{a'} e^{-2iq\Theta} F_{q, j''}(\gamma)]^* + [D_{a'} e^{-2iq\Theta} F_{q, j'}(\gamma)] [D_a e^{-2iq\Theta} F_{q, j''}(\gamma)]^* \right. \\
&\quad \left. - [e^{-2iq\Theta} F_{q, j'}(\gamma)] [D_a D_{a'} e^{-2iq\Theta} F_{q, j''}(\gamma)]^* - [D_a D_{a'} e^{-2iq\Theta} F_{q, j'}(\gamma)] [e^{-2iq\Theta} F_{q, j''}(\gamma)]^* \right), \\
F_{a'}^q(x, x') &= \sum_{j', j''} \frac{e^{-(E_{qj'} + E_{qj''})|\tau - \tau'|}}{4E_{qj'} E_{qj''}} \\
&\quad \times \left([e^{-2iq\Theta} F_{q, j'}(\gamma)] [D_{a'} e^{-2iq\Theta} F_{q, j''}(\gamma)]^* - [D_{a'} e^{-2iq\Theta} F_{q, j'}(\gamma)] [e^{-2iq\Theta} F_{q, j''}(\gamma)]^* \right), \tag{A.4}
\end{aligned}$$

where

$$D_a \equiv \partial_a - i\mathcal{A}_a^q(x), \quad D_{a'} \equiv \partial_{a'} + i\mathcal{A}_{a'}^q(x'). \tag{A.5}$$

In taking the $x' \rightarrow 0$ limit in (A.3), it is convenient to use the addition formula for the spherical harmonics

$$\sum_{m=-j}^j Y_{jm}^*(\theta, \phi) Y_{jm}(\theta', \phi') = F_{0j}(\gamma), \quad F_{0j}(\gamma) \equiv \frac{2j+1}{4\pi} P_j(\cos \theta), \tag{A.6}$$

where γ is the relative angle between the points (θ, ϕ) and (θ', ϕ') on S^2 . Taking derivatives

of this formula and using the definition of the vector harmonics in (4.8), we have

$$\begin{aligned}
\lim_{x' \rightarrow 0} \sum_{m=-j}^j Y_{jm}^*(\theta, \phi) Y_{jm}(\theta', \phi') &= F_{0j}(\theta), \\
\lim_{x' \rightarrow 0} \sum_{m=-j}^j Y_{jm}^*(\theta, \phi) \mathcal{X}_{jm}^{a'}(\theta', \phi') &= \frac{1}{\sqrt{j(j+1)}} \begin{pmatrix} -F'_{0j}(\theta) & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} \cos \phi & \sin \phi \\ -\sin \phi & \cos \phi \end{pmatrix}, \\
\lim_{x' \rightarrow 0} \sum_{m=-j}^j Y_{jm}^*(\theta, \phi) \mathcal{Y}_{jm}^{a'}(\theta', \phi') &= \frac{1}{\sqrt{j(j+1)}} \begin{pmatrix} 0 & F'_{0j}(\theta) \\ 0 & 0 \end{pmatrix} \begin{pmatrix} \cos \phi & \sin \phi \\ -\sin \phi & \cos \phi \end{pmatrix}, \\
\lim_{x' \rightarrow 0} \sum_{m=-j}^j \mathcal{X}_{jm}^{a*}(\theta, \phi) Y_{jm}(\theta', \phi') &= \frac{1}{\sqrt{j(j+1)}} \begin{pmatrix} F'_{0j}(\theta) \\ 0 \end{pmatrix}, \\
\lim_{x' \rightarrow 0} \sum_{m=-j}^j \mathcal{Y}_{jm}^{a*}(\theta, \phi) Y_{jm}(\theta', \phi') &= \frac{1}{\sqrt{j(j+1)}} \begin{pmatrix} 0 \\ -F'_{0j}(\theta) \end{pmatrix}, \\
\lim_{x' \rightarrow 0} \sum_{m=-j}^j \mathcal{X}_{jm}^{a*}(\theta, \phi) \mathcal{X}_{jm}^{a'}(\theta', \phi') &= \frac{1}{j(j+1)} \begin{pmatrix} -F''_{0j}(\theta) & 0 \\ 0 & -\frac{F'_{0j}(\theta)}{\sin \theta} \end{pmatrix} \begin{pmatrix} \cos \phi & \sin \phi \\ -\sin \phi & \cos \phi \end{pmatrix}, \\
\lim_{x' \rightarrow 0} \sum_{m=-j}^j \mathcal{Y}_{jm}^{a*}(\theta, \phi) \mathcal{Y}_{jm}^{a'}(\theta', \phi') &= \frac{1}{j(j+1)} \begin{pmatrix} -\frac{F'_{0j}(\theta)}{\sin \theta} & 0 \\ 0 & -F''_{0j}(\theta) \end{pmatrix} \begin{pmatrix} \cos \phi & \sin \phi \\ -\sin \phi & \cos \phi \end{pmatrix}.
\end{aligned} \tag{A.7}$$

We also have

$$\begin{aligned}
\lim_{x' \rightarrow 0} \left[e^{-2iq\Theta(x, x')} F_{qj}(\gamma) \right] &= F_{qj}(\theta), \\
\lim_{x' \rightarrow 0} D_a \left[e^{-2iq\Theta(x, x')} F_{qj}(\gamma) \right] &= \begin{pmatrix} F'_{qj}(\theta) \\ -iq \tan \frac{\theta}{2} F_{qj}(\theta) \end{pmatrix}, \\
\lim_{x' \rightarrow 0} D_{a'} \left[e^{-2iq\Theta(x, x')} F_{qj}(\gamma) \right] &= \begin{pmatrix} -F'_{qj}(\theta) & -iq \tan \frac{\theta}{2} F_{qj}(\theta) \\ 0 & 0 \end{pmatrix} \begin{pmatrix} \cos \phi & \sin \phi \\ -\sin \phi & \cos \phi \end{pmatrix}, \\
\lim_{x' \rightarrow 0} D_a D_{a'} \left[e^{-2iq\Theta(x, x')} F_{qj}(\gamma) \right] &= \begin{pmatrix} -F''_{qj}(\theta) & \frac{-iq(F_{qj}(\theta) + \sin \theta F'_{qj}(\theta))}{1 + \cos \theta} \\ \frac{iq(F_{qj}(\theta) - \sin \theta F'_{qj}(\theta))}{1 + \cos \theta} & -\frac{F'_{qj}(\theta)}{\sin \theta} + q^2 \tan^2 \frac{\theta}{2} F_{qj}(\theta) \end{pmatrix} \\
&\quad \times \begin{pmatrix} \cos \phi & \sin \phi \\ -\sin \phi & \cos \phi \end{pmatrix}.
\end{aligned} \tag{A.8}$$

Plugging in (A.7) and (A.8) into (A.3) and performing the integrals over ω and ϕ , we

obtain (4.43) and (4.47), with

$$\begin{aligned}
\mathcal{I}_D(j, j', j'') &\equiv \int_0^\pi d\theta \sin \theta F_{0j}(\theta) F_{qj'}(\theta) F_{qj''}(\theta), \\
\mathcal{I}_F(j, j', j'') &\equiv \int_0^\pi d\theta \sin \theta \tan \frac{\theta}{2} F'_{0j}(\theta) F_{qj'}(\theta) F_{qj''}(\theta), \\
\mathcal{I}_\phi(j, j', j'') &\equiv \int_0^\pi d\theta \sin \theta F'_{0j}(\theta) (F'_{qj'}(\theta) F_{qj''}(\theta) - F_{qj'}(\theta) F'_{qj''}(\theta)), \\
\mathcal{I}_E(j, j', j'') &\equiv \int_0^\pi d\theta \sin \theta A_E(\theta), \\
\mathcal{I}_B(j, j', j'') &\equiv \int_0^\pi d\theta \sin \theta A_B(\theta),
\end{aligned} \tag{A.9}$$

where

$$\begin{aligned}
A_E &= F''_{0j} (2F'_{qj'} F'_{qj''} - F_{qj'} F''_{qj''} - F''_{qj'} F_{qj''}) - \frac{F'_{0j}}{\sin^2 \theta} \frac{\partial}{\partial \theta} [F_{qj'} F_{qj''}] - \frac{4q^2}{\sin \theta} \tan^2 \frac{\theta}{2} F'_{0j} F_{qj'} F_{qj''}, \\
A_B &= \frac{1}{\sin \theta} \left[-F''_{0j} \frac{\partial}{\partial \theta} [F_{qj'} F_{qj''}] + F'_{0j} (2F'_{qj'} F'_{qj''} - F_{qj'} F''_{qj''} - F''_{qj'} F_{qj''}) \right] \\
&\quad - 4q^2 \tan^2 \frac{\theta}{2} F''_{0j} F_{qj'} F_{qj''}.
\end{aligned} \tag{A.10}$$

To simplify these expressions, we can use the fact that the monopole spherical harmonics $Y_{q,jm}(\theta, \phi)$ are eigenfunctions of the gauge-covariant laplacian on S^2 with eigenvalue $j(j+1) - q^2$:

$$(\nabla^\mu - i\mathcal{A}^{q\mu})(\nabla_\mu - i\mathcal{A}_\mu) Y_{q,jm}(\theta, \phi) = [j(j+1) - q^2] Y_{q,jm}(\theta, \phi). \tag{A.11}$$

Since $F_{qj}(\theta) = \sqrt{\frac{2j+1}{4\pi}} Y_{q,j(-q)}(\theta, 0)$, we also have

$$\left[\frac{1}{\sin \theta} \partial_\theta (\sin \theta \partial_\theta) - q^2 \tan^2 \frac{\theta}{2} + j(j+1) - q^2 \right] F_{qj}(\theta) = 0. \tag{A.12}$$

Eq. (A.12) implies that after integration by parts in the formula for \mathcal{I}_ϕ above, we have

$$\mathcal{I}_\phi(j, j', j'') = \int d\theta \sin \theta F_{0,j}(\theta) F_{q,j'}(\theta) F_{q,j''}(\theta) [j''(j''+1) - j'(j'+1)], \tag{A.13}$$

thus obtaining the simplified form for \mathcal{I}_ϕ presented in (4.48).

One can simplify \mathcal{I}_E as follows. From (A.12), it follows that

$$\frac{1}{\sin \theta} \partial_\theta (\sin \theta F''_{qj'}) = F'_{qj'} \left[-j'(j' + 1) + \frac{1}{\sin^2 \theta} + \frac{2q^2}{\sin \theta} \tan \frac{\theta}{2} \right] + \frac{2q^2}{\sin \theta} \tan^2 \frac{\theta}{2} F_{qj'} \quad (\text{A.14})$$

and similarly for $F_{qj''}$. After integration by parts in the second and third terms in the first paranthesis in (A.10) one obtains

$$\begin{aligned} A_E &= 2F''_{0j} F'_{qj'} F'_{qj''} + F'_{0j} (F''_{qj'} F'_{qj''} + F'_{qj'} F''_{qj''}) + \frac{q^2}{\cos^2 \frac{\theta}{2}} F'_{0j} (F_{qj'} F_{qj''})' \\ &\quad - F'_{0j} [j'(j' + 1) F'_{qj'} F_{qj''} + j''(j'' + 1) F_{qj'} F'_{qj''}] . \end{aligned} \quad (\text{A.15})$$

Integrating by parts in the last term and using (A.12), one has

$$\begin{aligned} A_E &= 2F''_{0j} F'_{qj'} F'_{qj''} + F'_{0j} (F''_{qj'} F'_{qj''} + F'_{qj'} F''_{qj''}) + \frac{q^2}{\cos^2 \frac{\theta}{2}} F'_{0j} (F_{qj'} F_{qj''})' \\ &\quad + F_{0j} F'_{qj'} F'_{qj''} [j'(j' + 1) + j''(j'' + 1)] \\ &\quad - F_{0j} F_{qj'} F_{qj''} \left[j'^2(j' + 1)^2 + j''^2(j'' + 1)^2 - \frac{q^2}{\cos^2 \frac{\theta}{2}} [j'(j' + 1) + j''(j'' + 1)] \right] . \end{aligned} \quad (\text{A.16})$$

Integrating by parts in the fourth term we get:

$$\begin{aligned} A_E &= 2F''_{0j} F'_{qj'} F'_{qj''} + F'_{0j} (F''_{qj'} F'_{qj''} + F'_{qj'} F''_{qj''}) + \frac{q^2}{\cos^2 \frac{\theta}{2}} F'_{0j} (F_{qj'} F_{qj''})' \\ &\quad - F'_{0j} [F_{qj'} F'_{qj''} j'(j' + 1) + F'_{qj'} F_{qj''} j''(j'' + 1)] \\ &\quad - F_{0j} F_{qj'} F_{qj''} (j'(j' + 1) - j''(j'' + 1))^2 . \end{aligned} \quad (\text{A.17})$$

Combining the third and fourth terms using (A.12) gives:

$$\begin{aligned} A_E &= 2F''_{0j} F'_{qj'} F'_{qj''} + F'_{0j} (F''_{qj'} F'_{qj''} + F'_{qj'} F''_{qj''}) + F'_{0j} F'_{qj''} \frac{1}{\sin \theta} (\sin \theta F'_{qj'})' \\ &\quad + F'_{0j} F'_{qj'} \frac{1}{\sin \theta} (\sin \theta F'_{qj''})' - F_{0j} F_{qj'} F_{qj''} (j'(j' + 1) - j''(j'' + 1))^2 . \end{aligned} \quad (\text{A.18})$$

Everything except for the last term is a total derivative, so

$$A_E = -F_{0j} F_{qj'} F_{qj''} (j'(j' + 1) - j''(j'' + 1))^2 . \quad (\text{A.19})$$

The simplified form for \mathcal{I}_E presented in (4.48) immediately follows. The simplification of \mathcal{I}_B is achieved through a similar integration by parts.

A.2 Check of gauge invariance

As discussed around (4.15), gauge invariance implies:

$$\begin{aligned} i\omega K_j^{q,\tau\tau}(\omega) &= -\sqrt{j(j+1)}K_j^{q,\tau E}(\omega), \\ i\omega K_j^{q,\tau E}(\omega) &= \sqrt{j(j+1)}K_j^{q,E}(\omega). \end{aligned} \tag{A.20}$$

We can check that the expression in (4.47) obey these relations by comparing residues at $\omega = \pm i(E_{qj'} + E_{qj''})$. We have

$$\text{Res}_{\omega=\pm i(E_{qj'}+E_{qj''})} \frac{1}{\omega^2 + (E_{qj'} + E_{qj''})^2} = \mp \frac{i}{E_{qj'} + E_{qj''}}. \tag{A.21}$$

From (4.47) and (4.48), one can check that, for instance, when $j' \neq j''$,

$$\begin{aligned} \text{Res}(i\omega K_j^{q,\tau\tau}(\omega)) &= i \frac{8\pi^2}{2j+1} \frac{(E_{qj'} - E_{qj''})(E_{qj'}^2 - E_{qj''}^2)}{2E_{qj'}E_{qj''}} \mathcal{I}_D(j, j', j'') \\ &= -\text{Res}\left(\sqrt{j(j+1)}K_j^{q,\tau E}(\omega)\right), \end{aligned} \tag{A.22}$$

so the first gauge invariance relation is verified. In checking this relation it was important that $\mathcal{I}_\phi(j, j', j'') = (E_{qj''}^2 - E_{qj'}^2)\mathcal{I}_D(j, j', j'')$. A similar check shows that the second gauge invariance condition in (A.20) is satisfied.

B Formulas for \mathcal{I}_B and \mathcal{I}_F

B.1 Results for $q = 0, 1/2$

We have only been able to perform the integrals as written in (4.48) for $q = 0, 1/2$. We had to use a different method to obtain results for arbitrary q ; the details are given in the next subsection. We give the results below.

For $q = 0$, we have

$$\begin{aligned} \mathcal{I}_B(j, j', j'') &= -\frac{(2j+1)(2j'+1)(2j''+1)(j'+j''-j)(j+j'-j'')(j+j''-j')}{32\pi^3} \\ &\quad \times \left[j + j' + j'' - \frac{1}{j + j' + j''} \right] \begin{pmatrix} j-1 & j'-1 & j''-1 \\ 0 & 0 & 0 \end{pmatrix}^2. \end{aligned} \tag{B.1}$$

For $q = 1/2$, we have

$$\begin{aligned}\mathcal{I}_F(j, j', j'') &= \mathcal{I}_D(j, j', j'') \times \begin{cases} j(j+1) - (j' - j'')^2 & \text{for } j + j' + j'' \text{ odd} \\ j(j+1) - (j' + j'' + 1)^2 & \text{for } j + j' + j'' \text{ even} \end{cases} \\ \mathcal{I}_B(j, j', j'') &= -\mathcal{I}_D(j, j', j'') \times \begin{cases} [(j' - j'')^2 - j(j+1)]^2 & \text{for } j + j' + j'' \text{ odd} \\ [(j' + j'' + 1)^2 - j(j+1)]^2 & \text{for } j + j' + j'' \text{ even} \end{cases}\end{aligned}\quad (\text{B.2})$$

B.2 Results for arbitrary q

For arbitrary q we were not able to perform the integrals in (4.48) directly. Rather, we employed a method developed in [33] to obtain expressions for the kernels. From these expressions we were able to read off explicit formulas for \mathcal{I}_B and \mathcal{I}_F . We briefly introduce the method below, and give the end results, which are somewhat complicated.

In this subsection we will use a different basis for monopole vector harmonics than the one used in the main text. We denote the harmonics in the new basis by $U_{q,jm}^\mu$, $V_{q,jm}^\mu$, $W_{q,jm}^\mu$. For completeness we provide the details needed for the calculation, but we do not elaborate on the technique. For a detailed discussion see [33].

The vector spherical harmonics that we need are tangent vectors to $S^2 \times \mathbb{R}$. There are a few natural choices of basis for this tangent space. The basis we used in the body of the text is just the basis induced from the round coordinates on $S^2 \times \mathbb{R}$, and is given explicitly in (A.1).

There is also a cartesian basis given by conformally mapping the natural frame basis of \mathbb{R}^3 to $S^2 \times \mathbb{R}$. We write the standard line element on \mathbb{R}^3 in spherical coordinates as

$$ds_{\mathbb{R}^3}^2 = d\vec{x}^2 = e^{2\tau} [d\tau^2 + d\theta^2 + \sin^2 \theta d\phi^2] \quad \vec{x} \equiv e^\tau \begin{pmatrix} \sin \theta \cos \phi & \sin \theta \sin \phi & \cos \theta \end{pmatrix}. \quad (\text{B.3})$$

The metric on $S^2 \times \mathbb{R}$ is obtained by rescaling the \mathbb{R}^3 metric (B.3) by $e^{-2\tau}$:

$$ds_{S^2 \times \mathbb{R}}^2 = d\theta^2 + \sin^2 \theta d\phi^2 + d\tau^2. \quad (\text{B.4})$$

We obtain the frame on $S^2 \times \mathbb{R}$ by the conformal transformation of the standard frame $e_{\mathbb{R}^3}^a = dx^a$ on \mathbb{R}^3 :

$$e^a = e^{-\tau} dx^a. \quad (\text{B.5})$$

Explicitly,

$$e_\mu^a = \begin{pmatrix} \cos(\phi) \sin(\theta) & \cos(\theta) \cos(\phi) & -\sin(\theta) \sin(\phi) \\ \sin(\theta) \sin(\phi) & \cos(\theta) \sin(\phi) & \cos(\phi) \sin(\theta) \\ \cos(\theta) & -\sin(\theta) & 0 \end{pmatrix} \quad (\text{B.6})$$

The frame satisfies the relation

$$e_\mu^a e_\nu^b g^{\mu\nu} = \delta^{ab}. \quad (\text{B.7})$$

There is yet another basis for the tangent space, in which the angular momentum operators, \vec{J}^2 , and J_z are diagonal. This basis is given by the vielbeins:

$$\begin{aligned} e^\pm &= \frac{1}{\sqrt{2}} e^{-\tau} (\mp dx + idy) \\ e^z &= e^{-\tau} dz, \end{aligned} \quad (\text{B.8})$$

with x, y, z coordinates on \mathbb{R}^3 .

In this raising and lowering basis the expressions of the vector spherical harmonics for $j > q$ are:

$$\begin{aligned} U_{q,jm}^s(\hat{n}) &= \begin{pmatrix} \sqrt{\frac{(j-m+1)(j-m+2)}{(2j+2)(2j+3)}} Y_{q,j+1,m-1}(\hat{n}) \\ -\sqrt{\frac{(j-m+1)(j+m+1)}{(j+1)(2j+3)}} Y_{q,j+1,m}(\hat{n}) \\ \sqrt{\frac{(j+m+1)(j+m+2)}{(2j+2)(2j+3)}} Y_{q,j+1,m+1}(\hat{n}) \end{pmatrix}, \\ V_{q,jm}^s(\hat{n}) &= \begin{pmatrix} -\sqrt{\frac{(j-m+1)(j+m)}{2j(j+1)}} Y_{q,j,m-1}(\hat{n}) \\ \frac{m}{\sqrt{j(1+j)}} Y_{q,jm}(\hat{n}) \\ \sqrt{\frac{(j-m)(j+m+1)}{2j(j+1)}} Y_{q,j,m+1}(\hat{n}) \end{pmatrix}, \\ W_{q,jm}^s(\hat{n}) &= \begin{pmatrix} \sqrt{\frac{(j+m-1)(j+m)}{2j(2j-1)}} Y_{q,j-1,m-1}(\hat{n}) \\ \sqrt{\frac{(j-m)(j+m)}{j(2j-1)}} Y_{q,j-1,m}(\hat{n}) \\ \sqrt{\frac{(j-m-1)(j-m)}{2j(2j-1)}} Y_{q,j-1,m+1}(\hat{n}) \end{pmatrix}. \end{aligned} \quad (\text{B.9})$$

Here, U, V, W all have total angular momentum j , and s runs over $\{+, z, -\}$.

If $j = q$, we only have two modes $U_{q,jm}^s(\hat{n})$ and $V_{q,jm}^s(\hat{n})$. If $j = q = 0$, only the $U_{q,jm}^s(\hat{n})$ mode is non vanishing. For $q \geq 1$, if we have $j = q - 1$ only the mode $U_{q,jm}^s(\hat{n})$ is non-zero. We don't encounter this case in this paper. From the raising and lowering basis we

can go to the coordinate basis by contracting these expressions with e_s^μ in (B.8). Then $U_{q,jm}^\mu, V_{q,jm}^\mu, W_{q,jm}^\mu$ are themselves vectors of ordinary monopole harmonics (multiplied by some coordinate dependent factors coming from the e_s^μ).

A key formula that we will use is

$$D^\mu (Y_{q,jm}(\hat{n})e^{-i\omega\tau}) = \left((j+i\omega)\sqrt{\frac{(j+1)^2-q^2}{(j+1)(2j+1)}} \quad -\frac{q(1-i\omega)}{\sqrt{j(j+1)}} \quad (j+1-i\omega)\sqrt{\frac{j^2-q^2}{j(2j+1)}} \right) \begin{pmatrix} U_{q,jm}^\mu e^{-i\omega\tau} \\ V_{q,jm}^\mu e^{-i\omega\tau} \\ W_{q,jm}^\mu e^{-i\omega\tau} \end{pmatrix}, \quad (\text{B.10})$$

where we have to take the scalar product with the coefficient vector.

After these preparatory steps we can present our method. We want to calculate the matrix elements of the kernel, $\mathbf{K}_j^q(\omega)$, but now in the basis given by (B.9). To do this we have to invert (4.10). As explained below (4.20), using rotational symmetry we can average over the quantum number m to get a formula that is easier to treat:

$$[\mathbf{K}_j^q(\omega)]_{XZ} = \frac{1}{2j+1} \sum_{m=-j}^j \int d^3x d^3x' \sqrt{g(x)} \sqrt{g(x')} X_{jm}^{\mu*}(\hat{n}) \mathcal{K}_{\mu\mu'}^q(x, x') Z_{jm}^{\mu'}(\hat{n}') e^{i\omega(\tau-\tau')}, \quad (\text{B.11})$$

where $X, Z \in \{U, V, W\}$.

$\mathcal{K}_{\mu\mu'}^q(x, x')$ is given in terms of Green's functions in (4.7). We use the first line of (4.38) to do the Fourier decomposition of $G^q(x, x')$ instead of the second line, which we used in the rest of the paper. The benefit of treating x and x' separately, rather than introducing the relative angle, γ is an algorithmic method to evaluate the kernels at the expense of having long formulas.

Plugging into (4.7) we get

$$\begin{aligned} \mathcal{K}_{\mu\mu'}^q(x, x') = & \sum_{\substack{j', m' \\ j'', m''}} \int \frac{d\omega'}{2\pi} \frac{d\omega''}{2\pi} G_{j'}(\omega') G_{j''}(\omega'') \left[D_\mu \left(Y_{q,j'm'}(\hat{n}) e^{-i\omega'\tau} \right) Y_{q,j''m''}^*(\hat{n}') e^{i\omega''\tau} \right] \\ & \times \left[Y_{q,j'm'}^*(\hat{n}') e^{i\omega'\tau'} D_{\mu'} \left(Y_{q,j''m''}(\hat{n}'') e^{-i\omega''\tau'} \right) \right] + \text{other distribution of derivatives}, \end{aligned} \quad (\text{B.12})$$

where we only wrote down explicitly the term coming from the first term in (4.7), and we grouped spherical harmonics at the same point inside square brackets. For the derivative of

the spherical harmonics we can use (B.10), and plugging this into (B.11) we get an expression involving one scalar and two vector spherical harmonics at the two spacetime points x, x' . We can then use (B.9) to reduce the whole expression to the sum of products involving six ordinary spherical harmonics, three at each spacetime point x, x' .¹² As anticipated this is a tedious, but algorithmic task.

We can actually perform the integral over x' with no work. Because we averaged over m in (B.11), the integrand depends only on the relative angle between \hat{n} and \hat{n}' . The integral with respect to \hat{n} is therefore independent of \hat{n}' , so we can choose \hat{n}' to point in the \hat{z} direction and replace the integral with respect to \hat{n}' by a factor of 4π . Using

$$Y_{q,\ell m}(\hat{z}) = \delta_{q,-m} \sqrt{\frac{2\ell+1}{4\pi}}, \quad (\text{B.13})$$

we get rid of three monopole harmonics. The remaining angular integral over the product of three harmonics can be evaluated using some properties of monopole harmonics [45].

$$Y_{q,\ell m}(\hat{n})^* = (-1)^{q+m} Y_{-q,\ell,-m}(\hat{n}), \quad (\text{B.14})$$

and

$$\begin{aligned} & \int d\hat{n} Y_{q,\ell m}(\hat{n}) Y_{q',\ell' m'}(\hat{n}) Y_{q'',\ell'' m''}(\hat{n}) \\ &= (-1)^{\ell+\ell'+\ell''} \sqrt{\frac{(2\ell+1)(2\ell'+1)(2\ell''+1)}{4\pi}} \begin{pmatrix} \ell & \ell' & \ell'' \\ q & q' & q'' \end{pmatrix} \begin{pmatrix} \ell & \ell' & \ell'' \\ m & m' & m'' \end{pmatrix}, \end{aligned} \quad (\text{B.15})$$

where $\begin{pmatrix} j & j' & j'' \\ m & m' & m'' \end{pmatrix}$ is the Wigner $3j$ -symbol.

After performing the frequency integrals over ω', ω'' , we get the answer as a sum of a large number of $3j$ -symbols, as many of the angular momentum indices on $Y_{q,jm}$ are shifted in (B.9). We can use some identities to reduce the number of $3j$ -symbols that appear in our final answer, but we are still left with long expressions. The same logic also gives $[\mathbf{F}_j^q(\omega)]_X$, and $D_j^q(\omega)$, though the latter does not involve any vector harmonics.

There is an easy translation between the integrals and the kernels (4.47). Because $V_{0,jm}$ is identical to the B -mode of the gauge field, we can read off \mathcal{I}_F and \mathcal{I}_B from $[\mathbf{F}_j^q(\omega)]_V$ and $[\mathbf{K}_j^q(\omega)]_{VV}$. To simplify the expressions somewhat, we introduce the following notation for

¹²It is more economical to use the frame basis (B.9) to contract vector indices between the vector harmonics.

the product of two $3j$ -symbol that will appear in our formulas

$$\begin{bmatrix} j & j'' & j' \\ 0 & m & -m \\ 0 & n & -n \end{bmatrix} \equiv \begin{pmatrix} j & j'' & j' \\ 0 & m & -m \end{pmatrix} \times \begin{pmatrix} j & j'' & j' \\ 0 & -n & n \end{pmatrix} \quad (\text{B.16})$$

With this notation, we have:

$$\mathcal{I}_F(j, j', j'') = \frac{1}{64q\pi^3} \times \left\{ \begin{array}{l} -(2j+1)(2j'+1)(2j''+1) \sqrt{(j'-q+1)(j'+q)(j''-q+1)(j''+q)} \times \begin{bmatrix} j & j'' & j' \\ 0 & -q & q \\ 0 & q-1 & 1-q \end{bmatrix} \\ -2(2j+1)q(2j'+1)(2j''+1) \times \begin{bmatrix} j & j'' & j' \\ 0 & -q & q \\ 0 & q & -q \end{bmatrix} \\ +(2j+1)(2j'+1)(2j''+1) \sqrt{(j'-q)(j'+q+1)(j''-q)(j''+q+1)} \times \begin{bmatrix} j & j'' & j' \\ 0 & -q & q \\ 0 & q+1 & -q-1 \end{bmatrix} \end{array} \right\} \quad (\text{B.17})$$

$$\begin{aligned}
\mathcal{I}_B(j, j', j'') &= \frac{1}{128qj'(j'+1)j''(j''+1)\pi^3} \times \\
&\left(\begin{aligned}
&\left(\frac{-(2j+1)j'(j'+1)(2j'+1)(j''+1)^2(q+j'')}{\times \sqrt{(-q^2+q+j'^2+j')(-q^2+q+j''^2-j'')(-j^2-j+j'^2+j'+j''(-2q+j''-1))}} \right) \\
&+ \left(\frac{-2(2j+1)j'(j'+1)(2j'+1)(j''+1)^2(j''-q)}{\times (q+j'')(j^2+j-j''(2q^2+j''-1)+j'^2(2j''-1)+j'(2j''-1))} \right) \\
&+ \left(\frac{(2j+1)j'(j'+1)(2j'+1)(q-j'')(j''+1)^2}{\times \sqrt{-(j'^2+j'-q(q+1))(q^2+q-j''^2+j'')(-j^2-j+j'^2+j'+j''(2q+j''-1))}} \right) \\
&+ \left(\frac{-(2j+1)(j'+1)^2(q+j')j''(j''+1)(2j''+1)}{\times \sqrt{(-q^2+q+j'^2-j')(-q^2+q+j''^2+j'')(-j^2-j+j'^2+j''^2-(2q+1)j'+j'')}} \right) \\
&+ \left(\frac{(2j+1)q(2j'+1)(2j''+1)\sqrt{(-q^2+q+j'^2+j')(-q^2+q+j''^2+j'')}}{\times (j'^4+2j'^3-(j^2+j+2j'^2+2j''-1)j'^2-(j^2+j+2j''(j''+1))j'+j''(j''+1)(-j^2-j+j'^2+j''))} \right) \\
&+ \left(\frac{-(2j+1)j'^2(-q+j'+1)j''(j''+1)(2j''+1)}{\times \sqrt{(-q^2+q+j'^2+3j'+2)(-q^2+q+j''^2+j'')(-j^2-j+j'^2+j''^2+2q+(2q+3)j'+j''+2)}} \right) \\
&+ \left(\frac{2(2j+1)(j'+1)^2(j'-q)(q+j')j''(j''+1)}{\times (2j''+1)(-j^2-j+j'^2+(j'')^2+j''-j'(-2q^2+2j'^2+2j''+1))} \right) \\
&+ \left(\frac{2(2j+1)q^2(2j'+1)(2j''+1)}{\times (j'^4+2j'^3-(j^2+j+2j'^2+2j''-1)j'^2-(j^2+j+2j''(j''+1))j'+j''(j''+1)(-j^2-j+j'^2+j''))} \right) \\
&+ \left(\frac{-2(2j+1)j'^2(-q+j'+1)(q+j'+1)j''(j''+1)}{\times (2j''+1)(-j^2-j-2q^2+j'^2+3j'^2+3j''+j'(-2q^2+2j'^2+2j''+3)+2)} \right) \\
&+ \left(\frac{-(2j+1)(j'+1)^2(j'-q)j''(j''+1)\sqrt{(-q+j'-1)(q+j')(j''-q)(q+j''+1)}}{\times (2j''+1)(-j^2-j+j'^2+(j'')^2+(2q-1)j'+j'')} \right) \\
&+ \left(\frac{-(2j+1)q(2j'+1)\sqrt{(j'-q)(q+j'+1)(j''-q)(q+j''+1)}(2j''+1)}{\times (j'^4+2j'^3-(j^2+j+2j'^2+2j''-1)j'^2-(j^2+j+2j''(j''+1))j'+j''(j''+1)(-j^2-j+j'^2+j''))} \right) \\
&+ \left(\frac{-(2j+1)j'^2(q+j'+1)j''(j''+1)(2j''+1)}{\times \sqrt{(-q^2-q+j'^2+3j'+2)(j''^2+j''-q(q+1))(-j^2-j+j'^2+j''^2-2q+(3-2q)j'+j''+2)}} \right) \\
&+ \left(\frac{(2j+1)j'(j'+1)(2j'+1)(q-j''-1)j''^2}{\times \sqrt{(-q^2+q+j'^2+j')(-q^2+q+j''^2+3j''+2)}(-j^2-j+(j')^2+j'+(j''+1)(2q+j''+2))} \right) \\
&+ \left(\frac{-2(2j+1)j'(j'+1)(2j'+1)j''^2(-q+j''+1)}{\times (q+j''+1)(-j^2-j-2q^2+j''^2-2q^2j''+3j''+j'^2(2j''+3)+j'(2j''+3)+2)} \right) \\
&+ \left(\frac{-(2j+1)j'(j'+1)(2j'+1)j''^2(q+j''+1)}{\times \sqrt{(j'-q)(q+j'+1)(-q+j''+1)(q+j''+2)}(-j^2-j+j'^2+j'+(j''+1)(-2q+j''+2))} \right)
\end{aligned} \right) \times \left. \begin{aligned}
&\begin{bmatrix} j & j''-1 & j' \\ 0 & -q & q \\ 0 & q-1 & 1-q \end{bmatrix} \\
&\times \begin{bmatrix} j & j''-1 & j' \\ 0 & -q & q \\ 0 & q & -q \end{bmatrix} \\
&\times \begin{bmatrix} j & j''-1 & j' \\ 0 & -q & q \\ 0 & q+1 & -q-1 \end{bmatrix} \\
&\times \begin{bmatrix} j & j'' & j'-1 \\ 0 & -q & q \\ 0 & q-1 & 1-q \end{bmatrix} \\
&\times \begin{bmatrix} j & j'' & j' \\ 0 & -q & q \\ 0 & q-1 & 1-q \end{bmatrix} \\
&\times \begin{bmatrix} j & j'' & j'+1 \\ 0 & -q & q \\ 0 & q-1 & 1-q \end{bmatrix} \\
&\times \begin{bmatrix} j & j'' & j'-1 \\ 0 & -q & q \\ 0 & q & -q \end{bmatrix} \\
&\times \begin{bmatrix} j & j'' & j' \\ 0 & -q & q \\ 0 & q & -q \end{bmatrix} \\
&\times \begin{bmatrix} j & j'' & j'+1 \\ 0 & -q & q \\ 0 & q & -q \end{bmatrix} \\
&\times \begin{bmatrix} j & j'' & j'-1 \\ 0 & -q & q \\ 0 & q+1 & -q-1 \end{bmatrix} \\
&\times \begin{bmatrix} j & j'' & j' \\ 0 & -q & q \\ 0 & q+1 & -q-1 \end{bmatrix} \\
&\times \begin{bmatrix} j & j'' & j'+1 \\ 0 & -q & q \\ 0 & q+1 & -q-1 \end{bmatrix} \\
&\times \begin{bmatrix} j & j''+1 & j' \\ 0 & -q & q \\ 0 & q-1 & 1-q \end{bmatrix} \\
&\times \begin{bmatrix} j & j''+1 & j' \\ 0 & -q & q \\ 0 & q & -q \end{bmatrix} \\
&\times \begin{bmatrix} j & j''+1 & j' \\ 0 & -q & q \\ 0 & q+1 & -q-1 \end{bmatrix}
\end{aligned} \right)
\end{aligned}
\tag{B.18}$$

One can check that for $q = 0, 1/2$ we get back the results of the previous subsection.

C Asymptotic expansions

The goal of this Appendix is to derive an asymptotic formula at large j and ω for the integrand in the expression (4.19) for $\delta\mathcal{F}_q$.

C.1 Small distance expansion of the Green's function

We start by obtaining a better understanding of the scalar Green's function (4.38). Isolating the phase factor appearing in (4.38), we can write

$$G^q(x, x') = e^{-2iq\Theta} \tilde{G}_q(\tau - \tau', \gamma), \quad \tilde{G}_q(\tau - \tau', \gamma) \equiv \sum_{j=q}^{\infty} F_{q,j}(\gamma) \frac{e^{-E_{qj}|\tau - \tau'|}}{2E_{qj}}, \quad (\text{C.1})$$

where γ and Θ are defined in (4.24) and (4.40), respectively, and the energy E_{qj} was defined in (4.41). Note that when $\theta' = \phi' = \tau' = 0$, we have $G_q(x, 0) = \tilde{G}_q(\tau, \theta)$.

The Green's function satisfies the differential equation

$$\left[\partial_\tau^2 + (\nabla_\mu - i\mathcal{A}_\mu^q)(\nabla^\mu - i\mathcal{A}^{q,\mu}) - \left(\mu^2 + \frac{1}{4} \right) \right] \tilde{G}_q(\tau, \theta) = -\frac{1}{2\pi \sin \theta} \delta(\tau) \delta(\theta), \quad (\text{C.2})$$

where $\mathcal{A}^q = q(1 - \cos \theta)d\phi$. It is convenient to change variables to

$$t = \sinh^2 \frac{\tau}{2}, \quad s = \sin^2 \frac{\theta}{2}, \quad (\text{C.3})$$

and solve eq. (C.2) at small t and s (where t and s are considered to be of the same order).

The first few terms in this series expansion are

$$\begin{aligned} \tilde{G}_q(s, t) &= \frac{1}{8\pi\sqrt{s+t}} + C_q + \frac{\mu_q^2\sqrt{s+t}}{4\pi} + \left[A_q t + \frac{s(-2A_q + C_q(1 + 4\mu_q^2))}{4} \right] \\ &+ \sqrt{s+t} \frac{(q^2 + 2\mu_q^2)(2s-t) + 3\mu_q^4(s+t)}{36\pi} \\ &+ B_q t^2 + st \left[A_q \left(\mu_q^2 - \frac{3}{4} \right) - 3B_q \right] + s^2 \frac{24B_q - 4A_q(3 + 4\mu_q^2) + C_q(9 + 16q^2 + 40\mu_q^2 + 16\mu_q^4)}{64} \\ &+ \dots, \end{aligned} \quad (\text{C.4})$$

where C_q , A_q , and B_q are integration constants. At each integer order in the expansion, we have one new integration constant that can be chosen to be the coefficient of t^n . When $s = \gamma = 0$, we have $F_{q,j}(0) = (2j + 1)/(4\pi)$, so

$$\tilde{G}_q(0, \tau) = \frac{1}{8\pi\sqrt{t}} + C_q + \frac{\mu_q^2\sqrt{t}}{4\pi} + A_q t + t^{3/2} \frac{3\mu_q^4 - 2\mu_q^2 - q^2}{36\pi} + B_q t^2 + \dots \quad (\text{C.5})$$

The integration constants can be determined from the spectral decomposition (C.1). Indeed, for $s = \gamma = 0$, we have

$$\tilde{G}_q(0, \tau) = \frac{1}{4\pi} \sum_{j=q}^{\infty} (j + 1/2) \frac{e^{-E_{qj}|\tau|}}{E_{qj}}, \quad (\text{C.6})$$

and this expression can be expanded at small τ and matched with (C.5). The small τ expansion is subtle, however, because naively expanding the summand at small τ results in divergent sums.

In order to only work with absolutely convergent sums, we can first consider the quantity

$$\begin{aligned} \tilde{G}_q^{\text{sub}} &= \frac{1}{4\pi} \sum_{n=q+1/2}^{\infty} e^{-n|\tau|} \left[1 - \frac{1 + |\tau|n}{2n^2} (\mu_q^2 - q^2) + \frac{3 + 3n|\tau| + n^2\tau^2}{8n^4} (\mu_q^2 - q^2)^2 + \dots \right] \\ &+ \frac{1}{4\pi} \sum_{n=q+1/2}^{\infty} \left[\frac{1}{2n^2} (\mu_q^2 - q^2) + \frac{n^2\tau^2 - 6}{16n^4} (\mu_q^2 - q^2)^2 + \dots \right]. \end{aligned} \quad (\text{C.7})$$

\tilde{G}_q^{sub} is a power series in $\mu_q^2 - q^2$. The first line in this expression was obtained by expanding $E_{qj} = \sqrt{(j + 1/2)^2 + \mu_q^2 - q^2}$ at small $\mu_q^2 - q^2$ and plugging this expansion into (C.6), thereby obtaining an expansion of $\tilde{G}_q(0, \tau)$. The expression in the second line subtracts the first few terms in the small τ expansion of the first line: at $(\mu_q^2 - q^2)^p$ order it contains terms up to order $\tau^{2(p-1)}$.¹³ The sums in (C.7) can be performed analytically, and then expanded at small t :

$$\begin{aligned} \tilde{G}_q^{\text{sub}} &= \frac{1}{4\pi} \left[\frac{1}{2\sqrt{t}} + \mu_q^2\sqrt{t} + t^{3/2} \frac{q^2 + 2\mu_q^2 - 3\mu_q^4}{9} + \dots \right] \\ &+ \frac{1}{4\pi} \left[-q + t \frac{q(1 + 2q^2 - 6\mu_q^2)}{6} + \dots \right], \end{aligned} \quad (\text{C.8})$$

where in the first line we isolated the terms non-analytic in t , and in the second line we

¹³Note that this implies that the $O((\mu_q^2 - q^2)^0)$ term is absent in the second line.

included the terms analytic in t . The non-analytic terms in this expression match exactly the non-analytic terms in (C.5), so the difference $\tilde{G}_q(0, \tau) - \tilde{G}_q^{\text{sub}}$ is analytic in t . In fact, using (C.5) and (C.7), it is not hard to see that the difference $\tilde{G}_q(0, \tau) - \tilde{G}_q^{\text{sub}}$ and its first few derivatives w.r.t. τ (or t) are expressed as absolutely convergent sums, so one can expand at small t (or small τ) by expanding the summands. One finds

$$\begin{aligned} \tilde{G}_q(0, \tau) - \tilde{G}_q^{\text{sub}} &= \frac{1}{4\pi} \sum_{n=q+1/2}^{\infty} \left[\left(\frac{n}{E_{(n-\frac{1}{2})q}} - 1 \right) + t \left(2nE_{(n-\frac{1}{2})q} - \mu_q^2 + q^2 - 2n^2 \right) \right. \\ &\quad \left. + \frac{2t^2}{3} \left(nE_{(n-\frac{1}{2})q}^3 - nE_{(n-\frac{1}{2})q} - n^2(n^2 - 1) - \frac{1}{2}(3n^2 - 1)(\mu_q^2 - q^2) - \frac{3}{8}(\mu_q^2 - q^2)^2 \right) + \dots \right]. \end{aligned} \quad (\text{C.9})$$

Adding back (C.8), we can obtain an expression for $\tilde{G}_q(0, \tau)$. It can be checked that this expression can be written more succinctly as

$$\begin{aligned} \tilde{G}_q(0, \tau) &= \frac{1}{4\pi} \left[\frac{1}{2\sqrt{t}} + \mu_q^2 \sqrt{t} + t^{3/2} \frac{q^2 + 2\mu_q^2 - 3\mu_q^4}{9} + \dots \right] \\ &\quad + \frac{1}{4\pi} \left[\frac{S_{-1}}{2} + S_1 t + \frac{1}{3}(S_3 - S_1)t^2 + \dots \right], \end{aligned} \quad (\text{C.10})$$

where the first line contains the terms non-analytic in t , and in the second line S_p is defined as the zeta-function regularized sum

$$S_p \equiv \sum_{j=q}^{\infty} (2j+1)(E_{qj})^p. \quad (\text{C.11})$$

In particular,

$$\begin{aligned} S_{-1} &= -\frac{q}{2} + \sum_{j=q}^{\infty} \left[\frac{2j+1}{E_{qj}} - 2 \right], \\ S_1 &= \frac{q(1+2q^2-6\mu_q^2)}{6} + \sum_{j=q}^{\infty} \left[(2j+1)E_{qj} - 2(j+1/2)^2 - \mu_q^2 + q^2 \right], \\ S_3 &= \frac{q(-7+10q^2-18q^4+30\mu_q^2(1+2q^2)-90\mu_q^4)}{240} \\ &\quad + \sum_{j=q}^{\infty} \left[(2j+1)(E_{qj})^3 - 2(j+1/2)^4 - 3(\mu_q^2 - q^2)(j+1/2)^2 - \frac{3}{4}(\mu_q^2 - q^2)^2 \right], \end{aligned} \quad (\text{C.12})$$

and so on. These expressions can easily be evaluated numerically. By comparing (C.10) to (C.5), we can extract the constants C_q , B_q , and A_q appearing in (C.10):

$$C_q = \frac{S_{-1}}{8\pi}, \quad A_q = \frac{S_1}{4\pi}, \quad B_q = \frac{S_3 - S_1}{12\pi}. \quad (\text{C.13})$$

Note that the constant C_q is the same as that appearing in (4.57). From (3.7) and (3.8), it is easy to see that if one tunes to criticality we have

$$C_q = 0, \quad A_q = \frac{\mathcal{F}_q^\infty}{4\pi}. \quad (\text{C.14})$$

The constant B_q can be computed numerically for any given q . We have $B_{1/2} \approx -0.00475009$ and $B_1 \approx -0.01255981$.

With (C.13), we now have a complete expression for the small distance expansion (C.4) of the Green's function. This expansion can be developed to higher orders if needed, and in fact the final formulas presented below were obtained after keeping one more order in (C.4).

C.2 UV asymptotic of the scalar kernel

Let us first examine the scalar kernel $D^q(x, x') = |G^q(x, x')|^2$. From (4.22) and (C.1), we have

$$D_j^q(\omega) = 2\pi \int d\tau \int d\theta \sin \theta e^{i\omega\tau} P_j(\cos \theta) \tilde{G}_q(\tau, \theta)^2. \quad (\text{C.15})$$

The large j and ω behavior of $D_j^q(\omega)$ can be obtained by plugging in the expansion (C.4) in (C.15) and evaluating the integrals *provided* that the terms we're Fourier transforming are non-analytic in s and t . (The analytic terms in s and t are *not* related in any way to the large ω and j behavior of $D_j^q(\omega)$.)

So at large ω and j we have

$$D_j^q(\omega) = 2\pi \int d\tau \int d\theta \sin \theta e^{i\omega\tau} P_j(\cos \theta) \left[\frac{1}{64\pi^2(s+t)} - \frac{\mathcal{F}_q^\infty(s-2t)}{32\pi^2\sqrt{s+t}} + \frac{8\pi B_q(3s^2 - 24st + 8t^2) + \mathcal{F}_q^\infty [32t^2\mu_q^2 + 4st(-3 + 8\mu_q^2) - s^2(3 + 20\mu_q^2)]}{256\pi^2\sqrt{s+t}} + \dots \right] \quad (\text{C.16})$$

In order to evaluate these integrals asymptotically at large ω and j , it is convenient to

re-expand the term inside the square brackets in terms of simple functions of

$$X \equiv \sqrt{2(\cosh \tau - \cos \theta)} = 2\sqrt{s+t}. \quad (\text{C.17})$$

For instance, we can write the term inside the square brackets as

$$\begin{aligned} D_j^q(\omega) = \text{F.T.} \left[\frac{1}{16\pi^2 X^2} - \frac{\mathcal{F}_q^\infty}{192\pi^2} (\nabla_{S^2}^2 - 2\partial_\tau^2) X^3 + \frac{B_q}{26880\pi} (8\partial_\tau^4 - 24\partial_\tau^2 \nabla_{S^2}^2 + 3\nabla_{S^2}^4) X^7 \right. \\ \left. + \frac{\mathcal{F}_q^\infty}{1290240\pi^2} (-38\partial_\tau^4 + (29 + 96\mu_q^2)\partial_\tau^2 \nabla_{S^2}^2 - (23 + 24\mu_q^2)\nabla_{S^2}^4) X^7 + \dots \right], \end{aligned} \quad (\text{C.18})$$

where we introduced the definition of the $S^2 \times \mathbb{R}$ Fourier transform of a function $f(\theta, \tau)$ as

$$\text{F.T.} [f] \equiv 2\pi \int d\tau d\theta \sin \theta e^{i\omega\tau} P_j(\cos \theta) f(\theta, \tau). \quad (\text{C.19})$$

Eqs. (C.18) and (C.16) agree up to the order in s and t to which (C.16) was valid.

Note that the $S^2 \times \mathbb{R}$ Fourier transform satisfies the properties

$$\begin{aligned} \text{F.T.} [\partial_\tau^2 f] &= -\omega^2 \text{F.T.} [f], \\ \text{F.T.} [\nabla_{S^2}^2 f] &= -j(j+1) \text{F.T.} [f], \end{aligned} \quad (\text{C.20})$$

which can be derived upon integration by parts twice in (C.19). These properties, together with the explicit Fourier transforms of powers of X given in Appendix C.6 give

$$\begin{aligned} D_j^q(\omega) = D_j^0(\omega) + \frac{\mathcal{F}_q^\infty}{2\pi} \frac{(j + \frac{1}{2})^2 - 2\omega^2}{[(j + \frac{1}{2})^2 + \omega^2]^3} + 6B_q \frac{3(j + \frac{1}{2})^4 - 24\omega^2(j + \frac{1}{2})^2 + 8\omega^4}{[(j + \frac{1}{2})^2 + \omega^2]^5} \\ + \frac{\mathcal{F}_q^\infty}{8\pi} \frac{8(j + \frac{1}{2})^4(1 - 3\mu_q^2) + 3\omega^2(j + \frac{1}{2})^2(32\mu_q^2 - 23) + 25\omega^4}{[(j + \frac{1}{2})^2 + \omega^2]^5} + O\left(\frac{1}{[(j + \frac{1}{2})^2 + \omega^2]^{7/2}}\right). \end{aligned} \quad (\text{C.21})$$

Here, $D_j^0(\omega)$ is the quantity defined in (4.28). Asymptotically, at large ω and j ,

$$D_j^0(\omega) = \frac{1}{8\sqrt{(j+\frac{1}{2})^2 + \omega^2}} + \frac{\omega^2 - (j+\frac{1}{2})^2}{64[(j+\frac{1}{2})^2 + \omega^2]^{5/2}} + \frac{11(j+\frac{1}{2})^4 - 62\omega^2(j+\frac{1}{2})^2 + 11\omega^4}{1024[(j+\frac{1}{2})^2 + \omega^2]^{9/2}} + O\left(\frac{1}{[(j+\frac{1}{2})^2 + \omega^2]^{7/2}}\right). \quad (\text{C.22})$$

C.3 The mixed kernel

Next, we examine the mixed kernel $F_\mu^q(x, x')$ defined in (4.7). Using $G^q(x, x') = e^{-2iq\Theta}\tilde{G}$, we can write

$$\begin{aligned} D'G^q &= e^{-2iq\Theta} [d' + i(\mathcal{A}^q)' - 2iqd'\Theta] \tilde{G}_q, \\ D'G^{q*} &= e^{2iq\Theta} [d' - i(\mathcal{A}^q)' + 2iqd'\Theta] \tilde{G}_q. \end{aligned} \quad (\text{C.23})$$

The definition of F_μ^q then implies

$$F_\mu^q(x, x') = 2i [\mathcal{A}_\mu^q(x') - 2\partial'_\mu\Theta(x, x')] \tilde{G}_q(\tau - \tau', \gamma)^2. \quad (\text{C.24})$$

From this expression it can be easily seen that $F_\tau^q(x, x') = 0$, which explicitly verifies the argument that $F_j^{q,\tau}(\omega) = F_j^{q,E}(\omega) = 0$ because of CP symmetry. For $F_j^{q,B}(\omega)$, we have

$$\begin{aligned} F_j^{q,B}(\omega) &= \frac{4\pi}{2j+1} \int d\tau d\theta d\phi \sin\theta \lim_{x' \rightarrow 0} \left[e^{i\omega(\tau-\tau')} \sum_{m=-j}^j Y_{jm}^*(\hat{x}) F_{\mu'}^q(x, x') \mathcal{Y}_{jm}^{\mu'}(\hat{x}') \right] \\ &= \frac{4\pi q}{\sqrt{j(j+1)}} \int d\tau d\theta \sin\theta e^{i\omega\tau} \tilde{G}_q^2(\tau, \theta) P_j^1(\cos\theta) \tan\frac{\theta}{2} \end{aligned} \quad (\text{C.25})$$

We should use

$$P_j^1(\cos\theta) = -\sin\theta P_j'(\cos\theta) \quad (\text{C.26})$$

and then

$$F_j^{q,B}(\omega) = -\frac{4\pi q}{\sqrt{j(j+1)}} \int d\tau d\theta \sin\theta e^{i\omega\tau} \tilde{G}_q^2(\tau, \theta) P_j'(\cos\theta) (1 - \cos\theta). \quad (\text{C.27})$$

We can integrate by parts and obtain

$$\begin{aligned}
F_j^{q,B}(\omega) &= \frac{2\pi q}{\sqrt{j(j+1)}} \int d\tau d\theta \sin\theta e^{i\omega\tau} P_j(\cos\theta) \frac{d}{d\cos\theta} \left[2\tilde{G}_q^2(\tau, \theta) (1 - \cos\theta) \right] \\
&= \text{F.T.} \left[\frac{d}{d\cos\theta} \left[2\tilde{G}_q^2(\tau, \theta) (1 - \cos\theta) \right] \right],
\end{aligned} \tag{C.28}$$

where we used the Fourier transform definition in (C.19). We can use the expansion of the Green's function (C.4), re-expanded in terms of $X = \sqrt{2(\cosh\tau - \cos\theta)}$ as in the scalar kernel case, to write

$$\begin{aligned}
F_j^{q,B}(\omega) &= \frac{q}{16\pi^2 \sqrt{j(j+1)}} \text{F.T.} \left[-\nabla_{S^2} \log X + \frac{1}{160} (4\partial_\tau^2 \nabla_{S^2}^2 - \nabla_{S^2}^4) (X^4 \log X) \right. \\
&\quad \left. + \frac{\mathcal{F}_q^\infty}{180} (2\nabla_{S^2}^4 - 7\partial_\tau^2 \nabla_{S^2}^2) X^5 + \dots \right].
\end{aligned} \tag{C.29}$$

This expression agrees with (C.28) in a small s and t expansion up to analytic terms in s and t . Using the Fourier transform properties (C.20) and the formulas in Appendix C.6, we can evaluate the integrals in (C.29) and expand at large ω and j . We obtain

$$\begin{aligned}
\left| F_j^{q,B}(\omega) \right|^2 &= q^2 \frac{(j + \frac{1}{2})^2}{64 [(j + \frac{1}{2})^2 + \omega^2]^3} - q^2 \frac{2(j + \frac{1}{2})^4 - 17\omega^2(j + \frac{1}{2})^2 + \omega^4}{256 [(j + \frac{1}{2})^2 + \omega^2]^5} \\
&\quad + O\left(\frac{1}{[(j + \frac{1}{2})^2 + \omega^2]^{7/2}} \right).
\end{aligned} \tag{C.30}$$

C.4 Gauge field kernel

For the gauge field kernel, we have

$$\begin{aligned}
K_{\mu\mu'}^q(x, x') &= 2\partial_\mu \tilde{G}_q(\tau - \tau', \gamma) \partial_{\mu'} \tilde{G}_q(\tau - \tau', \gamma) - 2\tilde{G}_q(\tau - \tau', \gamma) \partial_\mu \partial_{\mu'} \tilde{G}_q(\tau - \tau', \gamma) \\
&\quad - 4 [\mathcal{A}_\mu^q(x) + 2\partial_\mu \Theta(x, x')] [\mathcal{A}_{\mu'}^q(x') - 2\partial_{\mu'} \Theta(x, x')] \tilde{G}_q(\tau - \tau', \gamma)^2 \\
&\quad + 2g_{\mu\mu'} \delta(x - x') G^q(x, x').
\end{aligned} \tag{C.31}$$

It is easiest to start with calculating the Fourier modes of $K_{\tau\tau'}^q$ defined by

$$K_{\tau\tau'}^q(x, x') = \int \frac{d\omega}{2\pi} \sum_{j,m} K_j^{q,\tau\tau}(\omega) Y_{jm}(\hat{x}) Y_{jm}^*(\hat{x}') e^{-i\omega(\tau - \tau')}. \tag{C.32}$$

Using (C.31), we can extract $K_j^{q,\tau\tau}(\omega)$ from

$$K_j^{q,\tau\tau}(\omega) = 2G^q(x, x) + 2\pi \int d\tau \int d\theta \sin \theta e^{i\omega\tau} P_j(\cos \theta) \left[2\tilde{G}_q \partial_\tau^2 \tilde{G}_q - 2(\partial_\tau \tilde{G}_q)^2 \right], \quad (\text{C.33})$$

where we replaced $\partial_{\tau'} \rightarrow -\partial_\tau$ when acting on \tilde{G}_q . The first term in (C.33) comes from the last line of (C.31). When $q = 0$, we know from (4.33) that

$$K_j^{0,\tau\tau}(\omega) = \frac{j(j+1)}{2} D_j^0(\omega), \quad (\text{C.34})$$

where $D_j^0(\omega)$ was given in (4.28).

Next, we can calculate the difference $K_j^{q,\tau\tau}(\omega) - K_j^{0,\tau\tau}(\omega)$. Integrating by parts the first term in (C.33), we have:

$$\begin{aligned} K_j^{q,\tau\tau}(\omega) - K_j^{0,\tau\tau}(\omega) &= 2 [G^q(x, x) - G^0(x, x)] \\ &+ 2\pi \int d\tau \int d\theta \sin \theta e^{i\omega\tau} P_j(\cos \theta) \left[-4(\partial_\tau \tilde{G}_q)^2 - \omega^2 \tilde{G}_q^2 + 4(\partial_\tau \tilde{G}_0)^2 + \omega^2 \tilde{G}_0^2 \right]. \end{aligned} \quad (\text{C.35})$$

Since from (C.4), $G^q(x, x) - G^0(x, x) = C_q$, we obtain

$$\begin{aligned} K_j^{q,\tau\tau}(\omega) - K_j^{0,\tau\tau}(\omega) &= 2C_q - \omega^2 [D_j^q(\omega) - D_j^0(\omega)] \\ &- 8\pi \int d\tau \int d\theta \sin \theta e^{i\omega\tau} P_j(\cos \theta) \left[(\partial_\tau \tilde{G}_q)^2 - (\partial_\tau \tilde{G}_0)^2 \right], \end{aligned} \quad (\text{C.36})$$

where we also used (C.15).

Setting $C_q = 0$, we can write this expression as

$$K_j^{q,\tau\tau}(\omega) - K_j^{0,\tau\tau}(\omega) = -\omega^2 [D_j^q(\omega) - D_j^0(\omega)] - \text{F.T.} \left[4(\partial_\tau \tilde{G}_q)^2 - 4(\partial_\tau \tilde{G}_0)^2 \right]. \quad (\text{C.37})$$

Expanding at small s and t and keeping only non-analytic terms, we can write

$$\begin{aligned}
\text{F.T.} \left[4(\partial_\tau \tilde{G}_q)^2 - 4(\partial_\tau \tilde{G}_0)^2 \right] &= \frac{1}{8\pi^2} \text{F.T.} \left[-\mu_q^2 \nabla_{S^2} \log X + \mathcal{F}_q^\infty (\partial_\tau^2 - \nabla_{S^2}^2) X \right. \\
&+ \frac{q^2 + 2\mu_q^2}{480} (4\partial_\tau^2 \nabla_{S^2}^2 - \nabla_{S^2}^4) (X^4 \log X) + \frac{\pi B_q}{15} (2\partial_\tau^4 - 7\partial_\tau^2 \nabla_{S^2}^2 + \nabla_{S^2}^4) X^5 \\
&+ \frac{\mathcal{F}_q^\infty}{120} [3\partial_\tau^4 + 5(4\mu_q^2 - 3)\partial_\tau^2 \nabla_{S^2}^2 - 4\mu_q^2 \nabla_{S^2}^2] X^5 \\
&+ \frac{1}{967680} \left[-4(13q^2 + 20\mu_q^2)\partial_\tau^4 \nabla_{S^2}^2 + (64q^2 + 92\mu_q^2)\partial_\tau^2 \nabla_{S^2}^4 \right. \\
&\left. - (10q^2 + 17\mu_q^2)\nabla_{S^2}^6 \right] (X^8 \log X) + \dots \left. \right] \tag{C.38}
\end{aligned}$$

Using (C.20) and the formulas in Appendix C.6 expanded at large ω and j , we obtain

$$\begin{aligned}
K_j^{q,\tau\tau}(\omega) &= K_j^{0,\tau\tau}(\omega) + j(j+1) \left[\frac{\mu_q^2}{4[(j+1/2)^2 + \omega^2]^{3/2}} + \frac{\mathcal{F}_q^\infty}{2\pi} \frac{2(j+\frac{1}{2})^2 - \omega^2}{[(j+\frac{1}{2})^2 + \omega^2]^3} \right. \\
&+ \frac{(\mu_q^2 - 2q^2)(j+\frac{1}{2})^2 + (11\mu_q^2 + 8q^2)\omega^2}{32[(j+\frac{1}{2})^2 + \omega^2]^{7/2}} + 6B_q \frac{4(j+\frac{1}{2})^4 - 27(j+\frac{1}{2})^2\omega^2 + 4\omega^4}{[(j+\frac{1}{2})^2 + \omega^2]^5} \\
&+ \frac{\mathcal{F}_q^\infty}{8\pi} \frac{16(1 - \mu_q^2)(j+\frac{1}{2})^4 + (88\mu_q^2 - 87)(j+\frac{1}{2})^2\omega^2 - (1 + 16\mu_q^2)\omega^4}{[(j+\frac{1}{2})^2 + \omega^2]^5} \\
&+ \frac{(68q^2 + 15\mu_q^2)(j+\frac{1}{2})^4 - 2(478q^2 + 111\mu_q^2)(j+\frac{1}{2})^2\omega^2 + (824q^2 + 519\mu_q^2)\omega^4}{512[(j+\frac{1}{2})^2 + \omega^2]^{11/2}} \\
&\left. + O\left(\frac{1}{[(j+\frac{1}{2})^2 + \omega^2]^4}\right) \right]. \tag{C.39}
\end{aligned}$$

Next, we should consider $K_j^{q,BB}(\omega)$, which can be computed as

$$K_j^{q,BB}(\omega) = 2G^q(x, x) + \frac{4\pi}{2j+1} \int d\tau d\theta d\phi \sin\theta \lim_{x' \rightarrow 0} \left[e^{i\omega(\tau-\tau')} \sum_{m=-j}^j \mathcal{Y}_{jm}^i(\hat{x}) K_{q,ii'}(x, x') \mathcal{Y}_{jm}^{i'*}(\hat{x}') \right], \tag{C.40}$$

where the first term comes from the contact term in (C.31). Plugging things in, we obtain

$$K_j^{q,BB}(\omega) = 2G^q(x, x) + \frac{4\pi}{j(j+1)} \int d\tau d\theta e^{i\omega\tau} \left[P_j^1(\cos\theta) \left((\partial_\theta \tilde{G}_q)^2 - \tilde{G}_q \partial_\theta^2 \tilde{G}_q \right) - \frac{dP_j^1(\cos\theta)}{d\theta} \tilde{G}_q \left(\partial_\theta \tilde{G}_q + 8q^2 \csc\theta \sin^4 \frac{\theta}{2} \tilde{G}_q \right) \right]. \quad (\text{C.41})$$

We know from (4.33) that when $q = 0$,

$$K_j^{0,BB}(\omega) = \frac{\omega^2 + j^2}{2} D_{j-1}^0(\omega), \quad (\text{C.42})$$

so we can calculate $K_j^{q,BB}(\omega) - K_j^{0,BB}(\omega)$. Using (C.26) and integrating by parts the first term on the second line of (C.41), we obtain

$$K_j^{q,BB}(\omega) - K_j^{0,BB}(\omega) = 2C_q + \frac{8\pi}{j(j+1)} \int d\tau d\theta e^{i\omega\tau} \left[\frac{dP_j(\cos\theta)}{d\theta} (\partial_\theta \tilde{G}_q)^2 - \frac{d^2 P_j(\cos\theta)}{d\theta^2} 4q^2 \csc\theta \sin^4 \frac{\theta}{2} \tilde{G}_q^2 - \frac{dP_j(\cos\theta)}{d\theta} (\partial_\theta \tilde{G}_0)^2 \right]. \quad (\text{C.43})$$

After setting $C_q = 0$ and integrating by parts once the first and third terms under the integral sign and twice the second term, we can write

$$\begin{aligned} K_j^{q,BB}(\omega) &= K_j^{0,BB}(\omega) + \frac{1}{8\pi j(j+1)} \text{F.T.} \left[-\mu_q^2 \nabla_{S^2}^2 \frac{1}{X^2} + \mathcal{F}_q^\infty \nabla_{S^2}^2 \frac{1}{X} \right. \\ &\quad + \frac{1}{24} [4(q^2 + \mu_q^2) \partial_\tau^2 \nabla_{S^2}^2 + (2\mu_q^2 - 3q^2) \nabla_{S^2}^4] (X^2 \log X) \\ &\quad + \frac{\pi B}{2} [4\partial_\tau^2 \nabla_{S^2}^2 - \nabla_{S^2}^4] X^3 + \frac{\mathcal{F}_q^\infty}{96} [(3 - 16\mu_q^2) \partial_\tau^2 \nabla_{S^2}^2 + (8\mu_q^2 - 3) \nabla_{S^2}^4] X^3 \\ &\quad + \frac{1}{40320} \left[(17\mu_q^2 - 30q^2) \nabla_{S^2}^6 + 2(8\mu_q^2 + 69q^2) \partial_\tau^2 \nabla_{S^2}^4 \right. \\ &\quad \left. - 4(16\mu_q^2 + 21q^2) \partial_\tau^4 \nabla_{S^2}^2 \right] (X^6 \log X) + \dots \left. \right] \end{aligned} \quad (\text{C.44})$$

Using again (C.20) and the formulas in Appendix C.6 expanded at large ω and j , we obtain

$$\begin{aligned}
K_j^{q,BB}(\omega) &= K_j^{0,BB}(\omega) + \frac{\mu_q^2}{4\sqrt{(j+\frac{1}{2})^2 + \omega^2}} - \frac{\mathcal{F}_q^\infty}{2\pi [(j+\frac{1}{2})^2 + \omega^2]} \\
&+ \frac{3(\mu_q^2 - 2q^2)(j+\frac{1}{2})^2 + (9\mu_q^2 + 8q^2)\omega^2}{32 [(j+\frac{1}{2})^2 + \omega^2]^{5/2}} \\
&+ \frac{\mathcal{F}_q^\infty (8\mu_q^2 - 3)(j+\frac{1}{2})^2 + (3 - 16\mu_q^2)\omega^2}{8\pi [(j+\frac{1}{2})^2 + \omega^2]^3} + 6B_q \frac{(j+\frac{1}{2})^2 - 4\omega^2}{[(j+\frac{1}{2})^2 + \omega^2]^3} \\
&+ \frac{3(28q^2 + \mu_q^2)(j+\frac{1}{2})^4 - 10(7\mu_q^2 + 118q^2)(j+\frac{1}{2})^2\omega^2 + (808q^2 + 459\mu_q^2)\omega^4}{512 [(j+\frac{1}{2})^2 + \omega^2]^{9/2}} \\
&+ O\left(\frac{1}{[(j+\frac{1}{2})^2 + \omega^2]^3}\right).
\end{aligned} \tag{C.45}$$

C.5 Ultraviolet expansion in the \mathbb{CP}^{N-1} model

The ground state energy in the \mathbb{CP}^{N-1} model can be written as

$$\delta\mathcal{F}_q = \frac{1}{2} \int \frac{d\omega}{2\pi} \sum_{j=0}^{\infty} (2j+1) L_j^q(\omega), \tag{C.46}$$

where we defined

$$L_j^q(\omega) \equiv \log \frac{K_j^{q,\tau\tau}(\omega) \left[D_j^q(\omega) K_j^{q,BB}(\omega) - \left| F_j^{q,B}(\omega) \right|^2 \right]}{D_j^0(\omega) K_j^{0,\tau\tau}(\omega) K_j^{0,BB}(\omega)} \tag{C.47}$$

when $j \geq 1$, and

$$L_0^q(\omega) \equiv \log \frac{D_0^q(\omega)}{D_0^0(\omega)} \tag{C.48}$$

when $j = 0$. The large j and ω behavior of L_j^q can be easily determined from (C.21), (C.30), (C.39), and (C.45) as well as the asymptotic expansion of $D_j^0(\omega)$ in (C.22). We find that for

$j > 0$ we have

$$\begin{aligned}
L_j^q(\omega) &= \frac{8\mu_q^2}{(j + \frac{1}{2})^2 + \omega^2} + \frac{12\mathcal{F}_q^\infty}{\pi} \frac{(j + \frac{1}{2})^2 - \omega^2}{[(j + \frac{1}{2})^2 + \omega^2]^{5/2}} \\
&\quad - \frac{2(3q^2 + \mu_q^2(8\mu_q^2 - 1))(j + \frac{1}{2})^2 + (\mu_q^2(8\mu_q^2 - 5) - 4q^2)\omega^2}{[(j + \frac{1}{2})^2 + \omega^2]^3} \\
&\quad + 144B_q \frac{3(j + \frac{1}{2})^4 - 24(j + \frac{1}{2})^2\omega^2 + 8\omega^4}{[(j + \frac{1}{2})^2 + \omega^2]^{9/2}} \\
&\quad + \frac{3\mathcal{F}_q^\infty}{2\pi} \frac{(25 - 48\mu_q^2)(j + \frac{1}{2})^4 + 3(64\mu_q^2 - 55)(j + \frac{1}{2})^2\omega^2 + 20\omega^4}{[(j + \frac{1}{2})^2 + \omega^2]^{9/2}} \\
&\quad + O\left(\frac{1}{[(j + \frac{1}{2})^2 + \omega^2]^3}\right).
\end{aligned} \tag{C.49}$$

(From the results of this Appendix, one can construct an asymptotic expansion of $L_j^q(\omega)$ that is accurate up to terms that behave as $O(1/[(j + \frac{1}{2})^2 + \omega^2]^{7/2})$. The derivation of the terms not included in (C.49) is straightforward.) The first term in (C.49) yields a linear UV divergence. We explain how to regularize this UV divergence in Section 4.5.

When $j = 0$, the analog of (C.49) can be obtained from (C.21) alone:

$$L_0^q(\omega) = -\frac{8\mathcal{F}_q^\infty}{\pi\omega^3} + \frac{32(\mathcal{F}_q^\infty + 12\pi B_q)}{\pi\omega^5} - \frac{32(\mathcal{F}_q^\infty)^2}{\pi^2\omega^6} + O(1/\omega^7). \tag{C.50}$$

C.6 Fourier transforms on $S^2 \times \mathbb{R}$

Here we present some of the Fourier transforms needed in the previous parts of this Appendix. Recall that the definition of a Fourier transform of a function of θ and τ on $S^2 \times \mathbb{R}$ is

$$\text{F.T. } [f] \equiv 2\pi \int d\tau d\theta \sin\theta e^{i\omega\tau} P_j(\cos\theta) f(\theta, \tau). \tag{C.51}$$

We can calculate explicitly the Fourier transform of $1/X^{2\Delta}$, where $X = \sqrt{2(\cosh\tau - \cos\theta)}$ was defined in (C.17). The calculation proceeds by expanding $1/X^{2\Delta}$ at large τ and per-

forming the integrals term by term:

$$\begin{aligned}
\text{F.T.} \left[\frac{1}{X^{2\Delta}} \right] &= 2\pi \int d\tau d\theta \sin \theta e^{i\omega\tau} P_j(\cos \theta) \frac{1}{[2(\cosh \tau - \cos \theta)]^\Delta} \\
&= 2\pi \sum_{n=0}^{\infty} \int d\tau \frac{(\Delta + j)_n (\Delta)_j (\Delta - 1/2)_n}{(n + 3/2)_j (2n + 1)!} 2^{2n+1} e^{-(\Delta + j + 2n)|\tau| + i\omega\tau} \\
&= 2\pi \sum_{n=0}^{\infty} \frac{(\Delta + j)_n (\Delta)_j (\Delta - 1/2)_n}{(n + 3/2)_j (2n + 1)!} 2^{2n+2} \frac{\Delta + j + 2n}{(\Delta + j + 2n)^2 + \omega^2},
\end{aligned} \tag{C.52}$$

where $(a)_n$ denotes the Pochhammer symbol. The sum in the last expression can be performed analytically in terms of hypergeometric functions, but we will not find it helpful to do so here.

When $\Delta = \frac{1}{2} - m$ with $m \geq 0$ an integer, the infinite sum in (C.52) becomes finite because only the first $m + 1$ terms contribute. We have

$$\begin{aligned}
\text{F.T.} \left[\frac{1}{X} \right] &= \frac{4\pi}{(j + \frac{1}{2})^2 + \omega^2}, \\
\text{F.T.} [X] &= -\frac{8\pi}{[(j + \frac{3}{2})^2 + \omega^2] [(j - \frac{1}{2})^2 + \omega^2]}, \\
\text{F.T.} [X^3] &= \frac{96\pi}{[(j + \frac{5}{2})^2 + \omega^2] [(j + \frac{1}{2})^2 + \omega^2] [(j - \frac{3}{2})^2 + \omega^2]},
\end{aligned} \tag{C.53}$$

and so on. The general formula is

$$\text{F.T.} [X^{-1+2m}] = \frac{(-1)^m (2m)! (4\pi)}{\prod_{k=0}^m [(j + \frac{1}{2} - m + 2k)^2 + \omega^2]}. \tag{C.54}$$

As can be deduced from Section 4.3.1, we have

$$\text{F.T.} \left[\frac{1}{X^2} \right] = 16\pi^2 D_j^0(\omega). \tag{C.55}$$

By taking derivatives of (C.52) with respect to Δ and evaluating the resulting expression at

$\Delta = -2m$, where $m \geq 0$ is an integer, we can also calculate

$$\begin{aligned}
\text{F.T. } [\log X] &= -\frac{16\pi^2}{\omega^2 + j^2} D_{j+1}^0(\omega), \\
\text{F.T. } [X^2 \log X] &= \frac{16\pi^2 \times 6}{[\omega^2 + (j-1)^2][\omega^2 + (j+1)^2]} D_{j+2}^0(\omega), \\
\text{F.T. } [X^4 \log X] &= -\frac{16\pi^2 \times 120}{[\omega^2 + (j-2)^2][\omega^2 + j^2][\omega^2 + (j+2)^2]} D_{j+3}^0(\omega), \\
\text{F.T. } [X^6 \log X] &= \frac{16\pi^2 \times 5040}{\prod_{k=0}^3 [\omega^2 + (j-3+2k)^2]} D_{j+4}^0(\omega), \\
\text{F.T. } [X^8 \log X] &= -\frac{16\pi^2 \times 362880}{\prod_{k=0}^4 [\omega^2 + (j-4+2k)^2]} D_{j+5}^0(\omega),
\end{aligned} \tag{C.56}$$

and so on.

References

- [1] A. M. Polyakov, “Compact gauge fields and the infrared catastrophe,” *Phys.Lett.* **B59** (1975) 82–84.
- [2] X.-G. Wen and Y.-S. Wu, “Transitions between the quantum Hall states and insulators induced by periodic potentials,” *Phys.Rev.Lett.* **70** (1993) 1501–1504.
- [3] W. Chen, M. P. Fisher, and Y.-S. Wu, “Mott transition in an anyon gas,” *Phys.Rev.* **B48** (1993) 13749–13761, [cond-mat/9301037](#).
- [4] S. Sachdev, “Non-zero temperature transport near fractional quantum Hall critical points,” *Phys.Rev.* **B57** (1998) 7157.
- [5] W. Rantner and X.-G. Wen, “Electron spectral function and algebraic spin liquid for the normal state of underdoped high T_c superconductors,” *Phys.Rev.Lett.* **86** (2001) 3871.
- [6] W. Rantner and X.-G. Wen, “Spin correlations in the algebraic spin liquid: Implications for high- T_c superconductors,” *Phys.Rev.* **B66** (2002) 144501.
- [7] O. I. Motrunich and A. Vishwanath, “Emergent photons and new transitions in the O(3) sigma model with hedgehog suppression,” *Phys.Rev.* **B70** (2004) 075104, [cond-mat/0311222](#).

- [8] T. Senthil, A. Vishwanath, L. Balents, S. Sachdev, and M. P. A. Fisher, “Deconfined Quantum Critical Points,” *Science* **303** (Mar., 2004) 1490–1494, [arXiv:cond-mat/0311326](#).
- [9] T. Senthil, L. Balents, S. Sachdev, A. Vishwanath, and M. P. A. Fisher, “Quantum criticality beyond the Landau-Ginzburg-Wilson paradigm,” *Phys.Rev.* **70** (Oct., 2004) 144407, [arXiv:cond-mat/0312617](#).
- [10] M. Hermele, T. Senthil, M. P. A. Fisher, P. A. Lee, N. Nagaosa, and X.-G. Wen, “Stability of U(1) spin liquids in two dimensions,” *Phys.Rev.* **B70** (2004) 214437, [arXiv:cond-mat/0404751](#).
- [11] M. Hermele, T. Senthil, and M. P. Fisher, “Algebraic spin liquid as the mother of many competing orders,” *Phys.Rev.* **B72** (2005) 104404.
- [12] Y. Ran and X.-G. Wen, “Continuous quantum phase transitions beyond Landau’s paradigm in a large- N spin model,” [cond-mat/0609620](#).
- [13] R. K. Kaul, Y. B. Kim, S. Sachdev, and T. Senthil, “Algebraic charge liquids,” *Nature Physics* **4** (2008) 28–31.
- [14] R. K. Kaul and S. Sachdev, “Quantum criticality of U(1) gauge theories with fermionic and bosonic matter in two spatial dimensions,” *Phys.Rev.* **B77** (2008) 155105, [0801.0723](#).
- [15] S. Sachdev, “The landscape of the Hubbard model,” [1012.0299](#).
- [16] T. Appelquist and R. D. Pisarski, “High-Temperature Yang-Mills Theories and Three-Dimensional Quantum Chromodynamics,” *Phys.Rev.* **D23** (1981) 2305.
- [17] T. W. Appelquist, M. J. Bowick, D. Karabali, and L. Wijewardhana, “Spontaneous Chiral Symmetry Breaking in Three-Dimensional QED,” *Phys.Rev.* **D33** (1986) 3704.
- [18] T. Appelquist, D. Nash, and L. Wijewardhana, “Critical behavior in $(2 + 1)$ -dimensional QED,” *Phys.Rev.Lett.* **60** (1988) 2575.
- [19] T. Appelquist and U. W. Heinz, “Three-dimensional $O(N)$ theories at large distances,” *Phys.Rev.* **D24** (1981) 2169.
- [20] S. Coleman, *Aspects of symmetry: selected Erice lectures*. Cambridge University Press, 1988.

- [21] N. Read and S. Sachdev, “Spin-Peierls, valence-bond solid, and Neel ground states of low-dimensional quantum antiferromagnets,” *Phys.Rev.* **B42** (1990) 4568–4589.
- [22] N. Read and S. Sachdev, “Valence-bond and spin-Peierls ground states of low-dimensional quantum antiferromagnets,” *Phys.Rev.Lett.* **62** (1989) 1694–1697.
- [23] G. Murthy and S. Sachdev, “Action of hedgehog instantons in the disordered phase of the $(2 + 1)$ -dimensional $\mathbb{C}\mathbb{P}^{N-1}$ model,” *Nucl.Phys.* **B344** (1990) 557–595.
- [24] J. Lou, A. W. Sandvik, and N. Kawashima, “Antiferromagnetic to valence-bond-solid transitions in two-dimensional $SU(N)$ Heisenberg models with multispin interactions,” *Physical Review B* **80** (Nov., 2009) 180414, 0908.0740.
- [25] R. K. Kaul and A. W. Sandvik, “Lattice Model for the $SU(N)$ Néel to Valence-Bond Solid Quantum Phase Transition at Large N ,” *Physical Review Letters* **108** (Mar., 2012) 137201, 1110.4130.
- [26] S. Pujari, K. Damle, and F. Alet, “Néel-State to Valence-Bond-Solid Transition on the Honeycomb Lattice: Evidence for Deconfined Criticality,” *Physical Review Letters* **111** (Aug., 2013) 087203, 1302.1408.
- [27] M. S. Block, R. G. Melko, and R. K. Kaul, “Fate of $\mathbb{C}\mathbb{P}^{N-1}$ Fixed Points with q Monopoles,” *Physical Review Letters* **111** (Sept., 2013) 137202, 1307.0519.
- [28] R. K. Kaul and M. Block, “Numerical studies of various Neel-VBS transitions in $SU(N)$ antiferromagnets,” 1502.05128.
- [29] V. Borokhov, A. Kapustin, and X.-k. Wu, “Topological disorder operators in three-dimensional conformal field theory,” *JHEP* **0211** (2002) 049, hep-th/0206054.
- [30] V. Borokhov, A. Kapustin, and X.-k. Wu, “Monopole operators and mirror symmetry in three dimensions,” *JHEP* **0212** (2002) 044, hep-th/0207074.
- [31] M. A. Metlitski, M. Hermele, T. Senthil, and M. P. Fisher, “Monopoles in $\mathbb{C}\mathbb{P}^{N-1}$ model via the state-operator correspondence,” *Phys.Rev.* **B78** (2008) 214418, 0809.2816.
- [32] S. S. Pufu, “Anomalous dimensions of monopole operators in three-dimensional quantum electrodynamics,” *Phys.Rev.* **D89** (2014), no. 6 065016, 1303.6125.

- [33] E. Dyer, M. Mezei, and S. S. Pufu, “Monopole Taxonomy in Three-Dimensional Conformal Field Theories,” 1309.1160.
- [34] E. Dyer, M. Mezei, S. Pufu, and S. Sachdev, to appear.
- [35] S. S. Pufu and S. Sachdev, “Monopoles in 2 + 1-dimensional conformal field theories with global U(1) symmetry,” *JHEP* **1309** (2013) 127, 1303.3006.
- [36] S. Sachdev, “Compressible quantum phases from conformal field theories in 2+1 dimensions,” *Phys.Rev.* **D86** (2012) 126003, 1209.1637.
- [37] N. Iqbal, “Monopole correlations in holographically flavored liquids,” 1409.5467.
- [38] V. Borokhov, “Monopole operators in three-dimensional $\mathcal{N} = 4$ SYM and mirror symmetry,” *JHEP* **0403** (2004) 008, hep-th/0310254.
- [39] M. K. Benna, I. R. Klebanov, and T. Klose, “Charges of Monopole Operators in Chern-Simons Yang-Mills Theory,” *JHEP* **1001** (2010) 110, 0906.3008.
- [40] A. Gustavsson and S.-J. Rey, “Enhanced $\mathcal{N} = 8$ Supersymmetry of ABJM Theory on \mathbb{R}^8 and $\mathbb{R}^8/\mathbb{Z}_2$,” 0906.3568.
- [41] F. Benini, C. Closset, and S. Cremonesi, “Chiral flavors and M2-branes at toric CY4 singularities,” *JHEP* **1002** (2010) 036, 0911.4127.
- [42] F. Benini, C. Closset, and S. Cremonesi, “Quantum moduli space of Chern-Simons quivers, wrapped D6-branes and AdS4/CFT3,” *JHEP* **1109** (2011) 005, 1105.2299.
- [43] O. Aharony, P. Narayan, and T. Sharma, “On monopole operators in supersymmetric Chern-Simons-matter theories,” 1502.00945.
- [44] T. T. Wu and C. N. Yang, “Dirac Monopole Without Strings: Monopole Harmonics,” *Nucl.Phys.* **B107** (1976) 365.
- [45] T. T. Wu and C. N. Yang, “Some properties of monopole harmonics,” *Phys.Rev.* **D16** (1977) 1018–1021.
- [46] H.-C. Kim and S. Kim, “Semi-classical monopole operators in Chern-Simons-matter theories,” 1007.4560.
- [47] I. A. Shovkovy, “Magnetic Catalysis: A Review,” *Lect.Notes Phys.* **871** (2013) 13–49, 1207.5081.

- [48] J. Zinn-Justin, *Quantum Field Theory and Critical Phenomena*. International series of monographs on physics. Clarendon Press, 2002.
- [49] R. Rattazzi, V. S. Rychkov, E. Tonni, and A. Vichi, “Bounding scalar operator dimensions in 4D CFT,” *JHEP* **0812** (2008) 031, 0807.0004.
- [50] D. Simmons-Duffin, “A Semidefinite Program Solver for the Conformal Bootstrap,” 1502.02033.