



Asymptotic symmetries of Yang-Mills theory

Citation

Strominger, Andrew. 2014. "Asymptotic Symmetries of Yang-Mills Theory." J. High Energ. Phys. 2014 (7) (July). doi:10.1007/jhep07(2014)151.

Published version

[https://doi.org/10.1007/JHEP07\(2014\)151](https://doi.org/10.1007/JHEP07(2014)151)

Link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:29374081>

Terms of use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Open Access Policy Articles (OAP), as set forth at

<https://harvardwiki.atlassian.net/wiki/external/NGY5NDE4ZjgzNTc5NDQzMGIzZWZhMGFIOWI2M2EwYTg>

Accessibility

<https://accessibility.huit.harvard.edu/digital-accessibility-policy>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#)

ASYMPTOTIC SYMMETRIES OF YANG-MILLS THEORY

Andrew Strominger

*Radcliffe Institute for Advanced Study, Harvard University,
Cambridge, MA 02138, USA*

Abstract

Asymptotic symmetries at future null infinity (\mathcal{I}^+) of Minkowski space for electrodynamics with massless charged fields, as well as non-Abelian gauge theories with gauge group G , are considered at the semiclassical level. The possibility of charge/color flux through \mathcal{I}^+ suggests the symmetry group is infinite-dimensional. It is conjectured that the symmetries include a G Kac-Moody symmetry whose generators are "large" gauge transformations which approach locally holomorphic functions on the conformal two-sphere at \mathcal{I}^+ and are invariant under null translations. The Kac-Moody currents are constructed from the gauge field at the future boundary of \mathcal{I}^+ . The current Ward identities include Weinberg's soft photon theorem and its colored extension.

Contents

1	Introduction	1
2	Electrodynamics	4
2.1	The boundary data problem	6
2.2	Conformal transformations	9
2.3	The boundary current	10
2.4	Ward identities and Weinberg's theorem	11
2.5	Green functions	14
2.6	Kac-Moody level?	15
3	The non-Abelian case	16
4	Stringy realization?	17

1 Introduction

Over half a century ago, Bondi, van der Burg, Metzner and Sachs [1] argued that gravitational wave scattering in asymptotically Minkowskian spacetimes is governed by an infinite-dimensional symmetry algebra now known as the BMS algebra. The symmetry has consequences for any theory which can be coupled to gravity, and would be expected to play a central role in Minkowskian scattering theory. However, while some important results have been obtained, the full import of BMS symmetry has remained elusive.

The BMS generators are infinitesimal diffeomorphisms which preserve a prescribed structure at future null infinity, or \mathcal{I}^+ , but act nontrivially on the physical data residing there. \mathcal{I}^+ is the product of a conformal two-sphere (directions of outgoing null rays) with a null line (retarded times). The existence of infinitely many conformal two-spheres in \mathcal{I}^+ leads to both an infinity of charges - e.g. the Bondi masses for each S^2 - and an infinite-dimensional symmetry group. The BMS algebra has two types of elements. First there are the global $SL(2, C)$ conformal transformations of the S^2 which are related to Lorentz transformations. Second there are "supertranslations" along the the null generators of \mathcal{I}^+ . These are infinite in number because their magnitude can vary over the S^2 .

In a fascinating recent proposal, Barnich and Troessaert [2] and Banks [3] argued that the BMS algebra is, after all these years, *not* in fact the complete symmetry algebra of asymptotically Minkowskian spacetimes. In the original analysis of [1], it was required that

all diffeomorphisms be everywhere finite on \mathcal{I}^+ . In [2, 3] it was argued that this should be relaxed to allow for certain analytic singularities, or equivalently have holomorphic behavior on local patches of the conformal S^2 . This relaxation enhances the *global* $SL(2, C)$ subalgebra of BMS to two *local* Virasoro algebras, leading to a new “extended BMS symmetry”.

Two decades after BMS, BPZ [4] studied a mathematically similar problem: 2D field theories with a global $SL(2, C)$ symmetry. Unlike BMS, BPZ allowed symmetries with analytic singularities, which leads to two Virasoro algebras. The implications of this infinite symmetry algebra for 2D field theory are of course truly extraordinary. One hopes by analogy that the extended BMS symmetry has extraordinary implications for Minkowski scattering. It is plausible, as we explore in [5], that some of these implications are already known in the guise of Weinberg’s soft graviton theorem [6].

This paper will address a simpler, but very closely related, toy version of the gravitational problem. In electrodynamics with no massless charged particles, the symmetry group at both spatial infinity and \mathcal{I}^+ is a global $U(1)$. The $U(1)$ symmetry is generated by the associated conserved electric charge, which may be measured at either spatial infinity or \mathcal{I}^+ .

The situation is radically different if there are massless charged particles. Then there can be charge flux through \mathcal{I}^+ , and many different electric charges associated to the many asymptotic S^2 s. Evidently, the \mathcal{I}^+ symmetry is not a simple global $U(1)$. This raises our basic question:

What are the asymptotic symmetries at \mathcal{I}^+ of electrodynamics with massless charged particles?

This is a toy version of the question asked by BMS. It is of interest in its own right as well as for a warm-up to the gravitational case. Part of this problem is defining both what is meant by asymptotic symmetries and how they act.

We conjecture herein that these symmetries include¹ “large” $U(1)$ gauge transformations which approach (the real part of) a holomorphic function on local patches of the conformal S^2 on \mathcal{I}^+ and are constant along the null generators. These are the toy analogs of the singular conformal transformations of [2], and appear related to the Sky transformations of [7]. A current J_z tangent to the conformal S^2 at the future of \mathcal{I}^+ is constructed in radiation

¹ The conjecture does not preclude the possibility that the full asymptotic symmetry group is larger. In particular, one might consider, in analogy with the BMS supertranslations, gauge parameters which locally approach arbitrary (not necessarily holomorphic) functions at infinity. Such transformations would not show up in the analysis of this paper because they are ruled out by our gauge choice.

gauge from the asymptotic gauge field. Insertions of contour integrals of this current are then shown to generate infinitesimal large gauge transformations on the matter fields and obey a $U(1)$ Kac-Moody algebra. Moreover the large gauge transformation law of the current itself hints at a nonzero Kac-Moody level proportional to $\frac{1}{e^2}$.² While intriguing, a precise statement about a possible level must await a better understanding of the putative CFT_2 at \mathcal{I}^+ : for example determining the sign (or phase) requires an as-yet-to-be-discussed notion of adjoint. An asymptotic Kac-Moody symmetry fits well with the proposed extended BMS group, since the generators are ready to transform under the Virasoro generators which appear when gravity is coupled.

Much of this structure is in fact already known in a different language. Around the same time as BMS, Weinberg [6] derived soft-photon theorems which give relations among scattering amplitudes with and without soft photons. In a key motivation for the present work, Maldacena and Zhiboedov have shown [8] by transforming Weinberg's momentum-space formulae into position space coordinates on \mathcal{I}^+ , that the soft photon theorem can be rewritten in the form of a current algebra Ward identity. We will see that the relations among scattering amplitudes implied by Weinberg's theorem can be viewed as a consequence of the large gauge symmetry.

Weinberg does not encounter a non-zero level when considering multiple soft photon insertions. This is because his soft photon insertions are subtly different from our J_z : they are gauge invariant but nonlocal while J_z is local but not gauge invariant (see section 2.3).

The situation for non-Abelian gauge theories (in the Coulomb phase) is similar. In general color flux can pass through \mathcal{I}^+ , leading to the basic question

What are the asymptotic symmetries at \mathcal{I}^+ for non-Abelian gauge theories?

The analysis is very similar to the abelian case with charged massless particles: for simple gauge group G we find a G Kac-Moody algebra and again hints of a non zero level proportional to $\frac{1}{g_{\text{YM}}^2}$.

The analysis of this paper is restricted to the simplest nontrivial situations, as these already turn out to be quite subtle and rich. For example we work only at the semiclassical level and restrict to configurations which have no incoming charge flux on \mathcal{I}^- and decay to the vacuum in the far future.

This paper is organized as follows. Section 2 treats the $U(1)$ case. In 2.1 we describe the boundary conditions and falloffs at \mathcal{I}^+ in radiation gauge, and develop the asymptotic

²With a current normalization fixed by the periodicity of the gauge transformations.

expansion in the inverse radius. Attention is drawn to residual large gauge transformations which are locally holomorphic functions on S^2 . 2.2 briefly describes the conformal transformation properties of the asymptotic data. In 2.3 we define the Kac-Moody current J_z in terms of tangential components of the gauge field at the infinite future of \mathcal{I}^+ , denoted \mathcal{I}_+^+ . J_z is shown semiclassically to acquire poles where charged particles cross \mathcal{I}^+ , as appropriate to such a current. In 2.4 we recast the appearance of poles in J_z as a semiclassical Ward identity and discuss the relation to Weinberg's soft photon theorem. It is further shown that insertion of J_z contour integrals generate large gauge transformations of matter fields on local patches of the S^2 at \mathcal{I}^+ . 2.5 reviews the radiation-gauge Feynman two-point function for gauge field in our notation and computes a boundary gauge field correlator. 2.6 infers a nonzero Kac-Moody level from the coefficient of the gauge variation of J_z itself. The non-abelian case is very similar and sketched in section 3. Finally in section 4 we discuss the realization of this structure within string theory and speculate that, in the spirit of [12], the spacetime Kac-Moody current J_z is the lift of the familiar worldsheet Kac-Moody current associated to spacetime gauge symmetries.

2 Electrodynamics

In this section we consider a $U(1)$ gauge theory coupled to massless charged matter. The Maxwell electromagnetic action is

$$S_M = -\frac{1}{4e^2} \int d^4x \sqrt{-g} F_{\mu\nu} F^{\mu\nu}. \quad (2.1)$$

This has the gauge symmetry

$$\delta_{\hat{\varepsilon}} A_\mu = \partial_\mu \hat{\varepsilon} \quad (2.2)$$

with the periodicity

$$\hat{\varepsilon} \sim \hat{\varepsilon} + 2\pi. \quad (2.3)$$

Given a surface Σ in \mathcal{I}^+ , electromagnetic Bondi charges can be defined by³

$$Q^E = \frac{1}{e^2} \int_\Sigma *F, \quad Q^M = \frac{1}{2\pi} \int_\Sigma F. \quad (2.4)$$

We consider the case $Q^M = 0$ for all Σ . In a theory with massless electrically charged particles which reach \mathcal{I}^+ , Q^E will depend on the choice of Σ . We will work with the flat

³ Q^M is an integer with the given $\hat{\varepsilon}$ -periodicity. Q^E differs by a factor of e from some definitions of the electric charge, and is also set to an integer by Dirac quantization.

Minkowski metric in coordinates

$$ds_F^2 = -dv^2 - 2drdv + 2r^2\gamma_{z\bar{z}}dzd\bar{z}, \quad \int d^2z\gamma_{z\bar{z}} = 4\pi. \quad (2.5)$$

(v, z, \bar{z}) with $v = t - r$, are coordinates on \mathcal{I}^+ . For the special case $z = e^{i\phi} \tan \frac{\theta}{2}$, $\gamma_{z\bar{z}} = 2(1 + z\bar{z})^{-2}$. The nonzero connection coefficients are

$$\Gamma_{rz}^z = \frac{1}{r}, \quad \Gamma_{zz}^z = \partial_z \ln \gamma_{z\bar{z}}, \quad \Gamma_{z\bar{z}}^v = r\gamma_{z\bar{z}}, \quad \Gamma_{z\bar{z}}^r = -r\gamma_{z\bar{z}}. \quad (2.6)$$

Denoting the charges and surfaces at retarded time v by $Q^E(v)$ and Σ_v one has

$$Q^E(v) = \frac{1}{e^2} \int_{\Sigma_v} d^2z\gamma_{z\bar{z}}r^2F_{rv}. \quad (2.7)$$

For nonzero matter charge current j_ν^M the Maxwell equations $\nabla^\mu F_{\mu\nu} = e^2j_\nu^M$ are

$$-\gamma_{z\bar{z}}r^2\partial_v F_{rv} + \partial_z F_{\bar{z}v} + \partial_{\bar{z}} F_{zv} + \partial_r(\gamma_{z\bar{z}}r^2F_{rv}) = e^2\gamma_{z\bar{z}}r^2j_v^M, \quad (2.8)$$

$$\partial_z F_{\bar{z}r} + \partial_{\bar{z}} F_{zr} + \partial_r(\gamma_{z\bar{z}}r^2F_{rv}) = e^2\gamma_{z\bar{z}}r^2j_r^M, \quad (2.9)$$

$$r^2\partial_r(F_{rz} - F_{vz}) - r^2\partial_v F_{rz} + \partial_z(\gamma^{z\bar{z}}F_{\bar{z}z}) = e^2r^2j_z^M. \quad (2.10)$$

We then have

$$\partial_v Q^E(v) = - \int_{\Sigma_v} d^2z\gamma_{z\bar{z}}r^2j_v^M. \quad (2.11)$$

The causal geometry is depicted in figure 1.

In this paper we impose a number of simplifying restrictions in order to exhibit the basic phenomena in the simplest possible setting. We consider only theories which have no stable massive particles and revert to the vacuum in the far future. In the spirit of [1] we concentrate on the structure of \mathcal{I}^+ and ignore the parallel structure on \mathcal{I}^- , as well as the scattering between the two. We specialize to the case where there is no incoming charges, namely on \mathcal{I}^-

$$j_\mu^M(r, -\infty, z, \bar{z}) = 0, \quad Q^E(-\infty) = 0. \quad (2.12)$$

At the quantum level, this means the incoming state is annihilated by the local charge current. This is of course semiclassically consistent with nontrivial outgoing charge fluxes on \mathcal{I}^+ , subject to the restriction that the total integrated charge flux must vanish because the system reverts to the vacuum. We expect it is both possible and interesting to extend our analysis to more general cases.




Figure 1: Penrose diagram for Minkowski scattering. The figure depicts incoming photons (blue) from \mathcal{I}^- ($v = -\infty$) creating a matter charge current j^M (green) outgoing at \mathcal{I}^+ ($r = \infty$). The total charge $Q^E(v)$ is measured on a conformal sphere at \mathcal{I}^+ (parameterized by (z, \bar{z})), and depends on the retarded time v . The (red) cones are null surfaces of constant v .

2.1 The boundary data problem

In this subsection we consider the inverse of the initial data problem on \mathcal{I}^- , namely the final data problem on \mathcal{I}^+ . It is convenient to impose the temporal gauge condition

$$A_v = 0, \tag{2.13}$$

which becomes $A_t=0$ in standard Minkowskian coordinates. This leaves unfixed a residual v -independent gauge symmetry. We partially fix this by demanding in the far past :⁴

$$r^2\gamma_{z\bar{z}}\nabla^\mu A_\mu(z, \bar{z}, r, -\infty) = [\partial_z A_{\bar{z}} + \partial_{\bar{z}} A_z + (\partial_r - \partial_v)(\gamma_{z\bar{z}}r^2 A_r)]_{\mathcal{I}^-} = 0, \quad (2.14)$$

(2.13)-(2.14) is commonly referred to as radiation gauge. Still remaining are gauge transformations obeying

$$2\partial_z\partial_{\bar{z}}\hat{\varepsilon} + \gamma_{z\bar{z}}\partial_r(r^2\partial_r\hat{\varepsilon}) = 0 = \partial_v\hat{\varepsilon}. \quad (2.15)$$

Locally this allows the residual transformations

$$\begin{aligned} \hat{\varepsilon} &= \varepsilon(z) + \bar{\varepsilon}(\bar{z}), \\ \delta_\varepsilon A_z &= \partial_z \varepsilon. \end{aligned} \quad (2.16)$$

for holomorphic ε . Globally of course there are no holomorphic functions on S^2 except the constant, and (2.16) must have singularities at points. Nevertheless we shall consider below transformations in a patch on \mathcal{I}^+ for which the local solutions (2.16) are relevant. The transformations (2.16) are the electromagnetic analog of the local conformal transformations proposed in [2] as part of the extended BMS group. They create zero-momentum "boundary photons" and we will refer to them as large gauge transformations.

In radiation gauge one has the simple relations

$$F_{vz} = \partial_v A_z, \quad F_{vr} = \partial_v A_r. \quad (2.17)$$

The Maxwell equations may then be written

$$-\partial_v[\partial_z A_{\bar{z}} + \partial_{\bar{z}} A_z + (\partial_r - \partial_v)(\gamma_{z\bar{z}}r^2 A_r)] = e^2\gamma_{z\bar{z}}r^2 j_v^M, \quad (2.18)$$

$$2\partial_z\partial_{\bar{z}}A_r - \partial_v\partial_r(\gamma_{z\bar{z}}r^2 A_r) - \partial_r(\partial_z A_{\bar{z}} + \partial_{\bar{z}} A_z) = e^2\gamma_{z\bar{z}}r^2 j_r^M, \quad (2.19)$$

$$r^2(\partial_r^2 A_z - 2\partial_r\partial_v A_z) - r^2\partial_r\partial_z A_r + r^2\partial_v\partial_z A_r + \partial_z(\gamma^{z\bar{z}}(\partial_{\bar{z}} A_z - \partial_z A_{\bar{z}})) = e^2 r^2 j_z^M. \quad (2.20)$$

A typical wave solution near \mathcal{I}^+ is of the general form

$$A_r = 0, \quad A_z = a(z, \bar{z})e^{ikv}. \quad (2.21)$$

This motivates the asymptotic boundary condition at \mathcal{I}^+

$$A_z \sim O(1), \quad A_r \sim O\left(\frac{1}{r^2}\right). \quad (2.22)$$

⁴This condition is obstructed when there is incoming charge flux on \mathcal{I}^- .

This leading behavior of A_r insures that $Q^E(v)$ is always finite. We may then expand near \mathcal{I}^+

$$\begin{aligned} A_z(r, v, z, \bar{z}) &= \sum_{n=0}^{\infty} \frac{A_z^{(n)}(v, z, \bar{z})}{r^n}, \\ A_r(r, v, z, \bar{z}) &= \sum_{n=0}^{\infty} \frac{A_r^{(n)}(v, z, \bar{z})}{r^{n+2}}. \end{aligned} \quad (2.23)$$

For simplicity we consider the case where $j_r^M = j_z^M = 0$ near \mathcal{I}^+ . Current conservation then implies in this region

$$j_v^M(r, v, z, \bar{z}) = \frac{j_v^{M(0)}(v, z, \bar{z})}{r^2}. \quad (2.24)$$

The leading terms at large r in (2.18)-(2.20) then give the equations on \mathcal{I}^+

$$-\partial_v [\gamma^{z\bar{z}}(\partial_{\bar{z}} A_z^{(0)} + \partial_z A_{\bar{z}}^{(0)}) - \partial_v A_r^{(0)}] = e^2 j_v^{M(0)}, \quad (2.25)$$

$$2\partial_v A_z^{(1)} + \partial_v \partial_z A_r^{(0)} + \partial_z (\gamma^{z\bar{z}}(\partial_{\bar{z}} A_z^{(0)} - \partial_z A_{\bar{z}}^{(0)})) = 0 \quad (2.26)$$

$$2\partial_z \partial_{\bar{z}} A_r^{(0)} + \gamma_{z\bar{z}} \partial_v A_r^{(1)} + \partial_z A_{\bar{z}}^{(1)} + \partial_{\bar{z}} A_z^{(1)} = 0 \quad (2.27)$$

The boundary fields appearing here are constrained by the subleading term in (2.18)

$$-\partial_v [\gamma^{z\bar{z}}(\partial_{\bar{z}} A_z^{(1)} + \partial_z A_{\bar{z}}^{(1)}) - \partial_v A_r^{(1)}] = 0. \quad (2.28)$$

Integrating the constraint equation (2.25) along a null generator of \mathcal{I}^+ and using (2.14) gives the important relation

$$2\partial_{\bar{z}} A_z^{(0)} = -\gamma_{z\bar{z}} e^2 \int_{-\infty}^v dv' j_{v'}^{M(0)} + \gamma_{z\bar{z}} F_{vr}^{(0)} - F_{z\bar{z}}^{(0)}. \quad (2.29)$$

Integrating this once more with respect to v gives $A_r^{(0)}$ in terms of $A_z^{(0)}$. The other equations of motion determine $A_z^{(1)}$ in terms of $A_z^{(0)}$

$$A_z^{(1)} = -\partial_z \int_{-\infty}^v dv' \left[\gamma^{z\bar{z}} \partial_{\bar{z}} A_z^{(0)} + \frac{e^2}{2} \int_{-\infty}^{v'} dv'' j_{v''}^{M(0)} \right] \quad (2.30)$$

Finally integrating (2.28) with respect to v then gives $A_r^{(1)}$ in terms of $A_z^{(0)}$. Further subleading equations of motion similarly determine further subleading terms in the power expansion.

Hence the local free final data lies in the two unconstrained real functions $A_z^{(0)}$. This is as expected, because a massless spin one has two degrees of freedom and should be locally specified by two real (or one complex) functions on \mathcal{I}^+ . Noting that $\partial_v A_z^{(0)} = F_{vz}^{(0)}$, the free data is simply related to the radiation energy flux at \mathcal{I}^+ according to $T_{vv} \sim \frac{1}{r^2} \gamma^{z\bar{z}} F_{vz}^{(0)} F_{v\bar{z}}^{(0)}$.

2.2 Conformal transformations

In this subsection we briefly describe the action of \mathcal{I}^+ conformal transformations - a subset of the extended BMS transformations - on fields appearing in the $\frac{1}{r}$ expansion around \mathcal{I}^+ . Similarities with the familiar AdS case will be evident. The transformations are generated by the real part of the complex vector fields [2]

$$\begin{aligned}
\zeta_{\text{conf}}^a \partial_a &\equiv (\zeta^z - \frac{v}{2r} \gamma^{z\bar{z}} D_{\bar{z}} D_z \zeta^z) \partial_z + (-\frac{r}{2} D_z \zeta^z + \frac{v}{4} D^2 D_z \zeta^z) \partial_r - \frac{v}{2r} \gamma^{z\bar{z}} D_{\bar{z}}^2 \zeta^z \partial_{\bar{z}} + \frac{v}{2} D_z \zeta^z \partial_v \\
&= (1 + \frac{vR}{4r}) \zeta^z \partial_z - (\frac{r}{2} D_z \zeta^z + \frac{v}{4} D_z (R \zeta^z)) \partial_r - \frac{v}{2r} \gamma^{z\bar{z}} D_{\bar{z}}^2 \zeta^z \partial_{\bar{z}} + \frac{v}{2} D_z \zeta^z \partial_v \\
&= (1 + \frac{v}{2r}) \zeta^z \partial_z - (1 + \frac{v}{r}) \frac{D_z \zeta^z r}{2} \partial_r - \frac{v}{2r} \gamma^{z\bar{z}} D_{\bar{z}}^2 \zeta^z \partial_{\bar{z}} + \frac{v}{2} D_z \zeta^z \partial_v,
\end{aligned} \tag{2.31}$$

where $\partial_{\bar{z}} \zeta^z = 0$, D is the γ -covariant derivative with curvature R and $D^2 = \gamma^{z\bar{z}} (D_z D_{\bar{z}} + D_{\bar{z}} D_z)$. In the last line we use the fact that $R = 2$ for our choice of $\gamma_{z\bar{z}}$. These act on the flat Minkowski metric as

$$\mathcal{L}_\zeta ds^2 = -rv D_z^3 \zeta^z dz^2, \tag{2.32}$$

which is a subleading (in $\frac{1}{r}$) contribution to the induced S^2 metric. A massless scalar field has the expansion near \mathcal{I}^+

$$\phi(r, v, z, \bar{z}) = \sum_n \frac{\phi^{(n)}(v, z, \bar{z})}{r^{n+1}}. \tag{2.33}$$

Conformal transformations act on the leading expansion coefficient as

$$\mathcal{L}_\zeta \phi^{(0)}(v, z, \bar{z}) = \frac{1}{2} D_z \zeta^z (1 + v \partial_v) \phi^{(0)}(v, z, \bar{z}) + \zeta^z \partial_z \phi^{(0)}(v, z, \bar{z}). \tag{2.34}$$

Hence the v zero modes of these coefficients transform as two dimensional conformal fields with weights $(h, \bar{h}) = (\frac{1}{2}, \frac{1}{2})$. For the vector field one finds the v component obeys

$$\mathcal{L}_\zeta A_v = \zeta_{\text{conf}}^a \partial_a A_v + \frac{1}{2r} (\zeta^z A_z^{(0)} - \gamma^{z\bar{z}} D_z^2 \zeta^z A_{\bar{z}}^{(0)}) + \dots \tag{2.35}$$

The gauge choice $A_v = 0$ employed in this paper is not preserved by \mathcal{L}_ζ and explicitly breaks the symmetry. Hence, while useful for other purposes, it is not ideal for analyzing conformal properties. More generally the extended BMS symmetry potentially constrains the boundary correlations functions on \mathcal{I}^+ and the S-matrix. Indeed, the peeling theorems of Newman and Penrose [9] exploit the original BMS symmetry to constrain the behavior of gravitational radiation at \mathcal{I}^+ .

2.3 The boundary current

We consider final data which corresponds to solutions with finite total energy as measured at spatial infinity as well as vanishing initial and final charges. This requires that at the future ($v \rightarrow \infty$) of \mathcal{I}^+ (denoted \mathcal{I}_+^+ and depicted in figure 1) the field strengths go to zero:

$$F_{z\bar{z}}^{(0)}(\infty, z, \bar{z}) = 0 = F_{rv}^{(0)}(\infty, z, \bar{z}), \quad (2.36)$$

or equivalently

$$\partial_z A_{\bar{z}}^{(0)}(\infty, z, \bar{z}) = \partial_{\bar{z}} A_z^{(0)}(\infty, z, \bar{z}), \quad \partial_v A_r^{(0)}(\infty, z, \bar{z}) = 0. \quad (2.37)$$

(2.29) then reduces to

$$\partial_{\bar{z}} A_z^{(0)}(\infty, z, \bar{z}) = -\gamma_{z\bar{z}} \frac{e^2}{2} \int_{-\infty}^{\infty} dv' j_{v'}^{M(0)}. \quad (2.38)$$

For such finite-energy data we can define currents on \mathcal{I}_+^+

$$J_z(z, \bar{z}) = -\frac{4\pi}{e^2} A_z^{(0)}(\infty, z, \bar{z}). \quad (2.39)$$

A parallel construction is possible for $J_{\bar{z}}$. This current is sourced by charges which cross \mathcal{I}^+ at any value of retarded time

$$\partial_{\bar{z}} J_z(z, \bar{z}) = 2\pi\gamma_{z\bar{z}} \int_{-\infty}^{\infty} j_{v'}^{M(0)}(v', z, \bar{z}) dv'. \quad (2.40)$$

Consider for example of a set of massless point electric particles crossing \mathcal{I}^+ with the conserved charge current

$$r^2 j_v^M = \sum_k q_k \delta(v - v_k) \frac{\delta^2(z - z_k)}{\gamma_{z\bar{z}}}, \quad (2.41)$$

where

$$\sum_k q_k = 0, \quad (2.42)$$

so that the total initial and final electric charge vanishes. We then have

$$\partial_{\bar{z}} J_z = 2\pi \sum_k q_k \delta^2(z - z_k). \quad (2.43)$$

Using $\partial_{\bar{z}} \frac{1}{z} = 2\pi \delta^2(z)$ this is solved by

$$J_z = \sum_k \frac{q_k}{z - z_k}. \quad (2.44)$$

Locally, a holomorphic function can be added corresponding to large gauge transformations. We will see in the next subsection that the quantum version of this simple equality is the Ward identity for a $U(1)$ current algebra on \mathcal{I}_+^+ .

Note that in radiation gauge the presence of pointlike charged radiation passing through \mathcal{I}^+ implies the existence of singularities in the gauge potential $A_z^{(0)} \rightarrow -\frac{e^2}{4\pi} J_z$ even at \mathcal{I}_+^+ where the system reverts to the vacuum. Smoothing out the charge flux over z will also smooth out $A_z^{(0)}$, but it will necessarily remain nonzero in the far future.

It is interesting to compare the current J_z , viewed as a quantum operator, to the zero momentum limit of an operator annihilating a photon emerging at z on \mathcal{I}^+ . The gauge invariant operator

$$V_{z,\omega}(z, \bar{z}) = \int_{-\infty}^{\infty} dv e^{-i\omega v} F_{v\bar{z}}^{(0)}(v, z, \bar{z}), \quad \omega > 0 \quad (2.45)$$

annihilates a positive helicity photon of energy ω emerging at z while the Fourier transform of $F_{v\bar{z}}^{(0)}$ annihilates negative helicity photons. In the soft photon limit $\omega \rightarrow 0$ (2.45) becomes a total derivative and reduces to

$$V_{z,0}(z, \bar{z}) = A_z^{(0)}(\infty, z, \bar{z}) - A_z^{(0)}(-\infty, z, \bar{z}). \quad (2.46)$$

This is closely related to the current J_z defined in (3.8). Given our boundary conditions including (2.12) both are sourced by charge fields according to (2.44). However there is a difference: $V_{z,0}$ is nonlocal on \mathcal{I}^+ but invariant under large gauge transformations, while J_z is local but transforms nontrivially under large gauge transformations.

2.4 Ward identities and Weinberg's theorem

In this subsection we recast the pole formula (2.44) for the current as a semiclassical relation between correlation functions, identify it as a Kac-Moody Ward identity and relate it to Weinberg's soft photon theorem.

S-matrix elements with no charged particles coming in from \mathcal{I}^- and n charged particles with charges q_k going out at (v_k, z_k, \bar{z}_k) on \mathcal{I}^+ are constructed from correlation functions of the form

$$\langle 0 | V_1(v_1, z_1, \bar{z}_1) \dots V_n(v_n, z_n, \bar{z}_n) \dots | 0 \rangle. \quad (2.47)$$

Here V_k is an operator at $r = \infty$ on \mathcal{I}^+ which annihilates a matter particle of charge q_k at (v_k, z_k, \bar{z}_k) .⁵ As the counterpart of our assumption (2.12) of no charge flux across \mathcal{I}^- ,

⁵ It would be interesting to work out in detail the S-matrix in a basis with particles labelled by their (z, \bar{z}) coordinates on \mathcal{I}^\pm (rather than the usual momentum space labels) and study their conformal transformation properties. This is partially achieved in the twistor formulation.

the incoming particles, denoted by ..., are all taken here to be neutral. Arbitrary time and radius independent gauge transformations $f(z, \bar{z})$ of the matter fields on any fixed time slice with $v + r = t_0$ are generated by the matter part of Noether charge

$$Q_{t_0}^M[f] = \int_{v+r=t_0} d^3\Sigma^\mu f j_\mu^M \quad (2.48)$$

according to

$$[Q_{t_0}^M[f], V_k(r, t_0 - r, z, \bar{z})] = -f(z, \bar{z}) q_k V_k(r, t_0 - r, z, \bar{z}). \quad (2.49)$$

Consider the normal-ordered v -zero mode of the matter charge current on \mathcal{I}^+

$$q_{z\bar{z}}(z, \bar{z}) = \gamma_{z\bar{z}} \int_{-\infty}^{\infty} dv j_v^{0M}(v, z, \bar{z}). \quad (2.50)$$

Since the fixed time surface approaches \mathcal{I}^+ in the limit $t_0 \rightarrow \infty$, $q_{z\bar{z}}$ is the special case of $Q_\infty^M[f]$ where f is set to a delta function. It follows that

$$[q_{z\bar{z}}(z, \bar{z}), V_k(v_k, z_k, \bar{z}_k)] = -q_k V_k(v_k, z_k, \bar{z}_k) \delta^2(z - z_k). \quad (2.51)$$

Using this and $\langle 0|q_{z\bar{z}} = 0$, insertions of $q_{z\bar{z}}$ in correlation functions gives

$$\langle 0|V_1(v_1, z_1, \bar{z}_1) \dots V_n(v_n, z_n, \bar{z}_n) q_{z\bar{z}}(z, \bar{z}) \dots |0\rangle = \sum_{k=1}^n q_k \delta^2(z - z_k) \langle 0|V_1(v_1, z_1, \bar{z}_1) \dots V_n(v_n, z_n, \bar{z}_n) \dots |0\rangle. \quad (2.52)$$

The integrated constraint (2.29) can be written

$$\partial_{\bar{z}} J_z = 2\pi q_{z\bar{z}} + \frac{2\pi}{e^2} F_{z\bar{z}}^{(0)}(\infty, z, \bar{z}) - \frac{2\pi}{e^2} \gamma_{z\bar{z}} F_{vr}^{(0)}(\infty, z, \bar{z}). \quad (2.53)$$

The last two terms are the magnetic and electric field operators in the infinite future at \mathcal{I}_+^+ . If there are no operator insertions at \mathcal{I}_+^+ , and the incoming state has finite extent and energy, these terms vanishes in vacuum correlation functions and can be dropped.⁶ We thereby reproduce the operator version of the semiclassical relation (2.44)

$$\langle 0|V_1(v_1, z_1, \bar{z}_1) \dots V_n(v_n, z_n, \bar{z}_n) J_z(z, \bar{z}) \dots |0\rangle = \sum_{k=1}^n \frac{q_k}{z - z_k} \langle 0|V_1(v_1, z_1, \bar{z}_1) \dots V_n(v_n, z_n, \bar{z}_n) \dots |0\rangle. \quad (2.54)$$

This is not a new result: Weinberg's soft photon theorem [6] for amplitudes reads

$$\langle 0|V_1(p_1) \dots V_n(p_n) V_\gamma(\varepsilon, p_\gamma) \dots |0\rangle_{LSZ} \sim \sum_{k=1}^n \frac{q_k \varepsilon \cdot p_k}{p_\gamma \cdot p_k} \langle 0|V_1(p_1) \dots V_n(p_n) \dots |0\rangle_{LSZ}, \quad p_\gamma \rightarrow 0, \quad (2.55)$$

⁶These terms are relevant for multiple J_z insertions on \mathcal{I}^+ .

where the subscript denotes that correlators are LSZ reduced to amplitudes. Here the n particles are in momentum eigenstates p_k with $p_k^2 = 0$ and V_γ annihilates a soft photon with momentum p_γ and polarization ε . Maldacena and Zhiboedov have shown [8] that, for an appropriate normalization of V_γ , (2.54) is simply a rewriting of (2.55) in position space for the conformal sphere at \mathcal{I}^+ . References [6] and [8] actually consider an insertion of the closely related gauge invariant zero-energy photon vertex operator, denoted $V_{z,0}$ in equation (2.46). This differs from J_z by the insertion of A_z at spatial infinity which will not give charge-dependent terms in (2.54) (but is relevant for multiple current insertions).

It is interesting to see how the collinear momentum space singularity in (2.55) becomes the pole in (2.54) in the simplest context: a more detailed and complete treatment is in [8]. For this we need the transformation from our retarded coordinates to the usual flat Minkowski coordinates

$$\begin{aligned} ds^2 &= -dt^2 + d\vec{x} \cdot d\vec{x}, \\ t &= v + r, \\ x^1 + ix^2 &= r \frac{2z}{1 + z\bar{z}}, \\ x^3 &= r \frac{1 - z\bar{z}}{1 + z\bar{z}}, \end{aligned} \tag{2.56}$$

with $\vec{x} = (x^1, x^2, x^3)$ obeying $\vec{x} \cdot \vec{x} = r^2$. At late times and large r the wave packet for a massless particle with charge q and spatial momentum centered around \vec{p} becomes localized on the conformal sphere near the point

$$\frac{\vec{p}}{\omega} = \frac{\vec{x}}{r} = \frac{1}{1 + z\bar{z}}(z + \bar{z}, -iz + i\bar{z}, 1 - z\bar{z}). \tag{2.57}$$

where $\vec{p} \cdot \vec{p} = \omega^2$. Consider the example of a soft photon with momentum $p_\gamma^3 = \omega_\gamma$ which hits the asymptotic sphere at $z = 0$. The positive helicity spatial polarization vector obeying $\vec{\varepsilon} \cdot \vec{p}_\gamma = 0$ is

$$\vec{\varepsilon} = (1, -i, 0). \tag{2.58}$$

These obey

$$p_\gamma \cdot p = -2\omega\omega_\gamma \frac{z\bar{z}}{(1 + z\bar{z})}, \quad \varepsilon \cdot p = 2\omega \frac{\bar{z}}{(1 + z\bar{z})}. \tag{2.59}$$

The Weinberg soft factor is

$$\frac{q\varepsilon \cdot p}{p_\gamma \cdot p} = -\frac{1}{\omega_\gamma} \frac{q}{z}, \tag{2.60}$$

which exhibits the collinear pole at $z = 0$ as in (2.54). The factor of ω_γ is ultimately cancelled in (2.54) because it involves a zero mode of F_{vz} rather than A_z [8].

Let us now return to (2.54) and consider an integral of the current J_z convoluted with ε around a contour C encircling all the operator insertions at (z_1, \dots, z_n) . One immediately finds.

$$\begin{aligned} \langle 0|V_1(v_1, z_1, \bar{z}_1)\dots V_n(v_n, z_n, \bar{z}_n)\frac{1}{2\pi i}\oint_C \varepsilon(z)J_z dz\dots|0\rangle &= \sum_{k=1}^n q_k \varepsilon(z_k)\langle 0|V_1(v_1, z_1, \bar{z}_1)\dots V_n(v_n, z_n, \bar{z}_n)\dots|0\rangle \\ &= -i\delta_\varepsilon\langle 0|V_1(v_1, z_1, \bar{z}_1)\dots V_n(v_n, z_n, \bar{z}_n)|0\rangle. \end{aligned} \quad (2.61)$$

Hence large gauge transformations in the region bounded by a contour C are generated by insertions of contour integrals $\oint_C J$. This identifies Weinberg's soft-photon relations between S-matrix elements as a Ward identity of the large gauge symmetry (2.16).

2.5 Green functions

In this section we record the radiation-gauge two-point functions in the free quantum theory, and take their limits to obtain boundary correlators on \mathcal{I}^+ . The standard radiation gauge $A_t = 0 = \nabla^\mu A_\mu$ is the same as $A_v = 0 = \nabla^\mu A_\mu$. The time-ordered Feynman two-point function is, in the flat Minkowski coordinates (2.56)⁷

$$\frac{1}{e^2}\langle 0|T(A_i(x)A_j(x'))|0\rangle = \delta_{ij}D_F(x, x') - \partial_i\partial'_j\Delta(x, x'), \quad (2.62)$$

where $i, j = 1, 2, 3$, $x \sim (t, \vec{x})$,

$$D_F(x, x') = i\int \frac{d^4k}{(2\pi)^4} \frac{e^{ik\cdot(x-x')}}{(k_0^2 - \vec{k}^2 + i\varepsilon)}, \quad \nabla^2 D_F(x, x') = i\delta^4(x, x'), \quad (2.63)$$

is the Feynman two-point function for a massless scalar and

$$\Delta = -\bar{\partial}^{-2}D_F, \quad (2.64)$$

where

$$\begin{aligned} \bar{\partial}^{-2}\delta^3(\vec{x}, \vec{x}') &= -\frac{1}{4\pi\sqrt{(\vec{x} - \vec{x}')^2}} \\ &\equiv -\frac{1}{4\pi\ell}. \end{aligned} \quad (2.65)$$

is the inverse of the spatial laplacian $\bar{\partial}^2$. One finds

$$\partial_t^2\Delta = \frac{1}{4\pi^2((|t-t'| + i\varepsilon)^2 - \ell^2)} - \frac{i\delta(t-t')}{4\pi\ell}, \quad (2.66)$$

⁷See *e.g.* equation 14.54 in [10].

Integrating twice and suppressing the $i\varepsilon$ yields

$$\Delta = \frac{1}{8\pi^2} \left(\frac{t-t'}{\ell} \ln \frac{\ell-t+t'}{\ell+t-t'} - \ln(\ell^2 - (t-t')^2) \right) - i \frac{|t-t'|}{8\pi\ell}. \quad (2.67)$$

We have fixed the integration constant using (2.64). In Bondi coordinates we have

$$\ell^2 = (r-r')^2 + 4rr' \frac{(z-z')(\bar{z}-\bar{z}')}{(1+z\bar{z})(1+z'\bar{z}')}. \quad (2.68)$$

Commutators of gauge-variant operators are in general nonlocal. For example it follows from (2.62) that at equal times

$$[\partial_t A_i(t, \vec{x}), A_j(t, \vec{x}')] = -ie^2 \delta_{ij} \delta^3(\vec{x}, \vec{x}') - i \partial_i \partial'_j \frac{e^2}{4\pi\ell}. \quad (2.69)$$

The boundary correlators are limits of the two point functions in which we take $r = r' \rightarrow \infty$ with the other coordinates fixed. To leading order in $\frac{1}{r}$

$$\begin{aligned} \ell^2(x, x') &= 4r^2 \frac{(z-z')(\bar{z}-\bar{z}')}{(1+z'\bar{z}')(1+z\bar{z})} + \dots, \\ \Delta(x, x') &= -\frac{1}{8\pi^2} \left(2 \ln r + \ln \frac{4(z-z')(\bar{z}-\bar{z}')}{(1+z'\bar{z}')(1+z\bar{z})} + \dots \right). \end{aligned} \quad (2.70)$$

We then have

$$\langle 0 | A_z^{(0)}(v, z, \bar{z}) A_{\bar{z}}^{(0)}(v', z', \bar{z}') | 0 \rangle_{\mathcal{I}^+} = -e^2 \frac{\partial^2 \Delta}{\partial z \partial z'} + e^2 \partial_z \vec{x} \cdot \partial_{z'} \vec{x} D_F(x, x') = \frac{e^2}{8\pi^2 (z-z')^2} \left(1 - \frac{\ell^2}{4r^2} \right). \quad (2.71)$$

2.6 Kac-Moody level?

The Kac-Moody current J_z , which is proportional to the gauge potential, transforms non-trivially under large gauge transformations:⁸

$$\delta_\varepsilon J_z = -\frac{4\pi}{e^2} \partial_z \varepsilon. \quad (2.72)$$

Often such a transformation law for a Kac-Moody current indicates a nonzero level, in this case proportional to $\frac{4\pi}{e^2}$. However such an interpretation is too hasty here as we have not succeeded in obtaining this transformation law as some kind of commutator of J_z with itself. Moreover the level can be thought of as the norm of identity descendants, and a norm has not been introduced yet. In the usual 4D field theory adjoint $J_z^\dagger = J_{\bar{z}}$, which is not appropriate for a CFT₂ current. These issues presumably must be resolved before inferring a level from (2.72). Hence at this point we simply regard (2.72) as an interesting hint to guide further investigations.

⁸ In general the normalization of a $U(1)$ current is arbitrary. We have fixed this ambiguity by requiring equation (2.16) hold with the factor written and periodicity (2.3).

3 The non-Abelian case

The nonabelian case with group G is very similar to the abelian one, except one must keep track of a few matrix orderings. We will only sketch the results. Radiation gauge for the nonabelian theory is defined by

$$A_v = 0, \quad (3.1)$$

$$[\sqrt{-g}\nabla^\mu A_\mu]_{\mathcal{I}^-} = [\partial_z A_{\bar{z}} + \partial_{\bar{z}} A_z - r^2 \partial_v A_r + \partial_r r^2 A_r]_{\mathcal{I}^-} = 0. \quad (3.2)$$

This is the same as the Abelian case except that A_μ is now valued in the Lie algebra of G . We also assume here no color flux incoming on \mathcal{I}^- .

Expanding near \mathcal{I}^+

$$\begin{aligned} A_z(r, v, z, \bar{z}) &= \sum_{n=0}^{\infty} \frac{A_z^{(n)}(v, z, \bar{z})}{r^n}, \\ A_r(r, v, z, \bar{z}) &= \sum_{n=0}^{\infty} \frac{A_r^{(n)}(v, z, \bar{z})}{r^{n+2}}, \end{aligned} \quad (3.3)$$

at leading order we obtain

$$\begin{aligned} F_{z\bar{z}}^{(0)} &= \partial_z A_{\bar{z}}^{(0)} - \partial_{\bar{z}} A_z^{(0)} + [A_z^{(0)}, A_{\bar{z}}^{(0)}], \\ F_{vz}^{(0)} &= \partial_v A_z^{(0)}, \\ F_{rz}^{(0)} &= -D_z^{(0)} A_r^{(0)} - A_z^{(1)}, \\ F_{rv}^{(0)} &= -\partial_v A_r^{(0)}, \end{aligned} \quad (3.4)$$

where $D_z^{(0)} = \partial_z + [A_z^{(0)}, \]$. Radiation gauge locally leaves unfixed large nonabelian gauge transformations $\varepsilon(z)$:

$$\delta A_z^{(0)} = D_z^{(0)} \varepsilon. \quad (3.5)$$

The constraint equation on \mathcal{I}^+ is

$$-\partial_v (\partial_{\bar{z}} A_z^{(0)} + \partial_z A_{\bar{z}}^{(0)} - \gamma_{z\bar{z}} \partial_v A_r^{(0)}) = [A_z^{(0)}, \overleftrightarrow{\partial}_v A_{\bar{z}}^{(0)}] + \gamma_{z\bar{z}} g_{\text{YM}}^2 j_v^{(0)M}. \quad (3.6)$$

where j_v^{0M} is the leading component of any matter color current on \mathcal{I}^+ and the commutator is the gluon color current. Assuming the field strengths vanish at \mathcal{I}_+^+ , integrating (3.6) gives

$$-[\partial_{\bar{z}} A_z^{(0)} + \partial_z A_{\bar{z}}^{(0)}]_{v=\infty} = \int_{-\infty}^{\infty} dv \left([A_z^{(0)}, \overleftrightarrow{\partial}_v A_{\bar{z}}^{(0)}] + \gamma_{z\bar{z}} g_{\text{YM}}^2 j_v^{(0)M} \right). \quad (3.7)$$

Defining

$$J_z(z, \bar{z}) = -\frac{4\pi}{g_{\text{YM}}^2} A_z^{(0)}(\infty, z, \bar{z}), \quad (3.8)$$

adding $F_{z\bar{z}}^{(0)}|_{v=\infty} = 0$ to the left hand side of (3.7) and then subtracting $[A_z^{(0)}, A_{\bar{z}}^{(0)}]_{v=\infty}$ from both sides gives

$$\partial_{\bar{z}} J_z = 2\pi \int_{-\infty}^{\infty} dv \left(\gamma_{z\bar{z}} j_v^{0E} + \frac{2}{g_{\text{YM}}^2} [A_{\bar{z}}^{(0)}, \partial_v A_z^{(0)}] \right) \equiv 2\pi q_{z\bar{z}}. \quad (3.9)$$

This implies

$$J_z(z) = \int d^2 w \frac{q_{w\bar{w}}(w, \bar{w})}{(z-w)}. \quad (3.10)$$

As in the abelian case the gauge variation of the current

$$\delta_\varepsilon J_z = -\frac{4\pi}{g_{\text{YM}}^2} \partial_z \varepsilon + [J_z, \varepsilon] \quad (3.11)$$

hints at a level proportional to $\frac{4\pi}{g_{\text{YM}}^2}$. The Ward identity (2.54) becomes

$$\langle 0 | V_1(v_1, z_1, \bar{z}_1) \dots V_n(v_n, z_n, \bar{z}_n) J_z^a(z, \bar{z}) \dots | 0 \rangle = \sum_{k=1}^n \frac{T_k^a}{z - z_k} \langle 0 | V_1(v_1, z_1, \bar{z}_1) \dots V_n(v_n, z_n, \bar{z}_n) \dots | 0 \rangle \quad (3.12)$$

where we display the adjoint index a and T_k^a is a generator in the representation of and acts on the operator insertion at z_k . This is the soft gluon theorem transformed from momentum space to spherical coordinates on \mathcal{I}^+ .

4 Stringy realization?

Our arguments so far have generally applied to any 4D gauge theory with massless charges. However the reader familiar with string theory may have found some features familiar from the study of string scattering amplitudes. The role of the conformal sphere at \mathcal{I}^+ resembles that of the string worldsheet: see for example [13, 14]. Perhaps this connection can be made precise: we give here a few speculative comments.

Our analysis applies in particular to any perturbative closed string compactification to four dimensions with a non-abelian gauge symmetry G . For such theories the string worldsheet has a level k_{ws} G -current algebra with an OPE

$$j^a(z) j^b(w) \sim \frac{k_{ws} \delta^{ab}}{(z-w)^2} + \frac{f^{abc} j^c(w)}{(z-w)^2}. \quad (4.1)$$

We use roman case to distinguish the worldsheet coordinate z from the spacetime coordinate z . The four dimensional space-time string coupling g_s , and worldsheet level k_{ws} are related by [11]

$$\frac{k_{ws}}{g_s^2} \frac{V_6}{4(2\pi)^6} = \frac{4\pi}{g_{\text{YM}}^2}, \quad (4.2)$$

where V_6 is the compactification volume in string units. On the other hand this paper raised the possibility of a relation of the form

$$k_{\text{spacetime}} \sim \frac{4\pi}{g_{\text{YM}}^2} = \frac{k_{\text{ws}}}{g_{\text{S}}^2} \frac{V_6}{4(2\pi)^6}. \quad (4.3)$$

The world sheet current algebra (4.1) generated by $j_z^a(z)$ is of course not manifestly the same as the spacetime current algebra generated by $J_z^a(z)$. Nevertheless the action of our Kac-Moody on the conformal sphere at \mathcal{I}^+ resembles the action of the worldsheet Kac-Moody on the string worldsheet. It is often the case in string theory that worldsheet symmetries lift to space-time symmetries. Consider for example strings on $AdS_3 \times S^3 \times M^4$ with NS fluxes. The string world sheet has a level k_{ws} $SU(2)_L \times SU(2)_R$ current algebra. As shown in [12], this lifts to a *spacetime* current algebra in the boundary CFT_2 with level

$$k_{\text{spacetime}} = \frac{k_{\text{ws}}^2}{g_{\text{S}}^2} \frac{V_4}{(2\pi)^4}, \quad (4.4)$$

where V_4 is the string-unit volume of M_4 . This formula bears some similarity to (4.3), which one hopes might be derived by similar methods.

Acknowledgements

I would like to thank J. Bourjaily, M. Guica, P. Mitra, G. S. Ng and W. Song for stimulating conversations. I am particularly grateful to J. Maldacena for many discussions and for sharing the significant unpublished work [8]. This work is partly supported by DOE grant DE-FG02-91ER40654.

References

- [1] H. Bondi, M. G. J. van der Burg, A. W. K. Metzner, “Gravitational waves in general relativity VII. Waves from isolated axisymmetric systems”, Proc. Roy. Soc. Lond. A 269, 21 (1962); R. K. Sachs, “Gravitational waves in general relativity VIII. Waves in asymptotically flat space-time”, Proc. Roy. Soc. Lond. A 270, 103 (1962).
- [2] G. Barnich and C. Troessaert, “Symmetries of asymptotically flat 4 dimensional spacetimes at null infinity revisited,” Phys. Rev. Lett. **105**, 111103 (2010) [arXiv:0909.2617 [gr-qc]]; “Supertranslations call for superrotations,” PoS CNCFG **2010**, 010 (2010); [arXiv:1102.4632 [gr-qc]] “BMS charge algebra,” JHEP **1112**, 105 (2011) [arXiv:1106.0213 [hep-th]].

- [3] See footnote 18 in T. Banks, “A Critique of pure string theory: Heterodox opinions of diverse dimensions,” hep-th/0306074.
- [4] A. A. Belavin, A. M. Polyakov and A. B. Zamolodchikov, “Infinite Conformal Symmetry in Two-Dimensional Quantum Field Theory,” Nucl. Phys. B **241**, 333 (1984).
- [5] A. Strominger, in progress.
- [6] S. Weinberg, “Infrared photons and gravitons,” Phys. Rev. **140**, B516 (1965).
- [7] A. P. Balachandran and S. Vaidya, “Spontaneous Lorentz Violation in Gauge Theories,” Eur. Phys. J. Plus **128**, 118 (2013) [arXiv:1302.3406 [hep-th]].
- [8] J. Maldacena and A. Zhiboedov, ”Notes on Soft Factors” unpublished (2012) and private communication.
- [9] E. T. Newman, R. Penrose, ”An approach to gravitational radiation by a method of Spin coefficients”, J. Math. Phys. 3, 566 (1962); R. Penrose, ”Asymptotic properties of fields and spacetimes”, Phys. Rev. Lett. 10, 66 (1963).
- [10] J. D. Bjorken and S. D. Drell, “Relativistic quantum fields,” ISBN-0070054940.
- [11] J. Polchinski, “String theory. Vol. 2: Superstring theory and beyond,” Cambridge, UK: Univ. Pr. (1998) 531 p
- [12] A. Giveon, D. Kutasov and N. Seiberg, “Comments on string theory on AdS(3),” Adv. Theor. Math. Phys. **2**, 733 (1998) [hep-th/9806194].
- [13] D. J. Gross and P. F. Mende, “String Theory Beyond the Planck Scale,” Nucl. Phys. B **303**, 407 (1988).
- [14] A. K. H. Bengtsson, L. Brink and S. -S. Kim, “Counterterms in Gravity in the Light-Front Formulation and a D=2 Conformal-like Symmetry in Gravity,” JHEP **1303**, 118 (2013) [arXiv:1212.2776 [hep-th]].