



Disinhibition, an emerging pharmacology of learning and memory

Citation

Möhler, Hanns, and Uwe Rudolph. 2017. "Disinhibition, an emerging pharmacology of learning and memory." F1000Research 6 (1): F1000 Faculty Rev-101.
doi:10.12688/f1000research.9947.1. <http://dx.doi.org/10.12688/f1000research.9947.1>.

Published version

<https://doi.org/10.12688/f1000research.9947.1>

Link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:31731798>

Terms of use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material (LAA), as set forth at

<https://harvardwiki.atlassian.net/wiki/external/NGY5NDE4ZjgzNTc5NDQzMGIzZWZhMGFIOWI2M2EwYTg>

Accessibility

<https://accessibility.huit.harvard.edu/digital-accessibility-policy>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#)



Check for updates

REVIEW

Disinhibition, an emerging pharmacology of learning and memory [version 1; referees: 3 approved]

Hanns Möhler^{1,2}, Uwe Rudolph^{3,4}¹Institute of Pharmacology, University of Zurich, Zurich, Switzerland²Department of Chemistry and Applied Biosciences, Swiss Federal Institute of Technology Zurich, Zurich, Switzerland³Laboratory of Genetic Neuropharmacology, McLean Hospital, Belmont, MA, USA⁴Department of Psychiatry, Harvard Medical School, Boston, MA, USA**v1** First published: 03 Feb 2017, 6(F1000 Faculty Rev):101 (doi: 10.12688/f1000research.9947.1)

Latest published: 03 Feb 2017, 6(F1000 Faculty Rev):101 (doi: 10.12688/f1000research.9947.1)

Abstract

Learning and memory are dependent on interactive excitatory and inhibitory mechanisms. In this review, we discuss a mechanism called disinhibition, which is the release of an inhibitory constraint that effectively results in an increased activity in the target neurons (for example, principal or projection neurons). We focus on discussing the role of disinhibition in learning and memory at a basic level and in disease models with cognitive deficits and highlight a strategy to reverse cognitive deficits caused by excess inhibition, through disinhibition of $\alpha 5$ -containing GABA_A receptors mediating tonic inhibition in the hippocampus, based on subtype-selective negative allosteric modulators as a novel class of drugs.

Open Peer Review

Referee Status:

	Invited Referees		
	1	2	3
version 1			
published 03 Feb 2017			

F1000 Faculty Reviews are commissioned from members of the prestigious F1000 Faculty. In order to make these reviews as comprehensive and accessible as possible, peer review takes place before publication; the referees are listed below, but their reports are not formally published.

- Bernhard Luscher**, Pennsylvania State University USA, Pennsylvania State University USA, Pennsylvania State University USA
- James Rowlett**, Center for Psychiatric Neuroscience, University of Mississippi Medical Center USA
- Trevor Smart**, University College London UK

Discuss this article

Comments (0)

Corresponding author: Hanns Möhler (mohler@pharma.uzh.ch)

How to cite this article: Möhler H and Rudolph U. **Disinhibition, an emerging pharmacology of learning and memory [version 1; referees: 3 approved]** *F1000Research* 2017, **6**(F1000 Faculty Rev):101 (doi: [10.12688/f1000research.9947.1](https://doi.org/10.12688/f1000research.9947.1))

Copyright: © 2017 Möhler H and Rudolph U. This is an open access article distributed under the terms of the [Creative Commons Attribution Licence](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Competing interests: In the last three years, UR received compensation from Concert Pharmaceuticals for professional services. HM declares that he has no competing interests.

First published: 03 Feb 2017, **6**(F1000 Faculty Rev):101 (doi: [10.12688/f1000research.9947.1](https://doi.org/10.12688/f1000research.9947.1))

Introduction

Cognitive disabilities are abundant in brain disorders. Apart from a loss of neurons in neurodegenerative diseases and various neurological states, the cognitive deficits are frequently attributed to dysfunctional operations of microcircuits and neuronal networks. This is thought to be particularly the case in neurodevelopmental disorders such as autism spectrum disorders (ASDs) or Down syndrome (DS), in psychiatric disorders such as schizophrenia, but also in age-related cognitive decline. In line with this view, research on cognitive behavior has focused on the role of microcircuits.

Disinhibition was recognized in recent years as an emerging general mechanism of learning and memory. Memories are acquired and encoded within large-scale neuronal networks spanning different brain areas with the respective information being processed by projection neurons in a tightly controlled balance of excitation and inhibition. Recent findings demonstrate that salient events, such as a footshock in aversive learning, often elicit disinhibition of projection neurons (that is, “a selective and transient reduction of synaptic inhibition received by projection neurons that changes their computation”¹). The ensuing stronger firing of the projection neuron is likely to gate the induction of synaptic plasticity, required for memory formation. Disinhibition is displayed in different compartments of projection neurons, in diverse cortical areas, and on time scales ranging from milliseconds to days. Behavioral functions of disinhibition range from critical

period plasticity², addiction³, and Pavlovian learning to spatial navigation¹. In this article, we will explore the evidence suggesting that disinhibition could become a therapeutic principle for reversal of disease-related cognitive deficits.

Disinhibition in behaving animals

Research on disinhibition in behaving animals so far has been strongly focused on aversive learning by fear conditioning, a type of Pavlovian learning. During the acquisition of auditory fear conditioning, disinhibition plays a role in cortical auditory plasticity, which depends on the convergence of activity evoked by both an unconditioned stimulus (for example, footshock) and a conditioned stimulus (for example, tone). Footshocks, which drive learning in this paradigm, elicit strong firing in nearly all layer 1 vasoactive intestinal polypeptide (VIP) interneurons of the auditory cortex, driven by acetylcholine released from basal forebrain afferents. The enhanced activity of inhibitory layer 1 VIP interneurons inhibits layer 1 somatostatin (SOM)-containing GABAergic interneurons, which target the dendrites of principal cells, and parvalbumin (PV)-containing GABAergic interneurons in layer 2/3, which target the soma. With this disinhibitory circuit, the footshock stimulus results in a reduction of inhibition over the entire somatodendritic domain (Figure 1A). The firing of the cortical projection neurons is enhanced⁴.

Disinhibition is also recruited by implementing the conditioned tone stimulus as apparent in the amygdala. Principal cells in the




Figure 1. Targets of disinhibition. (A) Disinhibition refers to the selective and transient reduction of synaptic inhibition of a projection neuron due to suppression of interneuron firing by another population of interneurons. A circuit model is given for the cortical disinhibition elicited by an unconditioned stimulus (US) (footshock) in auditory Pavlovian fear learning (modified from 1). Blue color denotes the source of disinhibition, yellow color the inhibited interneurons, and green color the disinhibited pyramidal neurons. Flat bars denote inhibitory inputs. ACh, acetylcholine; CS, conditioned stimulus; PV, parvalbumin; VIP, vasoactive intestinal polypeptide. (B) Immunohistochemical distribution of the $\alpha 5$ subunit of GABA_A receptors in mouse brain with false color coding. High expression in hippocampus and cortical layer 5 is shown. (C) Scheme of the extrasynaptic localization of $\alpha 5$ GABA_A receptors on dendrites and dendritic spines of hippocampal pyramidal cells, representing the initial, tonic inhibitory control to incoming excitatory signals. Figure 1B provided courtesy of Jean-Marc Fritschy.

basolateral amygdala (BLA) are under the control of inhibitory GABAergic interneurons of which those containing SOM target the dendrites and those containing PV target the soma. During an auditory cue, PV interneurons are excited and indirectly disinhibit the dendrites of the BLA principal cells via SOM interneurons. Thus, activation of PV interneurons by the tone leads to a disinhibition of projection cell dendrites, the site where auditory inputs arrive. Importantly, the dendritic disinhibition overcomes the increases in perisomatic inhibition mediated by excited PV interneurons during the tone, which is the crucial factor in enhancing projection cell tone responses and promoting cue-shock associations in fear learning⁵.

In addition to its role in memory acquisition, disinhibition is a factor in memory expression. In auditory fear-conditioned mice, the presentation of a conditioned tone stimulus causes strong phasic inhibition associated with inhibition of a subset of PV-positive interneurons in the dorso-medial prefrontal cortex, selectively when mice display a fear reaction. The ensuing disinhibition of projection neurons is required for memory expression. Indeed, in naïve animals, optogenetic inhibition of the corresponding PV cells proved sufficient to induce freezing, indicating that this form of disinhibition is both necessary and sufficient for memory expression. Disinhibition permits stronger responses of the projection neurons to tones that likely drive fear expression in downstream areas such as the amygdala¹. Finally, the behavioral motor response of freezing is also triggered by disinhibition. An inhibitory pathway from the central nucleus of the amygdala to the periaqueductal gray (PAG) produces freezing by disinhibition of the excitatory PAG output to the pre-motor targets in the medulla⁶.

Genetic disinhibition: downregulation of $\alpha 5$ GABA_A receptors enhances learning and memory

To exploit disinhibition pharmacologically, circuits associated selectively with learning and memory would have to be targeted. Starting in 2002, the $\alpha 5$ GABA_A receptor subtype (defined as GABA_A receptors containing the $\alpha 5$ subunit and β and γ subunits) took center stage (Figure 1B, C). It is mainly expressed in hippocampal pyramidal cells and layer 5 pyramidal cell dendrites, is located extrasynaptically, generates tonic inhibition, and is capable of altering neural oscillations, which is considered to impact on cognitive behavior⁷⁻⁹. In genetically modified mice, a partial knockdown of $\alpha 5$ receptors improved hippocampus-dependent performance, as shown in trace fear conditioning¹⁰ and appetitive conditioning¹¹, and a full knockdown in $\alpha 5$ knockout mice resulted in improved performance in the water maze¹² and improved novel object recognition¹³, while hippocampus-independent learning was unaltered in both partial and full $\alpha 5$ knockout mice^{10,13}. Furthermore, although low-frequency stimulation elicited long-term depression in hippocampal slices of normal mice, the same stimulation elicited long-term potentiation (LTP) in slices from $\alpha 5$ knockout mice, suggesting that the receptor sets the threshold for eliciting LTP and contributes to memory formation¹³. These results make $\alpha 5$ GABA_A receptors a key target for a pharmacological enhancement of hippocampus-dependent learning and memory¹⁴⁻¹⁶.

However, the role of $\alpha 5$ GABA_A receptors in memory formation may not be as straightforward as described above but depends on the cognitive domain and the context and demand of the task. These findings are a cautionary note on $\alpha 5$ GABA_A receptors as targets for $\alpha 5$ -negative allosteric modulators ($\alpha 5$ -NAMs). For instance, $\alpha 5$ knockout mice exhibited a deficit in short-term memory but only when a particular task (puzzle box) became progressively more difficult, whereas long-term memory deficits were not affected¹⁷. A deficit in learning selectivity was apparent in mice with a reduced $\alpha 5$ expression as shown by a deficit in latent inhibition¹⁸ and prepulse inhibition¹⁹. The memory for location of objects was impaired in these mice, when tested in a complex setting²⁰.

In a conditional knockdown of *Gabra5* selectively in dentate gyrus, performance was impaired in tasks characterized by high memory interference. Such tasks included behavioral pattern separation, when mice had to distinguish between an aversive context and a similar safe context, reversal learning in the Morris water maze, fear extinction, and latent inhibition to conditioned freezing²¹. In contrast, in tasks characterized by low memory interference (for example, novel object recognition), initial spatial learning in the Morris water maze and fear conditioning to tone, performance was unaltered²¹. In brain development, a heterozygous $\alpha 5$ knockout, as well as single-cell deletion of $\alpha 5$ in newborn granule cells, caused severe alterations of migration and dendrite development in the dentate gyrus, indicating that relatively minor imbalances in $\alpha 5$ GABA_A receptor-mediated transmission may have major consequences for neuronal plasticity²².

Negative allosteric modulators of $\alpha 5$ GABA_A receptors: a new class of drugs

In view of the memory-enhancing findings in the genetically modified mouse models, partial NAMs acting at the benzodiazepine site of $\alpha 5$ GABA_A receptors were expected to improve performance in learning and memory. The first ligands of this type were $\alpha 5$ IA, L-655,708, and MRK-016 synthesized by the Merck group. The $\alpha 5$ IA was selective for $\alpha 5$ receptors only by efficacy and showed equal affinity for $\alpha 1$, $\alpha 2$, $\alpha 3$, and $\alpha 5$ receptors. It improved spatial learning without being anxiogenic or proconvulsant. In elderly volunteers, paired associate learning was enhanced but remained inconsistent. Renal toxicity prevented further development^{23,24}. L-655,708 showed preferential affinity for $\alpha 5$ GABA_A receptors but acted as NAM not only at $\alpha 5$ but also, at higher concentration, at $\alpha 1$, $\alpha 2$, and $\alpha 3$ receptors. It enhanced spatial learning and induced gamma oscillations in hippocampal slices without being proconvulsant, but its anxiogenic effect, presumably due to interactions with receptors other than $\alpha 5$, prevented further development²⁴. MRK-016, like L-655,708, showed preferential affinity for $\alpha 5$ receptors and displayed nootropic properties (for a review, see 14). Most interestingly, disinhibition by L-655,708 and MRK-016 was recently found to produce rapid (measured 24 hours after drug administration), sustained ketamine-like antidepressant effects in animal models of depression, which is likely due to disinhibition-induced plasticity²⁵. Motor recovery after stroke was also enhanced by L-655,708

after chronic dosing (42 days), although the infarct size was unaltered²⁶. Memory in young but not aged rats was improved by the $\alpha 5$ -NAM TB21007²⁷. The $\alpha 5$ -NAM PWZ-029 largely lacks functional selectivity, being a weak positive allosteric modulator (PAM) at $\alpha 1$, $\alpha 2$, and $\alpha 3$ GABA_A receptors^{28,29}. In a task assessing working memory, it slightly increased delayed matching-to-sample (DMTS) accuracy in rhesus monkeys, which is likely due to its action as $\alpha 5$ -NAM²⁹.

The imidazo-triazolo-benzodiazepine RO4938581 was the first partial $\alpha 5$ -NAM with a highly selective preference in both affinity (up to 40-fold) and efficacy (10-fold) for $\alpha 5$ GABA_A receptors versus $\alpha 1$, $\alpha 2$, and $\alpha 3$ GABA receptors^{30,31}. This compound rescued deficits in working memory and spatial memory³⁰ and improved executive functions in an object retrieval task in cynomolgus monkeys³⁰. Most importantly, RO4938581 showed no anxiogenic or pro-convulsive activity. Such agents hold the promise of novel treatments for neurological disorders with cognitive dysfunctions such as DS and cognitive impairment in psychiatric conditions.

Effectiveness of $\alpha 5$ -NAMs in a model of Down syndrome

The best characterized animal model of DS (trisomy 21) is the Ts65Dn mouse, which contains an extra segment of the ortholog mouse chromosome 16^{32–35}. The neuronal plasticity in Ts65Dn mice is thought to be obstructed by excessive GABAergic inhibition as low-dose pentylentetrazole, a non-selective GABA antagonist, reversed deficits in learning and memory³⁶. Remarkably, reducing $\alpha 5$ GABA_A receptor function by the NAM $\alpha 5$ IA was likewise sufficient to reverse the deficits in spatial reference learning and novel object recognition³⁷. The $\alpha 5$ -NAM RO4938581 also rescued the spatial performance of Ts65Dn mice, improved the neurogenesis in the dentate gyrus, and normalized the enhanced density of hippocampal GABAergic boutons^{38–40}.

Clinical trials with RG1662 (Basmisanil), a compound related to RO4938581, were initiated by Roche to improve cognitive disabilities in adults, adolescents, and children with DS (www.clinicaltrials.gov; Drug RG1662). The phase II trials were recently terminated due to lack of efficacy (Roche Press Release, June 28, 2016; <http://www.roche.com/media/store/statements.htm>). This outcome suggests either that the $\alpha 5$ -NAM was insufficiently effective to restore neuronal plasticity or alternatively that the hypothesis of excessive GABAergic inhibition obstructing neuronal plasticity does not extend to individuals with DS.

Schizophrenia: cognitive deficits and $\alpha 5$ -NAMs

Cognitive deficits are a core symptom of schizophrenia. They are disabling and difficult to treat. Cortical disinhibition has been proposed as an underlying pathological hallmark⁴¹. Indeed, in post-mortem brain, a striking deficit of GABAergic inhibition is apparent with several distinct cortical GABA interneurons being dysfunctional (basket cells, Chandelier neurons, SOM/NPY, or CCK interneurons) concomitant with a deficit of GAD67 and a change in GABA_A receptor expression, in particular, an increased $\alpha 2$ subunit expression on the axon initial segment⁴². A deficit in the excitatory drive to cortical PV interneurons is thought to

contribute to altered gamma oscillations and cognitive dysfunctions⁴³ as well as to negative symptoms⁴⁴. GABAergic therapeutic attempts to improve cognitive deficits focused initially on MK-0777, a ligand enhancing synaptic $\alpha 2$ and $\alpha 3$ receptor function^{45,46}. More recently, enhancing $\alpha 5$ receptor function was found to improve behavioral deficits, at least in the rat methylazoxymethanol (MAM) model of neurodevelopmental disorders, which is characterized largely by a reduction in PV interneurons. The $\alpha 5$ -PAM SH-053-2'F-R-CH₃²⁸—when administered either systemically or directly into the ventral hippocampus—reversed the hyperactivation of the dopaminergic system in the ventral tegmental area and dampened the amphetamine-induced increase in the locomotor response⁴⁷, although not under conditions of haloperidol withdrawal⁴⁸. It remains to be seen whether a selective $\alpha 5$ -PAM would impact on cognitive deficits.

On the other hand, cognitive deficits in patients with schizophrenia are frequently attributed to their inability to recruit high-frequency oscillations while performing cognitive tasks⁴⁹. Thus, disinhibition of principal cell activity with an $\alpha 5$ -NAM might be considered beneficial. In hippocampal slices, high-frequency oscillations were induced by applying the $\alpha 5$ -NAM L-655,708⁵⁰. Thus, it appears warranted to test whether selective $\alpha 5$ -NAMs are able to restore high-frequency oscillations and to ameliorate cognitive deficits in patients with schizophrenia.

Antidepressant action via $\alpha 5$ GABA_A receptors?

The GABA hypothesis of depression posits that a deficit in GABAergic inhibition of principal cells contributes to depression-like behavior^{51,52}. In this view, an increased inhibitory input to principal neurons might be desirable for circuit-based antidepressant activity. Indeed, a preliminary report indicates that the $\alpha 5$ -PAM SH-053-2'F-R-CH₃²⁸ displayed rapid antidepressant-like effects in female (but not male) mice exposed to unpredictable chronic mild stress⁵³. However, contrary to this view, a negative modulation of $\alpha 5$ GABA_A receptor with the $\alpha 5$ -NAM L-655,708 was recently shown to restore excitatory synaptic strength and to display rapid and sustained antidepressant-like actions after chronic stress in rats²⁵. The stress-induced anhedonia (sucrose preference test) and the deficit in social interactions (interaction test) were reversed by a single dose of L-655,708, measured 24 hours after administration²⁵. Clearly, the discrepancy in the potential antidepressant pharmacology of $\alpha 5$ GABA_A receptor ligands remains to be resolved with regard to animal models and ligand selectivity.

Genetic evidence for a contribution to anxiety regulation by $\alpha 5$ GABA_A receptors

Experiments in mice with partial or complete knockout of the $\alpha 5$ subunit revealed no evidence of an involvement in anxiety regulation^{10,12}. Nevertheless, recent studies showed that $\alpha 5$ GABA_A receptors in the central amygdala, specifically those on PKC δ ⁺ neurons, contribute to anxiety regulation. Both a brain area-specific cre-mediated $\alpha 5$ knockdown and a cell type-specific knockdown of $\alpha 5$ using a short hairpin RNA (shRNA) approach resulted in anxiety-related phenotypes in the elevated plus maze test and increased fear generalization³⁴. Thus, in wild-type animals, an enhanced $\alpha 5$ receptor-mediated inhibition of PKC δ ⁺

neurons would be expected to contribute to an anxiolytic response. In keeping with this view, a global enhancement selectively of $\alpha 5$ GABA_A receptor function was anxiolytic. Via a “restriction-of-function” approach in which three out of the four diazepam-sensitive α subunits ($\alpha 1$, $\alpha 2$ and $\alpha 3$), and $\alpha 5$ were rendered diazepam-insensitive by a histidine-to-arginine point mutation, it was shown that positive allosteric modulation by diazepam of $\alpha 5$ GABA_A receptors with high molecular specificity resulted in anxiolytic-like effects (elevated plus maze and light/dark choice test) at doses higher than those required for diazepam-induced anxiolysis in wild-type mice that would be sedative in wild-type animals⁵⁵. Thus, enhancing $\alpha 5$ GABA_A receptor function contributes to a reduction of anxiety-like behavior. In this context, it is important to note that a pharmacological reduction of $\alpha 5$ GABA_A receptor function by selective $\alpha 5$ -NAMs did not influence anxiety behavior, as described above.

The impaired aging brain: a role for $\alpha 5$ -PAM?

Increased hippocampal activation has become recognized as a signature of the aging human brain. Hippocampal activation, largely restricted to the CA3/DG regions, is modestly increased in older subjects but, in patients with mild cognitive impairment, greatly exceeds that of healthy age-matched controls⁵⁶. Reduction of excess neuronal activity by enhancing $\alpha 5$ GABA_A receptor-mediated tonic inhibition could potentially improve symptomatic functions and slow prodromal progression. At present, only a few $\alpha 5$ -PAM compounds are available: SH-053-2’F-R-CH₃²⁸, compound 44⁵⁷, compound 6⁵⁸, and MP-III-022⁵⁹. However, when interpreting behavioral responses, it has to be kept in mind that the $\alpha 5$ -PAMs display only limited selectivity for $\alpha 5$ GABA_A receptors. So far, the $\alpha 5$ -PAM compound 44 improved long-term memory of aged rats in the Morris water maze, and the $\alpha 5$ -PAM compound 6 improved memory performance of aged rats but not of young rats in a radial arm maze task²⁷. Their impact on cognitive behavior remains to be established. Thus, selective $\alpha 5$ -PAMs may warrant further testing on the potential amelioration of age-related memory impairments as they are related to hippocampal overactivity.

Autism spectrum disorders: amelioration by enhancing GABA_A receptor function

In a recent behavioral analysis of $\alpha 5$ knockout mice, some autism-like behavior was apparent. Social contacts as tested in the social proximity assay and the three-chamber social approach test were reduced in male mice. Self-grooming as a measure of stereotypy was increased and performance in the puzzle box, reflecting executive functions, was reduced without changes in locomotion or anxiety level¹⁷. In contrast to the behavior reported for the $\alpha 5$ knockout mice, autism-like behavior has not been reported to be induced pharmacologically by partial NAMs acting at $\alpha 5$ GABA_A receptors.

On the contrary, the observation by Zurek *et al.*¹⁷ is in line with the possibility that drugs that act as PAMs of $\alpha 5$ GABA_A receptors may ameliorate autism-like behaviors. Indeed, in ASDs, the combinatorial effects of genetic and environmental variables were postulated to cause an excess of excitation versus inhibition (E/I) within key cortical circuits during critical periods of development^{60–62}. In

post-mortem brains of patients with autism, a reduced expression of $\alpha 4$, $\alpha 5$, and $\beta 1$ subunits has been observed⁶³. Autistic traits in a broad range of ASD models share a reduction of GABAergic signaling within key cortical microcircuits as a common denominator, frequently apparent as a reduction of PV interneurons in neocortex and impaired oscillations^{64,65}. Attempts to alleviate the GABAergic deficit in ASDs pharmacologically¹⁶ were made in two animal models: the idiopathic BTBR model and the Dravet syndrome. In both cases, extremely low doses of benzodiazepines were found to be beneficial, although in a narrow dose range. Although the type of responsive GABA_A receptor remains to be established, these results warrant testing selective $\alpha 5$ -PAMs in ASDs.

The BTBR model

In the BTBR model of ASDs, a dramatic behavioral improvement in cognition and social interaction was observed after low-dose benzodiazepine treatment. Non-sedating doses of clonazepam (0.05 mg/kg) improved cognitive deficits, with memory in a context-dependent fear conditioning paradigm being improved in both the short term (30 minutes) and the long term (24 hours) after training, and spatial learning and memory in the Barnes maze was also improved. Likewise, hyperactivity was significantly reduced, as was stereotyped behavior⁶⁶. Importantly, deficits in social interactions were also improved by low-dose clobazam (0.05 mg/kg). This effect was GABA_A receptor subtype-specific since L-838,417, a partial agonist acting at the benzodiazepine site of $\alpha 2$, $\alpha 3$, and $\alpha 5$ GABA_A receptors sparing $\alpha 1$ receptors, was likewise effective⁶⁶. In contrast, zolpidem, acting preferentially on $\alpha 1$ receptors, exacerbated social deficits in BTBR mice. These findings support the notion that rebalancing of GABAergic transmission via $\alpha 2$, $\alpha 3$, or $\alpha 5$ GABA_A receptors can improve at least some of the ASD symptoms. It remains to be seen whether selective $\alpha 5$ -PAMs are similarly effective and offer a broader dose range than the benzodiazepines used so far.

Dravet syndrome

Dravet syndrome (DrS), also called severe myoclonic epilepsy of infancy, is an intractable developmental epilepsy syndrome caused by a heterozygous loss-of-function mutation in the *SCN1A* gene encoding the α subunit of the Na_v1.1 sodium channel. DrS is accompanied by neuropsychiatric comorbidities overlapping with ASDs⁶⁷. Specific heterozygous deletion of Na_v1.1 in forebrain GABAergic interneurons is sufficient to recapitulate the autistic-like behavioral and cognitive impairments such as hyperactivity, stereotyped behavior, social interaction deficits, and impaired cognition and spatial memory⁶⁸. Most remarkably, a very low dose of clonazepam (0.0625 mg/kg), which was neither anxiolytic nor sedative, completely rescued impaired social interaction and cognitive deficits in *Scn1a*^{+/-} mice⁶⁸. Clinically, clobazam is used in combination with the cytochrome P450 inhibitor stiripentol in the treatment of DrS in children⁶⁹. $\alpha 5$ -PAMs may offer an alternative approach for treatment, possibly with an improved therapeutic window.

Conclusions

The $\alpha 5$ GABA_A receptors are an exceptional target for an enhancement of learning and memory, as they are expressed largely in the hippocampus. They mediate extrasynaptic, tonic inhibition of principal cells.

Disinhibition of $\alpha 5$ GABA_A receptor function by genetic means improved hippocampus-dependent learning and memory in classic behavioral tasks but not in complex tasks involving high memory interference.

A variety of PAMs and NAMs have been developed for the $\alpha 5$ pharmacophore⁷⁰. Pharmacologically, NAMs acting at $\alpha 5$ GABA_A receptors improved cognitive and spatial performance in rodents and non-human primates. Cognitive behavioral deficits in DS Ts65Dn mice were normalized. Highly selective NAMs have reached the stage of clinical development. They will permit an assessment of their therapeutic potential in ameliorating cognitive deficits in neurological and psychiatric conditions, including schizophrenia⁷¹.

In contrast, other disease conditions may benefit from an enhancement of $\alpha 5$ GABA_A receptors. Hippocampal overactivity is a signature of the age-impaired brain, which may respond to $\alpha 5$ -PAM treatment to improve function. Furthermore, autism-like

behavior likewise may be ameliorated by $\alpha 5$ -PAM. The beneficial effect on stereotypies and cognitive impairments, seen with non-selective GABAergic drugs in two animal models, appears to justify a focus on $\alpha 5$ GABA_A receptors.

Competing interests

In the last three years, UR received compensation from Concert Pharmaceuticals for professional services. HM declares that he has no competing interests.

Grant information

UR was supported by grant R01GM086448 from the National Institute of General Medical Sciences and grant R01MH095905 from the National Institute of Mental Health, National Institutes of Health.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References



- Letzkus JJ, Wolff SB, Luthi A: **Disinhibition, a Circuit Mechanism for Associative Learning and Memory.** *Neuron*. 2015; **88**(2): 264–76.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Werker JF, Hensch TK: **Critical periods in speech perception: new directions.** *Annu Rev Psychol*. 2015; **66**: 173–96.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Lüscher C: **The Emergence of a Circuit Model for Addiction.** *Annu Rev Neurosci*. 2016; **39**: 257–76.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Letzkus JJ, Wolff SB, Meyer EM, *et al.*: **A disinhibitory microcircuit for associative fear learning in the auditory cortex.** *Nature*. 2011; **480**(7377): 331–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Wolff SB, Gründemann J, Tovote P, *et al.*: **Amygdala interneuron subtypes control fear learning through disinhibition.** *Nature*. 2014; **509**(7501): 453–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Tovote P, Esposito MS, Botta P, *et al.*: **Midbrain circuits for defensive behaviour.** *Nature*. 2016; **534**(7606): 206–12.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Fritschy JM, Mohler H: **GABA_A-receptor heterogeneity in the adult rat brain: differential regional and cellular distribution of seven major subunits.** *J Comp Neurol*. 1995; **359**(1): 154–94.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Mann EO, Mody I: **Control of hippocampal gamma oscillation frequency by tonic inhibition and excitation of interneurons.** *Nat Neurosci*. 2010; **13**(2): 205–12.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Bonin RP, Martin LJ, MacDonald JF, *et al.*: **$\alpha 5$ GABA_A receptors regulate the intrinsic excitability of mouse hippocampal pyramidal neurons.** *J Neurophysiol*. 2007; **98**(4): 2244–54.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Crestani F, Keist R, Fritschy JM, *et al.*: **Trace fear conditioning involves hippocampal $\alpha 5$ GABA_A receptors.** *Proc Natl Acad Sci U S A*. 2002; **99**(13): 8980–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Yee BK, Hauser J, Dolgov VV, *et al.*: **GABA receptors containing the $\alpha 5$ subunit mediate the trace effect in aversive and appetitive conditioning and extinction of conditioned fear.** *Eur J Neurosci*. 2004; **20**(7): 1928–36.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Collinson N, Kuenzi FM, Jarolimek W, *et al.*: **Enhanced learning and memory and altered GABAergic synaptic transmission in mice lacking the alpha 5 subunit of the GABA_A receptor.** *J Neurosci*. 2002; **22**(13): 5572–80.
[PubMed Abstract](#)
- Martin LJ, Zurek AA, MacDonald JF, *et al.*: **$\alpha 5$ GABA_A receptor activity sets the threshold for long-term potentiation and constrains hippocampus-dependent memory.** *J Neurosci*. 2010; **30**(15): 5269–82.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Atack JR: **GABA_A receptor subtype-selective modulators. II. $\alpha 5$ -selective inverse agonists for cognition enhancement.** *Curr Top Med Chem*. 2011; **11**(9): 1203–14.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Rudolph U, Mohler H: **GABA_A receptor subtypes: Therapeutic potential in Down syndrome, affective disorders, schizophrenia, and autism.** *Annu Rev Pharmacol Toxicol*. 2014; **54**: 483–507.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Möhler H: **The legacy of the benzodiazepine receptor: from flumazenil to enhancing cognition in Down syndrome and social interaction in autism.** *Adv Pharmacol*. 2015; **72**: 1–36.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Zurek AA, Kemp SW, Aga Z, *et al.*: **$\alpha 5$ GABA_A receptor deficiency causes autism-like behaviors.** *Ann Clin Transl Neurol*. 2016; **3**(5): 392–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Gerdjikov TV, Rudolph U, Keist R, *et al.*: **Hippocampal $\alpha 5$ subunit-containing GABA_A receptors are involved in the development of the latent inhibition effect.** *Neurobiol Learn Mem*. 2008; **89**(2): 87–94.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Hauser J, Rudolph U, Keist R, *et al.*: **Hippocampal $\alpha 5$ subunit-containing GABA_A receptors modulate the expression of prepulse inhibition.** *Mol Psychiatry*. 2005; **10**(2): 201–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Prut L, Prenosil G, Willadt S, *et al.*: **A reduction in hippocampal GABA_A receptor $\alpha 5$ subunits disrupts the memory for location of objects in mice.** *Genes Brain Behav*. 2010; **9**(5): 478–88.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Engin E, Zarnowska ED, Benke D, *et al.*: **Tonic Inhibitory Control of Dentate Gyrus Granule Cells by $\alpha 5$ -Containing GABA_A Receptors Reduces Memory Interference.** *J Neurosci*. 2015; **35**(40): 13698–712.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Deprez F, Vogt F, Floriou-Servou A, *et al.*: **Partial inactivation of GABA_A receptors containing the $\alpha 5$ subunit affects the development of adult-born dentate gyrus granule cells.** *Eur J Neurosci*. 2016; **44**(5): 2258–71.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Dawson GR, Maubach KA, Collinson N, *et al.*: **An inverse agonist selective for**

- alpha5 subunit-containing GABA_A receptors enhances cognition. *J Pharmacol Exp Ther.* 2006; **316**(3): 1335–45.
[PubMed Abstract](#) | [Publisher Full Text](#)
24. Atack JR: **Preclinical and clinical pharmacology of the GABA_A receptor alpha5 subtype-selective inverse agonist alpha5IA.** *Pharmacol Ther.* 2010; **125**(1): 11–26.
[PubMed Abstract](#) | [Publisher Full Text](#)
 25. **F** Fischell J, van Dyke AM, Kvarta MD, *et al.*: **Rapid Antidepressant Action and Restoration of Excitatory Synaptic Strength After Chronic Stress by Negative Modulators of Alpha5-Containing GABA_A Receptors.** *Neuropsychopharmacology.* 2015; **40**(11): 2499–509.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 26. **F** Clarkson AN, Huang BS, Macisaac SE, *et al.*: **Reducing excessive GABA-mediated tonic inhibition promotes functional recovery after stroke.** *Nature.* 2010; **468**(7321): 305–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 27. **F** Koh MT, Rosenzweig-Lipson S, Gallagher M: **Selective GABA_A α5 positive allosteric modulators improve cognitive function in aged rats with memory impairment.** *Neuropharmacology.* 2013; **64**: 145–52.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 28. Savić MM, Huang S, Furtmüller R, *et al.*: **Are GABA_A receptors containing alpha5 subunits contributing to the sedative properties of benzodiazepine site agonists?** *Neuropsychopharmacology.* 2008; **33**(2): 332–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
 29. **F** Soto PL, Ator NA, Rallapalli SK, *et al.*: **Allosteric modulation of GABA_A receptor subtypes: effects on visual recognition and visuospatial working memory in rhesus monkeys [corrected].** *Neuropsychopharmacology.* 2013; **38**(11): 2315–25.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 30. Ballard TM, Knoflach F, Prinssen E, *et al.*: **RO4938581, a novel cognitive enhancer acting at GABA_A alpha5 subunit-containing receptors.** *Psychopharmacology (Berl.).* 2009; **202**(1–3): 207–23.
[PubMed Abstract](#) | [Publisher Full Text](#)
 31. Knust H, Achermann G, Ballard T, *et al.*: **The discovery and unique pharmacological profile of RO4938581 and RO4882224 as potent and selective GABA_A alpha5 inverse agonists for the treatment of cognitive dysfunction.** *Bioorg Med Chem Lett.* 2009; **19**(20): 5940–4.
[PubMed Abstract](#) | [Publisher Full Text](#)
 32. Gardiner K, Herault Y, Lott IT, *et al.*: **Down syndrome: from understanding the neurobiology to therapy.** *J Neurosci.* 2010; **30**(45): 14943–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 33. Gardiner KJ: **Molecular basis of pharmacotherapies for cognition in Down syndrome.** *Trends Pharmacol Sci.* 2010; **31**(2): 66–73.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 34. Ruparella A, Pearn ML, Mobley WC: **Cognitive and pharmacological insights from the Ts65Dn mouse model of Down syndrome.** *Curr Opin Neurobiol.* 2012; **22**(5): 880–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 35. Liu C, Belichenko PV, Zhang L, *et al.*: **Mouse models for Down syndrome-associated developmental cognitive disabilities.** *Dev Neurosci.* 2011; **33**(5): 404–13.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 36. Fernandez F, Morishita W, Zuniga E, *et al.*: **Pharmacotherapy for cognitive impairment in a mouse model of Down syndrome.** *Nat Neurosci.* 2007; **10**(4): 411–3.
[PubMed Abstract](#) | [Publisher Full Text](#)
 37. **F** Braudeau J, Delatour B, Duchon A, *et al.*: **Specific targeting of the GABA-A receptor α5 subtype by a selective inverse agonist restores cognitive deficits in Down syndrome mice.** *J Psychopharmacol.* 2011; **25**(8): 1030–42.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 38. Möhler H: **Cognitive enhancement by pharmacological and behavioral interventions: the murine Down syndrome model.** *Biochem Pharmacol.* 2012; **84**(8): 994–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
 39. **F** Martínez-Cué C, Martínez P, Rueda N, *et al.*: **Reducing GABA_A α5 receptor-mediated inhibition rescues functional and neuromorphological deficits in a mouse model of down syndrome.** *J Neurosci.* 2013; **33**(9): 3953–66.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 40. Martínez-Cué C, Delatour B, Potier M: **Treating enhanced GABAergic inhibition in Down syndrome: use of GABA α5-selective inverse agonists.** *Neurosci Biobehav Rev.* 2014; **46 Pt 2**: 218–27.
[PubMed Abstract](#) | [Publisher Full Text](#)
 41. **F** Murray JD, Anticevic A, Gancsos M, *et al.*: **Linking microcircuit dysfunction to cognitive impairment: effects of disinhibition associated with schizophrenia in a cortical working memory model.** *Cereb Cortex.* 2014; **24**(4): 859–72.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 42. Lewis DA, González-Burgos G: **Neuroplasticity of neocortical circuits in schizophrenia.** *Neuropsychopharmacology.* 2008; **33**(1): 141–65.
[PubMed Abstract](#) | [Publisher Full Text](#)
 43. **F** Chung DW, Fish KN, Lewis DA: **Pathological Basis for Deficient Excitatory Drive to Cortical Parvalbumin Interneurons in Schizophrenia.** *Am J Psychiatry.* 2016; **173**(11): 1131–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 44. Lisman JE, Coyle JT, Green RW, *et al.*: **Circuit-based framework for understanding neurotransmitter and risk gene interactions in schizophrenia.** *Trends Neurosci.* 2008; **31**(5): 234–42.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 45. Lewis DA, Cho RY, Carter CS, *et al.*: **Subunit-selective modulation of GABA type A receptor neurotransmission and cognition in schizophrenia.** *Am J Psychiatry.* 2008; **165**(12): 1585–93.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 46. Buchanan RW, Keefe RS, Lieberman JA, *et al.*: **A randomized clinical trial of MK-0777 for the treatment of cognitive impairments in people with schizophrenia.** *Biol Psychiatry.* 2011; **69**(5): 442–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 47. **F** Gill KM, Lodge DJ, Cook JM, *et al.*: **A novel α5GABA_A-positive allosteric modulator reverses hyperactivation of the dopamine system in the MAM model of schizophrenia.** *Neuropsychopharmacology.* 2011; **36**(9): 1903–11.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 48. **F** Gill KM, Cook JM, Poe MM, *et al.*: **Prior antipsychotic drug treatment prevents response to novel antipsychotic agent in the methylazoxymethanol acetate model of schizophrenia.** *Schizophr Bull.* 2014; **40**(2): 341–50.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 49. Spencer KM, Nestor PG, Perlmuter R, *et al.*: **Neural synchrony indexes disordered perception and cognition in schizophrenia.** *Proc Natl Acad Sci U S A.* 2004; **101**(49): 17288–93.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 50. Glykys J, Mann EO, Mody I: **Which GABA_A receptor subunits are necessary for tonic inhibition in the hippocampus?** *J Neurosci.* 2008; **28**(6): 1421–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
 51. Luscher B, Shen Q, Sahir N: **The GABAergic deficit hypothesis of major depressive disorder.** *Mol Psychiatry.* 2011; **16**(4): 383–406.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 52. **F** Luscher B, Fuchs T: **GABAergic control of depression-related brain states.** *Adv Pharmacol.* 2015; **73**: 97–144.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 53. **F** Piantadosi SC, French BJ, Poe MM, *et al.*: **Sex-Dependent Anti-Stress Effect of an α5 Subunit Containing GABA_A Receptor Positive Allosteric Modulator.** *Front Pharmacol.* 2016; **7**: 446.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 54. Botta P, Demmou L, Kasugai Y, *et al.*: **Regulating anxiety with extrasynaptic inhibition.** *Nat Neurosci.* 2015; **18**(10): 1493–500.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 55. **F** Behlke LM, Foster RA, Liu J, *et al.*: **A Pharmacogenetic 'Restriction-of-Function' Approach Reveals Evidence for Anxiolytic-Like Actions Mediated by α5-Containing GABA_A Receptors in Mice.** *Neuropsychopharmacology.* 2016; **41**(10): 2492–501.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 56. **F** Bakker A, Krauss GL, Albert MS, *et al.*: **Reduction of hippocampal hyperactivity improves cognition in amnesic mild cognitive impairment.** *Neuron.* 2012; **74**(3): 467–74.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 57. Chambers MS, Atack JR, Broughton HB, *et al.*: **Identification of a novel, selective GABA_A alpha5 receptor inverse agonist which enhances cognition.** *J Med Chem.* 2003; **46**(11): 2227–40.
[PubMed Abstract](#) | [Publisher Full Text](#)
 58. van Niel MB, Wilson K, Adkins CH, *et al.*: **A new pyridazine series of GABA_A alpha5 ligands.** *J Med Chem.* 2005; **48**(19): 6004–11.
[PubMed Abstract](#) | [Publisher Full Text](#)
 59. **F** Stamenić TT, Poe MM, Rehman S, *et al.*: **Ester to amide substitution improves selectivity, efficacy and kinetic behavior of a benzodiazepine positive modulator of GABA_A receptors containing the α5 subunit.** *Eur J Pharmacol.* 2016; **791**: 433–43.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 60. Rubenstein JL, Merzenich MM: **Model of autism: increased ratio of excitation/inhibition in key neural systems.** *Genes Brain Behav.* 2003; **2**(5): 255–67.
[PubMed Abstract](#) | [Publisher Full Text](#)
 61. Rubenstein JL: **Three hypotheses for developmental defects that may underlie some forms of autism spectrum disorder.** *Curr Opin Neurol.* 2010; **23**(2): 118–23.
[PubMed Abstract](#) | [Publisher Full Text](#)
 62. **F** Yizhar O, Fennel LE, Prigge M, *et al.*: **Neocortical excitation/inhibition balance in information processing and social dysfunction.** *Nature.* 2011; **477**(7363): 171–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 63. Fatemi SH, Reutiman TJ, Folsom TD, *et al.*: **mRNA and protein levels for GABA_A α4, α5, β1 and GABA_B R1 receptors are altered in brains from subjects with autism.** *J Autism Dev Disord.* 2010; **40**(6): 743–50.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 64. Gogolla N, Leblanc JJ, Quast KB, *et al.*: **Common circuit defect of excitatory-**

- inhibitory balance in mouse models of autism. *J Neurodev Disord.* 2009; **1**(2): 172–81.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
65. Rossignol E: **Genetics and function of neocortical GABAergic interneurons in neurodevelopmental disorders.** *Neural Plast.* 2011; **2011**: 649325.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
66.  Han S, Tai C, Jones CJ, *et al.*: **Enhancement of inhibitory neurotransmission by GABA_A receptors having $\alpha_{2,3}$ -subunits ameliorates behavioral deficits in a mouse model of autism.** *Neuron.* 2014; **81**(6): 1282–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
67. Li BM, Liu XR, Yi YH, *et al.*: **Autism in Dravet syndrome: prevalence, features, and relationship to the clinical characteristics of epilepsy and mental retardation.** *Epilepsy Behav.* 2011; **21**(3): 291–5.
[PubMed Abstract](#) | [Publisher Full Text](#)
68. Han S, Tai C, Westenbroek RE, *et al.*: **Autistic-like behaviour in *Scn1a*^{-/-} mice and rescue by enhanced GABA-mediated neurotransmission.** *Nature.* 2012; **489**(7416): 385–90.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
69. Chiron C, Marchand MC, Tran A, *et al.*: **Stiripentol in severe myoclonic epilepsy in infancy: a randomised placebo-controlled syndrome-dedicated trial. STICLO study group.** *Lancet.* 2000; **356**(9242): 1638–42.
[PubMed Abstract](#) | [Publisher Full Text](#)
70.  Clayton T, Poe MM, Rallapalli S, *et al.*: **A Review of the Updated Pharmacophore for the Alpha 5 GABA(A) Benzodiazepine Receptor Model.** *Int J Med Chem.* 2015; **2015**: 430248.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
71. Millan MJ, Agid Y, Brune M, *et al.*: **Cognitive dysfunction in psychiatric disorders: characteristics, causes and the quest for improved therapy.** *Nat Rev Drug Discov.* 2012; **11**(2): 141–68.
[PubMed Abstract](#) | [Publisher Full Text](#)

Open Peer Review

Current Referee Status:



Editorial Note on the Review Process

F1000 Faculty Reviews are commissioned from members of the prestigious F1000 Faculty and are edited as a service to readers. In order to make these reviews as comprehensive and accessible as possible, the referees provide input before publication and only the final, revised version is published. The referees who approved the final version are listed with their names and affiliations but without their reports on earlier versions (any comments will already have been addressed in the published version).

The referees who approved this article are:

Version 1

- 1 **Trevor Smart**, Department of Neuroscience, Physiology & Pharmacology, University College London, London, UK
Competing Interests: No competing interests were disclosed.
- 2 **James Rowlett**, Departments of Psychiatry & Human Behaviour and Neurobiology & Anatomical Sciences, Center for Psychiatric Neuroscience, University of Mississippi Medical Center, Jackson, Mississippi, USA
Competing Interests: No competing interests were disclosed.
- 3 **Bernhard Luscher**, ^{1,2,3} ¹ Department of Biology, Pennsylvania State University, Pennsylvania, USA
² Department of Biochemistry and Molecular Biology, Pennsylvania State University, Pennsylvania, USA
³ Center for Molecular Investigation of Neurological Disorders, Pennsylvania State University, Pennsylvania, USA
Competing Interests: No competing interests were disclosed.