



1236The Preventability of Ventilator-Associated Events: The CDC Prevention Epicenters' Wake Up and Breathe Collaborative

Citation

Klompas, M., D. Anderson, W. Trick, H. M. Babcock, R. Sinkowitz-Cochran, E. W. Ely, J. Jernigan, et al. 2014. "1236The Preventability of Ventilator-Associated Events: The CDC Prevention Epicenters' Wake Up and Breathe Collaborative." *Open Forum Infectious Diseases* 1 (Suppl 1): S46. doi:10.1093/ofid/ofu051.123. <http://dx.doi.org/10.1093/ofid/ofu051.123>.

Published version

<https://doi.org/10.1093/ofid/ofu051.123>

Link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:35981890>

Terms of use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material (LAA), as set forth at

<https://harvardwiki.atlassian.net/wiki/external/NGY5NDE4ZjgzNTc5NDQzMGIzZWZhMGFIOWI2M2EwYTg>

Accessibility

<https://accessibility.huit.harvard.edu/digital-accessibility-policy>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#)

ORAL ABSTRACTS

1236. The Preventability of Ventilator-Associated Events: The CDC Prevention Epicenters' Wake Up and Breathe Collaborative

Michael Klompas, MD, MPH, FRCPC, FIDSA¹; Deverick Anderson, MD, MPH²; William Trick, MD³; Hilary M. Babcock, MD, MPH⁴; Ronda Sinkowitz-Cochran, MPH⁵; E. Wesley Ely, MD, MPH⁶; John Jernigan, MD, MS⁷; Shelley S. Magill, MD, PhD⁸; Rosie D. Lyles, MD, MHA⁹; Caroline O'Neil, MPH⁹; Michele Balas, PhD, RN¹⁰; Michael V. Murphy, BA¹; Christopher Cox, MD¹¹; Ebbing Lautenbach, MD, MPH, MSCE¹²; Daniel J. Sexton, MD, FIDSA¹³; Victoria J. Fraser, MD, FIDSA, FSHEA⁴; Robert A. Weinstein, MD, FIDSA, FSHEA¹⁴; Richard Platt, MD, MS, FSHEA¹; ¹Department of Population Medicine, Harvard Medical School and Harvard Pilgrim Health Care Institute, Boston, MA; ²Duke University Medical Center, Durham, NC; ³Stroger Medical Center, Chicago, IL; ⁴Division of Infectious Diseases, Washington University School of Medicine, St. Louis, MO; ⁵Centers for Disease Control and Prevention, Atlanta, GA; ⁶Vanderbilt University, Nashville, TN; ⁷Division of Healthcare Quality Promotion, Centers for Disease Control and Prevention, Atlanta, GA; ⁸Stroger Hospital of Cook County, Chicago, IL; ⁹Washington University, St. Louis, MO; ¹⁰The Ohio State University, Columbus, OH; ¹¹Duke University, Durham, NC; ¹²University of Pennsylvania School of Medicine, Philadelphia, PA; ¹³Division of Infectious Diseases, Duke University Medical Center, Durham, NC; ¹⁴Internal Medicine, Section of Infectious Diseases, Cook County Health and Hospitals System, Chicago, IL

Session: 153. SHEA Featured Abstract
Friday, October 10, 2014: 4:30 PM

Background. The Centers for Disease Control and Prevention replaced its ventilator-associated pneumonia definitions with ventilator ventilator-associated event (VAE)

definitions in early 2013. Little is known about preventing VAEs. We hypothesized that coordinated, daily spontaneous awakening trials (SATs) and spontaneous breathing trials (SBTs) might prevent VAEs by decreasing patients' sedative and ventilator exposures.

Methods. We nested a multicenter quality improvement collaborative within a prospective study of VAE surveillance in 20 adult medical, surgical, medical-surgical, and cardiac intensive care units between November 2011 and May 2013. Twelve units joined the collaborative and implemented an opt-out protocol for nurses and respiratory therapists to initiate paired daily SATs and SBTs. The 8 remaining units conducted surveillance alone. The collaborative included two in-person meetings, monthly written reports, conference calls, and data feedback. We evaluated temporal trends in VAE rates and other outcomes per episode of mechanical ventilation using generalized mixed effects regression models adjusted for unit, age, sex, reason for intubation, SOFA score, and comorbidity index.

Results. We tracked 5,164 consecutive episodes of mechanical ventilation: 3,425 in collaborative units and 1,739 in surveillance-only units. Within collaborative units, there were significant increases in SATs, SBTs, and the percentage of SBTs performed without sedation. Mean duration of mechanical ventilation decreased by 2.4 days (95% CI 1.7-3.1), ICU length-of-stay by 3.0 days (95% CI, 1.6-4.3), and hospital length-of-stay by 6.3 days (95% CI, 4.0-8.6). There were significant decreases in ventilator-associated conditions (OR 0.63, 95% CI 0.42-0.97) and infection-related ventilator-associated complications (OR 0.35, 95% CI 0.17-0.71) but not Possible or Probable Pneumonia (OR 0.51, 95% CI 0.19-1.3). There was a significant increase in self-extubations (OR 2.1, 95% CI 1.1-3.9) but no change in re-intubations within 24 hours (OR 0.96, 95% CI 0.66-1.4). Within surveillance-only units, there were no significant changes in SAT, SBT, or VAE rates.

Conclusion. Nurse and respiratory therapist led opt-out protocols for paired, daily SATs and SBTs were associated with significant reductions in length-of-stay and VAEs.

Disclosures. E. W. Ely, Abbott: Speaker's Bureau, Speaker honorarium; Hospira: Speaker's Bureau, Speaker honorarium; Orion: Speaker's Bureau, Speaker honorarium; J. Jernigan, Sage Products: Conducting a clinical trial for which contributed product is being provided to participating hospitals, Contributed Product; Molnlycke: Conducting a clinical trial for which contributed product is being provided to participating hospitals, Contributed product; D. J. Sexton, UpToDate: Editor, Royalties; National Football League: Consultant, Consulting fee and Educational grant; Cubist: Grant Investigator, Grant recipient; Johnson and Johnson: Consultant, Consulting fee