



The brown anole dewlap revisited: do predation pressure, sexual selection, and species recognition shape among-population signal diversity?

Citation

Baeckens, Simon, Tess Driessens, and Raoul Van Damme. 2018. "The brown anole dewlap revisited: do predation pressure, sexual selection, and species recognition shape among-population signal diversity?" PeerJ 6 (1): e4722. doi:10.7717/peerj.4722. <http://dx.doi.org/10.7717/peerj.4722>

Published version

<https://doi.org/10.7717/peerj.4722>

Link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:37160428>

Terms of use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material (LAA), as set forth at

<https://harvardwiki.atlassian.net/wiki/external/NGY5NDE4ZjgzNTc5NDQzMGIzZWZhMGFIOWI2M2EwYTg>

Accessibility

<https://accessibility.huit.harvard.edu/digital-accessibility-policy>

Share Your Story

The Harvard community has made this article openly available. Please share how this access benefits you. [Submit a story](#)

The brown anole dewlap revisited: do predation pressure, sexual selection, and species recognition shape among-population signal diversity?

Simon Baeckens^{1,2,*}, Tess Driessens^{1,*} and Raoul Van Damme¹

¹ Laboratory of Functional Morphology, Department of Biology, University of Antwerp, Wilrijk, Belgium

² Museum of Comparative Zoology, Department of Organismic and Evolutionary Biology, Harvard University, Cambridge, MA, USA

* These authors contributed equally to this work.

ABSTRACT

Animal signalling structures are amongst the most variable characteristics, as they are subjected to a diversity of selection pressures. A well-known example of a diverse signalling system in the animal kingdom is the dewlap of *Anolis* lizards. Dewlap characteristics can vary remarkably among and within species, and also between sexes. Although a considerable amount of studies have attempted to disentangle the functional significance of the staggering dewlap diversity in *Anolis*, the underlying evolutionary processes remain elusive. In this study, we focus on the contribution of biotic selective pressures in shaping geographic variation in dewlap design (size, colour, and pattern) and dewlap display behaviour at the intraspecific level. Notably, we have tried to replicate and extend previously reported results hereof in both sexes of the brown anole lizard (*Anolis sagrei*). To do this, we assembled a dataset consisting of 17 *A. sagrei* heterogeneous island populations from the Caribbean and specifically tested whether predation pressure, sexual selection, or species recognition could explain interpopulational variation in an array of dewlap characteristics. Our findings show that in neither males nor females estimates of predation pressure (island size, tail break frequency, model attack rate, presence of predatory *Leiocephalus* lizards) or sexual selection (sexual size dimorphism) could explain variation in dewlap design. We did find that *A. sagrei* males from larger islands showed higher dewlap display intensities than males from smaller islands, but the direct connection with predation pressure remains ambiguous and demands further investigation. Last, we could show indirect support for species recognition only in males, as they are more likely to have a 'spotted' dewlap pattern when co-occurring with a higher number of syntopic *Anolis* species. In conclusion, we found overall limited support for the idea that the extensive interpopulational variability in dewlap design and use in *A. sagrei* is mediated by variation in their biotic environment. We propose a variety of conceptual and methodological explanations for this unexpected finding.

Submitted 13 March 2018

Accepted 16 April 2018

Published 8 May 2018

Corresponding author

Simon Baeckens,
simon.baeckens@uantwerp.be

Academic editor

David Roberts

Additional Information and
Declarations can be found on
page 15

DOI 10.7717/peerj.4722

© Copyright

2018 Baeckens et al.

Distributed under

Creative Commons CC-BY 4.0

OPEN ACCESS

Subjects Animal Behavior, Ecology, Evolutionary Studies, Zoology

Keywords *Anolis sagrei*, Caribbean islands, Clay models, Dewlap diversity, Display behaviour, Lizards, Predation pressure, Visual signalling system

INTRODUCTION

Even more than most other animal traits, signalling structures are subjected to a diversity of selection pressures (Johnstone, 1997; Smith & Harper, 2003). To be effective, they have to be clear and conspicuous, often under a variety of environmental conditions (Endler, 1992). A single signalling structure is often used to convey different messages to multiple receivers, and therefore must be capable of reaching several sensing systems (Loyau, Jalme & Sorci, 2005; Finkbeiner, Briscoe & Reed, 2014). At the same time, any form of transmission is prone to eavesdropping by predators or parasites, and thus signalling structures should not be *too* prominent (Roberts *et al.*, 2001; Clark, 2004; Bernal, Rand & Ryan, 2006). Adding to the complexity, as there are often multiple ways in which the same message (e.g. good genes) can be conveyed (e.g. by an acrobatic display, a bright-red crest, a particular odour), signalling structures also seem highly liable to the capriciousness of genetic drift (Richards-Zawacki, Yeager & Bart, 2013; Clark *et al.*, 2015) and mate choice (Møller & Pomiankowski, 1993). It should not come as a surprise, then, that signalling structures are amongst the most variable animal characteristics (Zuk & Tinghitella, 2008), and that understanding their evolution has proved particularly challenging.

The dewlap, an extendible flap of skin attached to the throat, of *Anolis* lizards is no exception. Typically male, but also female, anoles display their often brightly coloured dewlap in a variety of contexts and the resultant signal is said to function in social and sexual communication (Greenberg & Noble, 1944; Jenssen, 1970; Crews, 1975; Carpenter, 1978), in species recognition (Rand & Williams, 1970; Losos, 1985), and in predator deterrence (Leal & Rodríguez-Robles, 1995, 1997a, 1997b). Dewlaps can differ greatly in size, shape, colour, and patterning, among species, among populations within species, and between sexes (Nicholson, Harmon & Losos, 2007). In 2009, Vanhooydonck and colleagues studied differences in dewlap characteristics among seven island populations of the brown anole (*Anolis sagrei*) from the Bahamas. They reported that dewlap pattern and size have evolved under different selection regimes. Notably, their data showed that diversity in dewlap pattern is best explained by the number of syntopic *Anolis* species (thus, *species recognition* as the hypothesised driving force), whereas variation in relative dewlap size is primarily explained by the presence or absence of predatory *Leiocephalus* lizards (*natural selection*) and to some extent by sexual size dimorphism (*sexual selection*; in males only). Relative dewlap size in males and females appeared to be larger on islands where *A. sagrei* occurred in sympatry with *Leiocephalus* lizards. Based on this finding, the authors suggested that the *A. sagrei* dewlap functions at least partly as a pursuit-deterrence signal.

In this study, we have tried to replicate Vanhooydonck's results for both males and females using an extended dataset. To the data from the seven Bahamian islands (i.e. Acklins, Andros, Chub Cay, Crooked Island, Grand Bahama, Pidgeon Cay, Staniel Cay) reported in Vanhooydonck *et al.* (2009), we added *A. sagrei* populations from Cayman Brac, Cuba, Grand Cayman, Jamaica, Little Cayman, San Salvador, South Abaco, and South Bimini. Moreover, we measured two additional dewlap

characteristics (dewlap colour and use) that have been suggested to play a critical role in anole diversification and speciation ([Sigmund, 1983](#); [Losos, 1985](#); [Ord, Stamps & Losos, 2010](#); [Macedonia et al., 2013](#); [Ng et al., 2013, 2017](#)). Using similar proxies for quantifying selective regimes as Vanhooydonck and co-workers (i.e. island size, tail break frequency (TBF), model attack rate, presence of predatory *Leiocephalus* lizards, number of syntopic *Anolis* species, sexual size dimorphism), we here test whether interpopulational variation in *A. sagrei* dewlap characteristics (design and display) can be explained by predation, species recognition, and/or sexual selection hypotheses.

MATERIALS AND METHODS

Animals

We collected data on adult *A. sagrei* lizards from nine populations in the Caribbean during the breeding seasons of 2012, 2013, and 2015 (March–September; [Lee et al., 1989](#)). Data on one additional population (San Salvador) was collected outside the breeding season (January 2013). Details on the sampling locations of our populations are provided in [Fig. 1](#) and [Table S1](#). Individuals were caught by noose and kept individually in plastic bags for maximum 48 h before being released back at the exact location of capture. We measured snout–vent length (SVL) with callipers (Mitutoyo CD-15DC, accuracy 0.01 mm) and quantified dewlap design for each captured lizard. We caught a total of 282 males and 245 females (raw data for 143 males and 117 females were kindly provided by Bieke Vanhooydonck from the study by [Vanhooydonck et al., 2009](#)). To quantify dewlap use, we observed the behaviour of another 235 males and 189 females. All work was carried out in accordance with the University of Antwerp animal welfare standards and protocol (ECD 2011-64) and the local environmental agencies (The Bahamas Environment, Science & Technology Commission, The Ministry of The Environment and Housing; Department of Environment Cayman Islands; Centre for Marine Science, University of the West Indies Jamaica).

Dewlap measurements

Dewlap size

We used a technique outlined by [Vanhooydonck et al. \(2005\)](#) to estimate the surface area of the dewlap in every lizard caught. In short, we held the lizard on its left side against a 1 cm² gridded paper and gently pulled the base of the ceratobranchial forward with a pair of forceps until the dewlap was fully extended parallel to the grid. We then photographed the dewlap, using a Nikon D70 camera mounted on a tripod. We used Adobe Photoshop CS3 extended software (AP CS3, version 10.0) to trace the outer edge of the dewlap on the digital images and to calculate absolute dewlap area for all lizards (17 populations). To remove effects of overall size, we regressed log₁₀-transformed dewlap size against log₁₀-transformed SVL (for males and females separately). The obtained unstandardized residuals of all individuals were then averaged per population and used as estimate of relative dewlap size.




Figure 1 Sampling locations of the populations of study across the Caribbean. (1) Soroa (Cuba), population 1; (2) Soroa (Cuba) population 2; (3) Grand Cayman; (4) Santa Clara (Cuba); (5) South Bimini; (6) Chub Cay; (7) Andros; (8) Crooked Island; (9) Acklins; (10) San Salvador; (11) Staniel Cay; (12) Pidgeon Cay; (13) Grand Bahama; (14) South Abaco; (15) Cayman Brac; (16) Little Cayman; (17) Jamaica. [Full-size](#) DOI: 10.7717/peerj.4722/fig-1

Dewlap pattern

In most brown anoles, the alteration of red and yellow-coloured patches on the dewlap gives rise to a ‘pattern’ that can be categorised into three types (*Nicholson, Harmon & Losos, 2007; Driessens et al., 2015*). ‘Solid’ dewlaps are uniformly coloured; ‘marginal’ dewlaps have an evenly reddish coloured centre and a yellowish margin; and ‘spotted’ dewlaps have yellowish spots scattered across the reddish centre, regardless of the presence of a margin. One of us (T.D.) assigned each of the 425 male and 362 female dewlaps from the 17 study populations to one of the pattern categories on the basis of high-quality digital photos. We then determined the percentage of individuals attributed for the respective categories, per sex and per population.

Dewlap colour

We measured dewlap reflectance at the centre of the dewlap, using an Avantes spectrometer (AvaSpec-2048 USB2-UA-50; Avantes, Apeldoorn, the Netherlands, range 250–1000 nm) and deuterium-halogen light source (AvaLight-DHS; Avantes, Apeldoorn, the Netherlands) equipped with a fibre-optic probe. The probe was mounted within a metal holder to ensure readings at a fixed distance from the surface and was held perpendicular to the surface of the maximally extended dewlap. Reflectance data were

collected for wavelengths from 300 to 700 nm, including the lower range of photon absorption by UV-sensitive photoreceptor cones published for anoles ([Fleishman, Loew & Leal, 1993](#)). To investigate dewlap colour variation, we interpolated each spectrum to 1 nm wavelength intervals and extracted four variables following [Ng et al. \(2013\)](#): brightness, hue, and relative reflectance in UV (RF 365 nm) and in red (RF 655 nm) ([Montgomerie, 2006](#)). We calculated brightness as the total area under the uncorrected spectral curve (300–700 nm) ([Andersson, Örnborg & Andersson, 1998](#); [Smiseth et al., 2001](#)). For the remaining three colour variables, we corrected each spectrum for brightness by making the area under the curve equal to 1 ([Endler, 1990](#)). Hue was defined as the cut-on wavelength, i.e. the midpoint between baseline and maximum reflectance ([Andersson, Örnborg & Andersson, 1998](#); [Keyser & Hill, 2000](#); [Saks, McGraw & Horak, 2003](#); [Cummings, 2007](#)). We decided to extract relative reflectance specifically in UV (365 nm) and red (655 nm), as the *A. sagrei* dewlap spectrum shows maxima and a high level of intraspecific variation at both wavelengths ([Steffen & McGraw, 2007](#); T. Driessens, 2017, personal observation). Spectral measurements were carried out for 242 males and 217 females in total, distributed across nine populations. We do not have spectral data for the seven population sampled by [Vanhooydonck et al. \(2009\)](#), and for the population from Central Cuba (Santa Clara), due to technical problems with the spectrometer. All analyses of spectral data were run in R using the ‘pavo’ package ([Maia et al., 2013](#)).

Display behaviour

We observed each lizard ($N = 20\text{--}30$ males and $N = 8\text{--}25$ females per population; 10 populations in total) for 10 min in their natural environment, using a high definition camera (Sony, HDR-CX260VE). We first located lizards by walking silently through their natural habitat until an apparently undistributed individual was spotted. We then filmed the lizard from a distance using the camera zoom function in order to minimize any disturbance caused by our presence. Recordings were only made during sunny or partly clouded weather conditions and between 9 AM and 4:30 PM to avoid possible confounding effects of weather and time on the lizard’s activity level ([Huey, 1982](#); [Hertz, Huey & Stevenson, 1993](#)). All behavioural recordings were scored offline, using the software JWatcher 1.0 ([Blumstein, Evans & Daniel, 2000](#)). We calculated the dewlap extension (DE) rate per individual (‘DE rate’); that is, the number of times the dewlap was extended per minute. We then calculated the average DE per population. Moreover, we calculated the proportion of individuals per population that were observed dewlapping at least once during the 10 min observation window (‘prop. DE’). Calculations were performed for each sex separately. We combined these two measures for displaying intensity into a single measure by feeding them into a principal component analyses (for each sex separately). The analysis produced a single component that explained >90% of the variation in males and females, and was highly positive correlated with DE rate and proportion DE. We used this combined measure (‘PC dewlap display’) to index the intensity of dewlap displaying behaviour in males and females of our study populations.

Selection proxies

Predation pressure

As in *Vanhooydonck et al. (2009)*, we used island size, TBF, and the presence/absence of *Leiocephalus* lizards as proxies for predation pressure in the respective populations. Firstly, island size is a crude estimator of predation intensity, because larger islands tend to house larger numbers of predators, like raptors and snakes (*Losos & Schluter, 2000; Ricklefs & Bermingham, 2004*). Also, in the Bahamas, the survival rate of *A. sagrei* is substantially lower on larger islands with more bird species (*Schoener & Schoener, 1982*). Information on island area was obtained from the literature (*Losos, Irschick & Schoener, 1994; Vanhooydonck et al., 2009; Bradley & Rey-Millet, 2013*) or taken from websites (<http://islands.unep.ch> and <http://www.geographia.com>). Secondly, we concur with *Vanhooydonck et al. (2009)* and many other authors (*Schoener & Schoener, 1980; Turner et al., 1982; Fox, Perea-Fox & Franco, 1994; Bateman & Fleming, 2009; Donihue et al., 2016; Itescu et al., 2017a, 2017b*) that TBF can be a questionable measure of actual predation risk, and use it here for sake of conformity and in combination with other estimates of predation. This index was calculated as the ratio of the number of lizards with a regenerated tail to the total number of lizards captured for each population and separately per sex. Thirdly, the results of *Vanhooydonck et al. (2009)* suggest that the presence/absence of curly-tailed lizards (*Leiocephalus carinatus*) from their study islands in the Bahamas constitutes an important factor in the evolution of dewlap size. Curly-tailed lizards may exert their influence through competition for arthropod food or directly, by preying on anoles (*Schoener, Slade & Stinson, 1982; Schoener, Spiller & Losos, 2002; Losos, Schoener & Spiller, 2004; Losos et al., 2006; López-Darias, Schoener & Spiller, 2012; Lapiedra, Chejanovski & Kolbe, 2017*). For the smaller islands (<30 km²), we relied on data from the literature for deciding whether the *A. sagrei* populations were or were not syntopic with curly-tailed lizards. However, for the larger islands, we established a circle of 10 km radius (*Dean, Smith & Engeman, 2004*) around our study sites and considered them ‘under *Leiocephalus* pressure’ only when a curly-tailed lizard was seen within that area by us or by consulted local herpetologists. Finally, in addition to the three proxies of predation intensity used by *Vanhooydonck et al. (2009)*, we tallied the number of clay models of *Anolis* lizards attacked by predators. This technique has been used successfully to estimate predation rate (especially by birds) in other anole studies (*Brodie, 1993; Moore & Robinson, 2004; Husak et al., 2006; Steffen, 2009*). We first constructed hundreds of models by pouring brown-coloured clay (Plastalina, Claytoon, Valencia, Spain) in a mould made from an *A. sagrei* specimen (*Fig. 2A*). On location, we placed approximately 120 clay models per sampling locality with a distance of 4–6 m in between. Models were randomly distributed on natural perches for *A. sagrei* lizards (trunk-ground ecomorph; *Losos, 2009*), including trunks, branches, stones, and litter on the ground. After leaving the sampling site undisturbed for 48 h, we recollected the clay models and scored for predator marks (i.e. clear attacks of birds, lizards and/or rodents, *Fig. 2B*). The proportion of attacked models was calculated as the number of recollected attacked models divided by



Figure 2 Clay model of a brown anole used for estimating predation pressure. Photograph of a clay *Anolis* model used for estimating predation pressure: (A) an intact model; (B) a recaptured model with predation marks. Picture by Tess Driessens. [Full-size](#) DOI: [10.7717/peerj.4722/fig-2](https://doi.org/10.7717/peerj.4722/fig-2)

the total number of recaptured models, per population. Data on the proportion of attacked clay models could be collected for nine *A. sagrei* populations only.

Sexual selection

We used sexual size dimorphism (SSD) as a proxy for the intensity of sexual selection in each of our study populations. SSD has been widely used to gauge the strength of sexual selection in lizards, and anoles in particular, because there is strong evidence that a large body size increases a male's competitiveness and thereby its access to females (Ord, Blumstein & Evans, 2001; Stuart-Fox & Ord, 2004; Ord & Martins, 2006; and references therein). Following Vanhooydonck *et al.* (2009), we here defined SSD as mean SVL in males divided by mean SVL in females, per population.

Species recognition

To test the 'species recognition' hypothesis, we noted the presence of any other *Anolis* species within our sampling areas, as in Vanhooydonck *et al.* (2009).

Statistics

In this study, we considered interpopulational variation and therefore used population means and proportions of individuals per population as data points. We applied arcsine square root transformations to all proportion data (dewlap pattern, TBF, model attack rate) to meet normality assumptions (Sokal & Rohlf, 1995).

Our statistical analyses differed from the ones adopted by Vanhooydonck *et al.* (2009) in two important ways. Firstly, Vanhooydonck *et al.* (2009) adopted an information-theoretic approach to compare seven plausible models of selection for dewlap size and pattern. In doing so, they tested which combination of multiple predictor variables best describes the variation in dewlap characteristics. We are reluctant to take this approach because of the limited number of data points (seven populations in their case, nine to 17 in our extended dataset) relative to the number of predictors (see also Burnham & Anderson, 2002; Garamszegi, 2011). Rather, we opted for univariate regression analyses, linking individual predictor variables to individual dewlap characteristics. This allowed us to make full use of the information available for a particular pair of predictor and

response variable (not all variables could be measured in all populations). Second, in analysing the data here, we took phylogenetic relationships among the study populations into account. The phylogeny used in the comparative analyses is the one proposed by *Driessens et al. (2017)*, which is based on mtDNA haplotypes obtained by *Kolbe et al. (2004)*. Phylogenetic regression analyses were conducted using the `pgls()` function in the ‘caper’ package in R (*Revell, 2010; Orme et al., 2013*). This method uses maximum likelihood to simultaneously estimate the regression model and the phylogenetic signal (Pagel’s λ) of the residual error (*Garland & Ives, 2000; Revell, 2010*). It has been shown to do better than a priori tests of phylogenetic signal to estimate the appropriateness of phylogenetically corrected tests, especially when sample sizes are smaller than 20 (*Blomberg, Garland & Ives, 2003; Revell, 2010; Kamilar & Cooper, 2013*). Comparisons of dewlap characteristics between islands with and without *Leiocephalus* lizards were conducted using the `phy1ANOVA()` function in the ‘phytools’ package in R (*Revell, 2012*). Because data from one population (San Salvador) could only be obtained outside the breeding season (see Materials and Methods, section ‘Animals’), we have run all analyses with and without inclusion of this population. Results were nearly identical and we will, therefore, report results for the complete dataset only. Raw *P*-values were corrected for multiple testing using the Benjamini–Hochberg (BH) procedure (*Benjamini & Hochberg, 1995*). All statistical analyses were conducted in R version 3.2.1 (*R Development Core Team, 2016*).

RESULTS

The raw data on dewlap characteristics can be found on Dryad (<https://doi.org/10.5061/dryad.4572v>), and the data on island size, TBF, model attack rate, presence/absence of curly-tailed lizards, SSD, and number of *Anolis* species, is presented in [Table 1](#).

None of the four measures assumed to index the intensity of predation in our study populations correlated significantly with any of the dewlap design characteristics ([Table 2](#), all $P > 0.06$). Dewlap size, colour, and pattern also did not differ consistently between populations syntopic or not with the predatory curly-tailed lizard ([Table 2](#), all $P > 0.13$). These results suggest that interpopulational differences in dewlap design may not be driven by differences in predation intensity. We did find evidence for an effect of island size on dewlap display behaviour in males. Male anoles on larger islands scored higher on PC dewlap display, indicating that they used their dewlap more often than conspecifics on smaller islands ($P = 0.002$). All other proxies for predation pressure (TBF, model attack rate, presence/absence of curly-tailed lizards) did not have a comparable effect on male display intensity (all $P > 0.15$). Neither did we find any relationship between predation intensity and female display rate (all $P > 0.20$).

In neither males nor females, differences in SSD significantly contributed to interpopulational variation in dewlap characteristics ([Table 3](#), all $P > 0.11$). In populations exhibiting stronger size dimorphism, males nor females had dewlaps that were consistently larger, brighter, or of a different hue than in populations with limited SSD. Nor did they have dewlaps that reflected more in the UV or red region. We found also no evidence for a relationship between SSD and the proportion of different types of

Table 1 Data on selection indices.

Population	Island size	TBF		Model attack rate	<i>L. carinatus</i>	SSD (<i>N</i>)	Number of <i>Anolis</i> species	<i>Anolis</i> species (excl. <i>A. sagrei</i>)
		Males (<i>N</i>)	Females (<i>N</i>)					
Acklins	310.8	0.55 (10)	0.09 (12)	–	1	1.43 (22)	2	<i>A. carolinensis</i>
Andros	5,957	0.50 (23)	0.45 (18)	–	0	1.23 (41)	4	<i>A. angusticeps</i> , <i>A. carolinensis</i> , <i>A. distichus</i>
Cayman Brac	38	0.29 (28)	0.10 (29)	0.008	1	1.33 (57)	2	<i>A. maynardi</i>
Chub Cay	15.76	0.35 (20)	0.35 (16)	–	1	1.32 (36)	4	<i>A. angusticeps</i> , <i>A. carolinensis</i> , <i>A. distichus</i>
Crooked Island	238.28	0.46 (23)	0.48 (20)	–	1	1.25 (43)	2	<i>A. carolinensis</i>
Grand Bahama	1,373	0.56 (24)	0.38 (11)	–	1	1.33 (35)	3	<i>A. carolinensis</i> , <i>A. distichus</i>
Grand Cayman	197	0.26 (27)	0.10 (29)	0.070	1	1.28 (56)	2	<i>A. conspersus</i>
Jamaica	10,911	0.29 (32)	0.29 (23)	0.073	0	1.24 (55)	3	<i>A. lineatopus</i> , <i>A. grahami</i>
Little Cayman	28	0.59 (28)	0.43 (27)	0.034	1	1.29 (55)	2	<i>A. maynardi</i>
Pidgeon Cay	0.019	0.47 (16)	0.25 (8)	–	0	1.21 (24)	2	<i>A. carolinensis</i>
San Salvador	163	0.41 (27)	0.48 (14)	0.067	1	1.35 (41)	2	<i>A. distichus</i>
Santa Clara	105,006	0.67 (27)	0.58 (24)	0.020	0	1.33 (51)	2	<i>A. allisoni</i>
Soroa 1	105,006	0.42 (23)	0.38 (21)	–	0	1.24 (44)	3	<i>A. homolechis</i> , <i>A. porcatus</i>
Soroa 2	105,006	0.50 (22)	0.46 (24)	0.019	0	1.32 (46)	3	<i>A. homolechis</i> , <i>A. porcatus</i>
South Abaco	1,145.9	0.30 (26)	0.29 (21)	0.008	1	1.28 (47)	2	<i>A. smaragdinus</i>
South Bimini	10.36	0.44 (27)	0.36 (23)	0.000	1	1.30 (50)	4	<i>A. angusticeps</i> , <i>A. carolinensis</i> , <i>A. distichus</i>
Staniel Cay	5.18	0.37 (26)	0.33 (20)	–	0	1.32 (46)	3	<i>A. carolinensis</i> , <i>A. distichus</i>

Notes: Island size, tail break frequency (TBF) for males and females, proportion of attacked clay models, presence/absence of *Leiocephalus carinatus* lizards (0 = absent, 1 = present), sexual size dimorphism (SSD), and total number of co-occurring *Anolis* species; ‘–’ represents missing data. Sample sizes (*N*) used to calculate TBF and SSD are also provided.

dewlap patterns (i.e. solid, marginal, or spotted). Together, these findings do not suggest that differences in the intensity of sexual selection among the populations contribute to among-island variation in dewlap design.

Finally, we found no significant relationship between the number of co-occurring *Anolis* species and relative dewlap size, brightness, hue, or reflectance in UV or red regions (Table 3, all $P > 0.31$). We neither found an effect on dewlap display behaviour ($P > 0.58$). Interestingly, males—but not females—of populations with higher numbers of syntopic congeneric species were more likely to have a ‘spotted’ dewlap pattern ($P = 0.001$ in males).

Running standard (traditional) regression analyses without the incorporation of phylogenetic relationships revealed similar results.

DISCUSSION

Overall, we found limited support for the idea that the extensive among-population variability in dewlap characteristics in *A. sagrei* is mediated by variation in their biotic environment. SSD, an index of sexual selection, varied considerably among our study populations, but did not correlate with any of the structural aspects of the dewlap

Table 2 Data on predation pressure.

Predation pressure		Island size			Tail break frequency (TBF)			Model attack rate			<i>L. carinatus</i>					
Dewlap variables		<i>N</i>	<i>b</i>	<i>SE</i>	<i>P</i>	<i>N</i>	<i>b</i>	<i>SE</i>	<i>P</i>	<i>N</i>	<i>b</i>	<i>SE</i>	<i>P</i>	<i>N</i>	<i>F</i>	<i>P</i>
Males																
Relative dewlap size	17	3.7×10^{-7}		4.2×10^{-7}	0.392	17	0.241	0.131	0.086	9	-0.038	0.237	0.877	17	0.678	0.461
Pattern 'solid'	17	-1.8×10^{-6}		2.6×10^{-6}	0.498	17	0.757	0.569	0.204	9	1.684	1.009	0.139	17	0.686	0.457
Pattern 'marginal'	17	5.5×10^{-8}		2.5×10^{-6}	0.982	17	-0.587	0.525	0.281	9	1.475	1.411	0.331	17	0.004	0.961
Pattern 'spotted'	17	1.5×10^{-6}		2.4×10^{-6}	0.558	17	-0.104	0.539	0.850	9	-3.006	1.183	0.351	17	0.779	0.458
Colour brightness	9	1.4×10^{-2}		9.9×10^{-3}	0.197	9	-2405	4044	0.571	8	-3709	4880	0.476	9	0.422	0.575
Colour hue	9	-5.0×10^{-5}		4.2×10^{-5}	0.275	9	-14.88	12.08	0.257	8	-2.11	22.03	0.927	9	3.545	0.135
Colour RF365 nm	9	-3.6×10^{-9}		2.1×10^{-9}	0.135	9	-7.3×10^{-4}	6.0×10^{-4}	0.261	8	-1.3×10^{-4}	1.2×10^{-3}	0.917	9	2.979	0.137
Colour RF655 nm	9	2.2×10^{-9}		3.9×10^{-9}	0.592	9	9.6×10^{-4}	1.1×10^{-3}	0.409	8	1.1×10^{-3}	1.8×10^{-3}	0.571	9	3.088	0.147
PC dewlap display	10	1.8×10^{-5}		2.9×10^{-6}	0.002	10	2.587	1.656	0.157	9	-2.203	3.360	0.533	10	2.277	0.210
Females																
Relative dewlap size	17	4.7×10^{-7}		2.8×10^{-7}	0.114	17	-0.002	0.073	0.980	9	-0.254	0.153	0.140	17	0.018	0.920
Pattern 'solid'	17	-4.8×10^{-6}		3.0×10^{-6}	0.131	17	-0.946	0.548	0.105	9	-0.506	1.381	0.725	17	0.004	0.950
Pattern 'marginal'	17	5.4×10^{-6}		2.7×10^{-6}	0.062	17	1.135	0.461	0.176	9	1.693	1.251	0.218	17	0.106	0.746
Pattern 'spotted'	17	-1.9×10^{-7}		1.8×10^{-6}	0.916	17	0.253	0.437	0.571	9	-1.689	0.967	0.124	17	0.636	0.487
Colour brightness	9	6.1×10^{-3}		1.0×10^{-2}	0.577	9	-5349	2112	0.176	8	-4188	5199	0.451	9	0.322	0.633
Colour hue	9	-3.9×10^{-5}		4.9×10^{-5}	0.455	9	-3.61	13.18	0.792	8	18.18	25.05	0.495	9	3.421	0.113
Colour RF365 nm	9	-3.6×10^{-9}		2.0×10^{-9}	0.115	9	-6.7×10^{-4}	4.1×10^{-4}	0.146	8	-5.1×10^{-5}	1.1×10^{-3}	0.965	9	0.924	0.399
Colour RF655 nm	9	7.1×10^{-9}		2.8×10^{-9}	0.236	9	1.2×10^{-3}	6.6×10^{-4}	0.120	8	2.2×10^{-4}	1.8×10^{-3}	0.907	9	0.459	0.555
PC dewlap display	10	8.8×10^{-6}		6.4×10^{-6}	0.204	10	0.846	1.955	0.676	9	-2.285	4.065	0.591	10	0.040	0.877

Notes:

Univariate pgs regression analyses of dewlap design and display versus estimates of predation intensity (island size, tail break frequency, and model attack rate). Phylogenetic analyses of variance incorporating dewlap design and use versus presence or absence of predatory curly-tailed lizards. Results are shown separately per sex; *b* indicates the regression coefficient and *SE*, its standard error. Significant results (BH-corrected *P*-value) are in bold. See 'Statistics' section for more details.

Table 3 Sexual selection and species recognition.

Dewlap variables	Sexual selection				Species recognition			
	Sexual size dimorphism (SSD)				Number of <i>Anolis</i> species			
	<i>N</i>	<i>b</i>	<i>SE</i>	<i>P</i>	<i>N</i>	<i>b</i>	<i>SE</i>	<i>P</i>
Males								
Relative dewlap size	17	−0.014	0.323	0.966	17	0.011	0.022	0.632
Pattern ‘solid’	17	−0.937	1.178	0.439	17	−0.081	0.134	0.554
Pattern ‘marginal’	17	−0.250	1.094	0.822	17	−0.168	0.115	0.168
Pattern ‘spotted’	17	−0.984	1.035	0.357	17	0.471	0.094	0.001
Colour brightness	9	−1,0549	12,179	0.415	9	564.9	652.9	0.416
Colour hue	9	67.45	39.45	0.131	9	−2.552	2.617	0.362
Colour RF365 nm	9	0.001	0.002	0.701	9	1.1×10^{-4}	1.4×10^{-4}	0.473
Colour RF655 nm	9	−0.001	0.003	0.741	9	-2.2×10^{-4}	2.4×10^{-3}	0.388
PC dewlap display	10	8.327	5.290	0.154	10	0.243	0.421	0.580
Females								
Relative dewlap size	17	−0.026	0.211	0.907	17	0.015	0.015	0.349
Pattern ‘solid’	17	1.354	1.641	0.422	17	0.076	0.185	0.687
Pattern ‘marginal’	17	−0.677	1.696	0.696	17	0.089	0.171	0.609
Pattern ‘spotted’	17	−0.447	1.324	0.740	17	0.084	0.089	0.362
Colour brightness	9	−6,202	12,159	0.626	9	63.93	664.9	0.926
Colour hue	9	65.08	49.10	0.227	9	−3.250	3.000	0.314
Colour RF365 nm	9	−0.001	0.002	0.751	9	7.9×10^{-5}	1.4×10^{-4}	0.595
Colour RF655 nm	9	0.003	0.003	0.401	9	-6.0×10^{-5}	2.4×10^{-4}	0.811
PC dewlap display	10	3.462	8.755	0.703	10	−0.257	0.486	0.612

Notes:

Univariate ppls regression analyses of dewlap design and display versus sexual size dimorphism (SSD) and total number of co-occurring *Anolis* species. Results are shown separately per sex; *b* indicates the regression coefficient and *SE*, its standard error. Significant results (BH-corrected *P*-value) are in bold. See ‘Statistics’ section for more details.

considered, or with the intensity of displaying behaviour. Of the four proxies we used to assess relative predation intensity, none explained differences in dewlap design, and only one (i.e. island size) was associated with increased dewlap use. Our results do corroborate the hypothesis that the complexity of dewlap patterning functions in species recognition—at least in males. The relative size or colour characteristics of the dewlap, however, did not change consistently with the number of co-occurring congeneric species.

Vanhooydonck et al. (2009), using information on a subset of the populations studied here, concluded that predation pressure, especially the presence of predatory *L. carinatus* lizards, plays a significant role in the evolution of relative dewlap size. They argued that anoles may evolve larger dewlaps when in syntopy with these saurophagous lizards, because a large size would benefit the pursuit deterrence function of the dewlap. The predator deterrence hypothesis holds that prey performs eye-catching displays to warn the predator that its presence has been detected and that pursuits are likely to be futile or even dangerous (*Hasson, 1991*). It is a well-established fact that

many lizard species will indeed engage in conspicuous displaying behaviour when confronted with a predator (reviewed in [Greene, 1988](#)). However, in most cases, these displays involve tail vibrations, curling or waving (although not that common in anoles as in other lizard groups; [Dial, 1986](#); [Hasson, Hibbard & Ceballos, 1989](#); [Cooper, 2001, 2007, 2010, 2011](#); [Telemeco, Baird & Shine, 2011](#); [York & Bairds, 2016](#)), and they are more likely intended to deflect the predator's attack towards less vulnerable, expendable body parts, rather than to discourage pursuit. [Leal & Rodríguez-Robles \(1995, 1997a, 1997b\)](#) have argued that dewlapping in *A. cristatellus* and *A. cuvieri* may act as a pursuit deterrence signal, but the evidence is weak. In laboratory conditions, specimens of *A. cristatellus* were reported to dewlap 'only rarely' when a live native snake predator was introduced into their cage ([Leal & Rodríguez-Robles, 1995](#)); in the field, specimens of the same species did not extend their dewlaps more often when a snake model was moved into their territory ([Leal & Rodríguez-Robles, 1997a](#)). In their paper on *A. cuvieri*, [Leal & Rodríguez-Robles \(1997b\)](#) reported DEs in response to a snake model in only one individual out of a total of five. Moreover, several recent studies on *A. sagrei* found no evidence for increased DE rates in response to predatory birds ([Simon, 2007](#); [Elmasri et al., 2012](#)), snakes ([Yee et al., 2013](#)), or curly-tailed lizards ([Driessens, Vanhooydonck & Damme, 2014](#); [Steinberg et al., 2014](#)). For these reasons, we are sceptical about the role of the dewlap as a pursuit-deterrent and, hence, about predation pressure as a driver for dewlap size evolution. Accordingly, our analyses show very little evidence for a link between the used proxies of predation pressure and dewlap size—or any other structural aspect of the dewlap.

We are fully aware of the difficulty of measuring predation pressure. Each of the methods we employed has been criticized. Firstly, tail break frequencies may reflect predator inefficiency, rather than intensity ([Schoener & Schoener, 1980](#); [Turner et al., 1982](#); [Jaksic & Greene, 1984](#); [Fox, Perea-Fox & Franco, 1994](#)). Secondly, stationary clay models do not adequately mimic natural organisms with respect to traits such as odour, anti-predator postures, or movement ([Rangen, Clark & Hobson, 2000](#); [Thompson & Burhans, 2004](#); [Cooper, Caldwell & Vitt, 2008](#); [Santos & Cannatella, 2011](#); [Paluh, Hantak & Saporito, 2014](#)). Moreover, our model attack rates estimated especially predation by birds, but other predators like snakes and lizards can impose high predation threats as well ([Schoener, Slade & Stinson, 1982](#); [Henderson & Crother, 1989](#); [Rodríguez-Robles, 1992](#); [Rodríguez-Cabrera et al., 2016](#)). Thirdly, island size and the presence/absence of *L. carinatus* can be considered as very crude estimates of predator pressure—at best. On the other hand, some esteemed studies have established curly-tailed lizards as important drivers of morphological and behavioural diversification in *A. sagrei* ([Losos, Schoener & Spiller, 2004](#); [Losos et al., 2006](#); [Lapiedra, Chejanovski & Kolbe, 2017](#)). Lastly, prompting even more caution, none of the four respective proxies of predation intensity used in this study varied in concert (correlation analyses, all $R < 0.22$ and all $P > 0.24$). Perhaps the number of predatory species present on each island may provide more accurate information on the role of predation pressure in shaping dewlap design. Yet, greater species richness does not necessarily translate into higher predation rates as each predatory species may be less abundant or may include anoles as a smaller part of the diet ([Losos, 2009](#)). Combining the total number of predatory species with measures of their abundance and diet composition

might be most appropriate, but is hardly feasible in the field when surveying multiple sites in a short period of time. Admittedly, an accurate quantification of predation pressure in the field is very challenging. Our findings that predation pressure does not contribute to the evolution of dewlap design in the brown anole lizard across the islands included in our study remains therefore highly tentative and demands further research.

For dewlap displays, we did find that *A. sagrei* males from larger islands showed higher display intensities than males from smaller islands. Yet, part of this relationship might be driven by the artefact that the three populations sampled on Cuba (three times the same 'large' value for island size) exhibited high display rates. Besides, it is highly questionable whether the positive correlation between island size and male display rate is truly because larger islands harbour more predators. Island size is known to influence many ecologically relevant variables (e.g. habitat complexity, community richness; [Ricklefs & Lovette, 1999](#); [Losos & Schluter, 2000](#)) that were not considered here. Future studies are required to clarify and interpret our result showing that males use dewlaps more on large islands (i.e. non-independent island size data points from Cuba, islands size as accurate index of predation pressure).

The model that best explained the variation in male relative dewlap size in Vanhooydonck et al.'s dataset also contained sexual selection (SSD) as a predictor variable: on islands with high SSD (assumed to reflect high intensity of sexual selection), males tended to have larger dewlaps ([Vanhooydonck et al., 2009](#)). However, in our extended dataset, we found no indication that differences in SSD among islands are reflected in relative dewlap size, or any other dewlap characteristic. We can think of three ways to explain this result. First, sexual selection is simply not acting on dewlap traits. This sounds highly improbable, because the dewlap has all the characteristics of a sexually secondary trait: it is highly dimorphic in adult individuals, and exhibits the typical developmental pattern with sex-specific growth trajectories once the age of maturity is reached ([Vanhooydonck et al., 2015](#)). Several studies have shown that male brown anoles use their dewlap during territorial disputes ([Scott, 1984](#)) and/or during courtship ([Simon, 2011](#); [Driessens, Vanhooydonck & Damme, 2014](#)). At least in males, aspects of the dewlap reveal information on the individual signalling that is highly relevant in a sexual selection context ([Vanhooydonck et al., 2005](#); [Driessens et al., 2015](#)). Second, the effect of differences in the intensity of sexual selection among populations may be offset or overruled by some other factor. Natural selection may be counterbalancing or constraining any effects that divergent sexual selection is having on the among-island variation in dewlap characteristics (but see above). Dewlap traits may also be under differential selection for reasons not considered here. For instance, the climatic conditions and structural habitats in which our study populations live vary considerably. Physical aspects of the environment have been shown to influence the effectiveness of signals to a great extent in several animal species ([Hebets & Papaj, 2005](#); [Stuart-Fox, Moussalli & Whiting, 2007](#); [Baeckens et al., 2018](#)), including anoles ([Leal & Fleishman, 2004](#); [Ord et al., 2007](#); [Ng et al., 2013](#)). Moreover, [Driessens et al. \(2017\)](#) established that in *A. sagrei*, the dewlap of brown anoles occurring in 'xeric' environments differ from those inhabiting 'mesic' environment in dewlap colour, pattern, and use. They argue that the strong

relationship between signal design and prevailing environmental conditions might result from differential selection on signal efficacy. A third explanation for the apparent lack of a relationship between sexual selection and dewlap design is methodological: SSD may simply not be a good proxy for the intensity of sexual selection. Although male-biased SSD in lizards is traditionally linked to intrasexual selection (*Trivers, 1976; Stamps, 1983, 1999; Stamps, Losos & Andrews, 1997; Ord, Blumstein & Evans, 2001*), evolutionary shifts in male aggression, territoriality and (relative) home range size explain but a small proportion of evolutionary changes in SSD (*Cox, Skelly & John-Alder, 2003*), suggesting that other factors may be at play. Recently, *Bonneaud et al. (2016)* presented evidence that sex-specific developmental plasticity may contribute to adult SSD in brown anoles: on six Bahamian islands, the amount of food biomass explained variation in male, but not female body size, giving rise to significant differences in SSD. Clearly, future studies should invest in collecting information on more reliable estimates of the intensity of sexual selection.

Our analyses lend partial support to the finding that dewlap ‘patterning’ may play a role in species recognition. Male brown anoles are more likely to have a ‘spotted’ dewlap when co-occurring with several other *Anolis* species. Whereas *Vanhooydonck et al. (2009)* found this to be true in both sexes, the effect was only significant in males in our extended dataset. Perhaps males that can broadcast their species identity properly are less likely to be attacked by non-conspecific males. Indeed, male anoles can behave very aggressively towards conspecific males, while at the same time ignoring males of other *Anolis* species (*Losos, 1985*). Females, on the other hand, may communicate their species identity in some other way (perhaps via head bobbing patterns).

Rand & Williams (1970) coined the species recognition hypothesis for explaining dewlap diversity in *Anolis* five decades ago. Subsequent behavioural studies on pairs of anole species have offered further support to the hypothesis (*Losos, 1985; Macedonia & Stamps, 1994; Macedonia et al., 2013*), but a broad-scale, phylogenetically informed analysis could not provide statistical corroboration (*Nicholson, Harmon & Losos, 2007*). In this study, we only hold qualitative information based on personal observations and data from the literature to speculate on how the species recognition hypothesis might explain the success of a ‘spotted’ dewlap pattern in *A. sagrei* males when co-occurring with a high number of other *Anolis* species. As the species recognition hypothesis predicts that sympatric species will have different dewlap configurations (*Losos & Chu, 1998*), we expect the spotted dewlap pattern of *A. sagrei* to be unique in places where many *Anolis* species occur. Indeed, the dewlap patterning of the majority of syntopic species we observed is described as solid (*A. allisoni, A. angusticeps, A. carolinensis, A. conspersus, A. homolechis, A. maynardi, A. porcatius*) or marginal (*A. grahami*) (*Nicholson, Harmon & Losos, 2007; Vanhooydonck et al., 2009*), which strengthens the idea that a spotted dewlap might function as a species recognition signal in *A. sagrei*. Two other syntopic species (*A. distichus* and *A. lineatopus*), however, are known to vary in their dewlap patterning among populations (*Nicholson, Harmon & Losos, 2007; Ng et al., 2013*), and we do not have the data to confirm whether they have a non-spotted dewlap when co-occurring with *A. sagrei*. The brown anole seems clearly an interesting study animal

to deeper explore the possibility of sexual character displacement, but this will require detailed data on the signals of the co-occurring species and behavioural experiments.

CONCLUSION

Using a comparative approach, and based on estimates of predation pressure, sexual selection, and species recognition, we find limited evidence for the hypothesis that the vast among-population variability in dewlap characteristics in *A. sagrei* is driven by variation in their biotic environment. In this work, we offer a number of explanations for this unexpected finding and stress that this outcome might also be (partially) attributed to our study design, in specific, the choice and number of study islands. The initial study by [Vanhooydonck et al. \(2009\)](#) focused on a (although small) set of relatively homogeneous Bahamian islands (i.e. small variation in biotic features, such as, island size, number of conspecifics, predation pressure); in contrast, our extended dataset includes some ‘mega-islands’ (such as Cuba and Jamaica) and non-native populations (Grand Cayman, Jamaica; [Kolbe et al., 2004](#)), which might obscure clear patterns of dewlap variation. Moreover, although our comprehensive dataset encompasses over 600 lizards from 17 different populations, an increase in sample size would have strengthened our findings. Although sample size affects science in general, if correlations are weak or non-significant (as many in this study), regression analyses typically demand high sample sizes to reach high statistical power. Therefore, we encourage further research on the role of the biotic environment in shaping the intraspecific diversity in dewlap design. Ideally, comparative studies should include more similar-sized island populations from the brown anole’s native range (e.g. only focussing on islets in the Bahamas), or scholars should opt for an experimental approach with a controlled and replicated study design in natural populations (as in, e.g. [Losos, Schoener & Spiller, 2004](#); [Calsbeek & Cox, 2010](#)).

ACKNOWLEDGEMENTS

We acknowledge B. Vanhooydonck and co-workers for sharing the raw data of seven study populations included in this study. We further thank S. De Decker, J. Harvey, A. Herrel, J. Husak, P. Maillis, J. Mertens, D. Norris, V. Rivalta, L. Schettino, E. Schramme, B. Scott-Edwards, M. Valley, L. Vandervorst, and B. Vanhooydonck for help during data collection of the other 10 study populations. Thanks go to S. Van Dongen for statistical assistance. Finally, the authors thank the three reviewers for their constructive feedback on a previous draft of the manuscript.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

This research was financed by the Research Fund Flanders (FWO) awarded to Tess Driessens. Simon Baeckens is a postdoctoral fellow supported by the Belgian American Education Foundation (B.A.E.F.). Additional expenses for field missions were provided by Leopold III

fund and the University of Antwerp (DOCOP). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:

Research Fund Flanders (FWO).

Belgian American Education Foundation (B.A.E.F.).

Leopold III.

University of Antwerp (DOCOP).

Competing Interests

The authors declare that they have no competing interests.

Author Contributions

- Simon Baeckens performed the experiments, prepared figures and/or tables, authored or reviewed drafts of the paper, approved the final draft.
- Tess Driessens conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, approved the final draft.
- Raoul Van Damme conceived and designed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, approved the final draft.

Animal Ethics

The following information was supplied relating to ethical approvals (i.e. approving body and any reference numbers):

The Ethical Committee for Animal Testing at the University of Antwerp approved protocols for the use of live animals in this study.

Field Study Permissions

The following information was supplied relating to field study approvals (i.e. approving body and any reference numbers):

Field experiments were approved by the local environmental agencies (The Bahamas Environment, Science & Technology Commission; The Ministry of The Environment and Housing; Department of Environment Cayman Islands; Centre for Marine Science, University of the West Indies Jamaica).

Data Availability

The following information was supplied regarding data availability:

The raw data are included in [Tables 1–3](#).

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.4722#supplemental-information>.

REFERENCES

- Andersson SJ, Örnborg J, Andersson M. 1998. Ultraviolet sexual dimorphism and assortative mating in blue tits. *Proceedings of the Royal Society B: Biological Sciences* 265(1395):445–450 DOI 10.1098/rspb.1998.0315.
- Baeckens S, Martín J, Garcia-Roa R, Pafilis P, Huyghe K, Van Damme R. 2018. Environmental conditions shape the chemical signal design of lizards. *Functional Ecology* 32(2):566–580 DOI 10.1111/1365-2435.12984.
- Bateman P, Fleming P. 2009. To cut a long tail short: a review of lizard caudal autotomy studies carried out over the last 20 years. *Journal of Zoology* 277(1):1–14 DOI 10.1111/j.1469-7998.2008.00484.x.
- Benjamini Y, Hochberg Y. 1995. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society Series B: Statistical Methodology* 57(1):289–300.
- Bernal XE, Rand AS, Ryan MJ. 2006. Acoustic preferences and localization performance of blood-sucking flies (*Corethrella coquillett*) to tungara frog calls. *Behavioral Ecology* 17(5):709–715 DOI 10.1093/beheco/arl003.
- Blomberg SP, Garland T, Ives AR. 2003. Testing for phylogenetic signal in comparative data: behavioural traits are more labile. *Evolution* 57(4):717–745 DOI 10.1111/j.0014-3820.2003.tb00285.x.
- Blumstein DT, Evans CS, Daniel JC. 2000. JWatcher V1.0. An introductory user's guide. Available at <http://www.jwatcher.ucla.edu> (accessed March 2012).
- Bonneaud C, Marnocha E, Herrel A, Vanhooydonck B, Irschick DJ, Smith TB. 2016. Developmental plasticity affects sexual size dimorphism in an anole lizard. *Functional Ecology* 30(2):235–243 DOI 10.1111/1365-2435.12468.
- Bradley BR, Rey-Millet Y. 2013. *A Photographic Guide to the Birds of the Cayman Islands*. London: Bloomsbury Publishing Plc.
- Brodie ED III. 1993. Differential avoidance of coral snake banded patterns by free ranging avian predators in Costa Rica. *Evolution* 47(1):227–235 DOI 10.1111/j.1558-5646.1993.tb01212.x.
- Burnham KP, Anderson DR. 2002. *Model Selection and Multimodel Inference: A Practical Information-Theoretic Approach*. New York: Springer.
- Calsbeek R, Cox RM. 2010. Experimentally assessing the relative importance of predation and competition as agents of selection. *Nature* 465(7298):613–616 DOI 10.1038/nature09020.
- Carpenter CC. 1978. Ritualistic social behavior in lizard. In: Greenberg N, MacLean PD, eds. *Behavior and Neurology of Lizards*. Rockville: NIMH, 253–257.
- Clark RW. 2004. Timber rattlesnakes (*Crotalus horridus*) use chemical cues to select ambush sites. *Journal of Chemical Ecology* 30(3):607–617 DOI 10.1023/b:joec.0000018632.27010.1e.
- Clark DL, Macedonia JM, Rowe JW, Stuart MA, Kemp DJ, Ord TJ. 2015. Evolution of displays in Galapagos lava lizards: comparative analyses of signallers and robot playbacks to receivers. *Animal Behaviour* 109:33–44 DOI 10.1016/j.anbehav.2015.08.002.
- Cooper WE. 2001. Multiple roles of tail display by the curly-tailed lizard *Leiocephalus carinatus*: Pursuit deterrent and deflective roles of a social signal. *Ethology* 107(12):1137–1149 DOI 10.1046/j.1439-0310.2001.00754.x.
- Cooper WE. 2007. Escape and its relationship to pursuit-deterrent signalling in the Cuban curly-tailed lizard *Leiocephalus carinatus*. *Herpetologica* 63(2):144–150 DOI 10.1655/0018-0831(2007)63[144:eairtp]2.0.co;2.

- Cooper WE. 2010.** Timing during predator-prey encounters, duration and directedness of a putative pursuit-deterrent signal by the zebra-tailed lizard, *Callisaurus draconoides*. *Behaviour* **147**(13):1675–1691 DOI [10.1163/000579510x528215](https://doi.org/10.1163/000579510x528215).
- Cooper WE Jr. 2011.** Pursuit deterrence, predation risk, and escape in the lizard *Callisaurus draconoides*. *Behavioral Ecology and Sociobiology* **65**(9):1833–1841 DOI [10.1007/s00265-011-1191-5](https://doi.org/10.1007/s00265-011-1191-5).
- Cooper WE Jr, Caldwell JP, Vitt LJ. 2008.** Conspicuousness and vestigial escape behavior by two dendrobatid frogs, *Dendrobates auratus* and *Oophaga pumilio*. *Behaviour* **146**(3):325–349 DOI [10.1163/156853909x410946](https://doi.org/10.1163/156853909x410946).
- Cox RM, Skelly SL, John-Alder HB. 2003.** A comparative test of adaptive hypotheses for sexual size dimorphism in lizards. *Evolution* **57**(7):1653–1669 DOI [10.1111/j.0014-3820.2003.tb00371.x](https://doi.org/10.1111/j.0014-3820.2003.tb00371.x).
- Crews D. 1975.** Effects of different components of male courtship behavior on environmentally induced ovarian recrudescence and mating preferences in the lizard *Anolis carolinensis*. *Animal Behaviour* **23**:349–356 DOI [10.1016/0003-3472\(75\)90083-4](https://doi.org/10.1016/0003-3472(75)90083-4).
- Cummings ME. 2007.** Sensory trade-offs predict signal divergence in surfperch. *Evolution* **61**(3):530–545 DOI [10.1111/j.1558-5646.2007.00047.x](https://doi.org/10.1111/j.1558-5646.2007.00047.x).
- Dean CL, Smith HT, Engeman RM. 2004.** Geographic distribution: *Leiocephalus carinatus armouri* (Little Bahama curly-tailed lizard). *Herpetological Review* **35**:82.
- Dial BE. 1986.** Tail display in two species of iguanid lizards—a test of the predator signal hypothesis. *American Naturalist* **127**(1):103–111 DOI [10.1086/284471](https://doi.org/10.1086/284471).
- Donihue CM, Brock KM, Foufopoulos J, Herrel A. 2016.** Feed or fight: testing the impact of food availability and intraspecific aggression on the functional ecology of an island lizard. *Functional Ecology* **30**(4):566–575 DOI [10.1111/1365-2435.12550](https://doi.org/10.1111/1365-2435.12550).
- Driessens T, Baeckens S, Balzarolo M, Vanhooydonck B, Huyghe K, Van Damme R. 2017.** Climate-related environmental variation in a visual signalling device: the male and female dewlap in *Anolis sagrei* lizards. *Journal of Evolutionary Biology* **30**(10):1846–1861 DOI [10.1111/jeb.13144](https://doi.org/10.1111/jeb.13144).
- Driessens T, Huyghe K, Vanhooydonck B, Van Damme R. 2015.** Messages conveyed by assorted facets of the dewlap, in both sexes of *Anolis sagrei*. *Behavioral Ecology and Sociobiology* **69**(8):1251–1264 DOI [10.1007/s00265-015-1938-5](https://doi.org/10.1007/s00265-015-1938-5).
- Driessens T, Vanhooydonck B, Damme R. 2014.** Deterring predators, daunting opponents or drawing partners? Signaling rates across diverse contexts in the lizard *Anolis sagrei*. *Behavioral Ecology and Sociobiology* **68**(2):173–184 DOI [10.1007/s00265-013-1669-4](https://doi.org/10.1007/s00265-013-1669-4).
- Elmasri OL, Moreno MS, Neumann CA, Blumstein DT. 2012.** Response of brown anoles (*Anolis sagrei*) to multimodal signals from a native and novel predator. *Current Zoology* **58**(6):791–796 DOI [10.1093/czoolo/58.6.791](https://doi.org/10.1093/czoolo/58.6.791).
- Endler JA. 1990.** On the measurements and classification of color in studies of animal color pattern. *Biological Journal of the Linnean Society* **41**(4):315–352 DOI [10.1111/j.1095-8312.1990.tb00839.x](https://doi.org/10.1111/j.1095-8312.1990.tb00839.x).
- Endler JA. 1992.** Signals, signal conditions, and the direction of evolution. *American Naturalist* **139**:S125–S153 DOI [10.1086/285308](https://doi.org/10.1086/285308).
- Finkbeiner SD, Briscoe AD, Reed RD. 2014.** Warning signals are seductive: relative contributions of color and pattern to predator avoidance and mate attraction in *Heliconius* butterflies. *Evolution* **68**(12):3410–3420 DOI [10.1111/evo.12524](https://doi.org/10.1111/evo.12524).
- Fleishman LJ, Loew ER, Leal M. 1993.** Ultraviolet vision in lizards. *Nature* **365**(6445):397 DOI [10.1038/365397a0](https://doi.org/10.1038/365397a0).

- Fox SF, Perea-Fox S, Franco RC. 1994. Development of the tail autotomy adaptation in lizards under disparate levels of predation at high and low elevations in Mexico. *Southwestern Naturalist* 39:311–322.
- Garamszegi LZ. 2011. Information-theoretic approaches to statistical analyses in behavioural ecology: an introduction. *Behavioral Ecology and Sociobiology* 65(1):1–11 DOI 10.1007/s00265-010-1028-7.
- Garland T Jr, Ives AR. 2000. Using the past to predict the present: confidence intervals for regression equations in phylogenetic comparative methods. *American Naturalist* 155(3):346–364 DOI 10.1086/303327.
- Greenberg G, Noble GK. 1944. Social behavior of the American chameleon (*Anolis carolinensis* Voigt). *Physiological Zoology* 17(4):392–439 DOI 10.1086/physzool.17.4.30151738.
- Greene HW. 1988. Antipredator mechanisms in reptiles. In: Gans C, Huey RB, eds. *Biology of the Reptilia*. 16. New York: John Wiley & Sons, 1–152.
- Hasson O. 1991. Pursuit-deterrent signals: communication between prey and predator. *Trends in Ecology & Evolution* 6(10):325–329 DOI 10.1016/0169-5347(91)90040-5.
- Hasson O, Hibbard R, Ceballos G. 1989. The pursuit deterrent function of tail-wagging in the zebra-tailed lizard (*Callisaurus draconoides*). *Canadian Journal of Zoology* 67(5):1205–1209 DOI 10.1139/z89-174.
- Hebets EA, Papaj DR. 2005. Complex signal function: developing a framework of testable hypothesis. *Behavioral Ecology and Sociobiology* 57(3):197–214 DOI 10.1007/s00265-004-0865-7.
- Henderson RW, Crother B. 1989. Biogeographic patterns of predation in West Indian colubrid snakes. In: Woods CA, ed. *Biogeography of the West Indies Past, Present, and Future*. Gainesville: Sandhill Crane Press, Inc., 479–518.
- Hertz PE, Huey RB, Stevenson RD. 1993. Evaluating temperature regulation by field-active ectotherms—the fallacy of the inappropriate question. *American Naturalist* 142(5):796–818 DOI 10.1086/285573.
- Huey RB. 1982. Temperature, physiology, and the ecology of reptiles. In: Gans C, Pough FH, eds. *Biology of the Reptilia*. New York: Academic Press, 25–91.
- Husak JF, Macedonia JM, Fox SF, Saucedo RC. 2006. Predation cost of conspicuous male coloration in collared lizards (*Crotaphytus collaris*): an experimental test using clay-covered model lizards. *Ethology* 112(6):572–580 DOI 10.1111/j.1439-0310.2005.01189.x.
- Itescu Y, Schwarz R, Meiri S, Pafilis P. 2017a. Intraspecific competition, not predation, drives lizard tail loss on islands. *Journal of Animal Ecology* 86(1):66–74 DOI 10.1111/1365-2656.12591.
- Itescu Y, Schwarz R, Meiri S, Pafilis P. 2017b. Lizard tail-loss rates on islands are not governed by longer life spans. *Israel Journal of Ecology and Evolution* 63(2):53–56 DOI 10.1163/22244662-06301011.
- Jaksic FM, Greene ER. 1984. Correlates of tail losses in twelve species of *Liolaemus* lizards. *Journal of Herpetology* 14(2):137–141 DOI 10.2307/1563844.
- Jenssen TA. 1970. The ethoecology of *Anolis nebulosus*. *Journal of Herpetology* 4(1/2):1–38 DOI 10.2307/1562700.
- Johnstone RA. 1997. The evolution of animal signals. In: Krebs JR, Davies NB, eds. *Behavioural Ecology: An Evolutionary Approach*. Oxford: Blackwell Publishing, 155–178.
- Kamilar JM, Cooper N. 2013. Phylogenetic signal in primate behavior, ecology and life history. *Philosophical Transactions of the Royal Society B: Biological Sciences* 368(1618):20120341 DOI 10.1098/rstb.2012.0341.

- Keyser AJ, Hill G. 2000. Structurally based plumage coloration is an honest signal of quality in male blue grosbeaks. *Behavioral Ecology* 11(2):202–209 DOI 10.1093/beheco/11.2.202.
- Kolbe JJ, Glor RE, Schettino LR, Lara AC, Larson A, Losos JB. 2004. Genetic variation increases during biological invasion by a Cuban lizard. *Nature* 431(7005):177–181 DOI 10.1038/nature02807.
- Leal M, Fleishman LJ. 2004. Differences in visual signal design and detectability between allopatric populations of *Anolis* lizards. *American Naturalist* 163(1):26–39 DOI 10.1086/379794.
- Leal M, Rodríguez-Robles JA. 1995. Antipredator responses of *Anolis cristatellus* (Sauria: Polychrotidae). *Copeia* 1995(1):155–161 DOI 10.2307/1446810.
- Leal M, Rodríguez-Robles JA. 1997a. Signalling displays during predator–prey interactions in a Puerto Rican anole, *Anolis cristatellus*. *Animal Behaviour* 54(5):1147–1154 DOI 10.1006/anbe.1997.0572.
- Leal M, Rodríguez-Robles JA. 1997b. Antipredator responses of the Puerto Rican giant anole, *Anolis cuvieri* (Squamata:Polychrotidae). *Biotropica* 29(3):372–375 DOI 10.1111/j.1744-7429.1997.tb00440.x.
- Lapiedra O, Chejanovski Z, Kolbe JJ. 2017. Urbanization and biological invasion shape animal personalities. *Global Change Biology* 23(2):592–603 DOI 10.1111/gcb.13395.
- Lee JC, Clayton D, Eisenstein S, Perez I. 1989. The reproductive cycle of *Anolis sagrei* in southern Florida. *Copeia* 1989(4):930–937 DOI 10.2307/1445979.
- López-Darias M, Schoener T, Spiller DA. 2012. Predators determine how weather affects the spatial niche of lizard prey: exploring niche dynamics at a fine scale. *Ecology* 93(12):2512–2518 DOI 10.1890/12-0483.1.
- Losos JB. 1985. An experimental demonstration of the species-recognition role of *Anolis* dewlap color. *Copeia* 1985(4):905–910 DOI 10.2307/1445240.
- Losos JB. 2009. *Lizards in an Evolutionary Tree: Ecology and Adaptive Radiation of Anoles*. Berkeley: University of California Press.
- Losos JB, Chu L. 1998. Examination of factors potentially affecting dewlap size in Caribbean anoles. *Copeia* 1998(2):430–438 DOI 10.2307/1447437.
- Losos JB, Irschick DJ, Schoener TW. 1994. Adaptation and constraint in the evolution of specialization of Bahamian *Anolis* lizard. *Evolution* 48(6):1786–1789 DOI 10.2307/2410508.
- Losos JB, Schluter D. 2000. Analyses of an evolutionary species-area relationship. *Nature* 408(6814):847–850 DOI 10.1038/35048558.
- Losos JB, Schoener TW, Langerhans RB, Spiller DA. 2006. Rapid temporal reversal in predator-driven natural selection. *Science* 314(5802):1111 DOI 10.1126/science.1133584.
- Losos JB, Schoener TW, Spiller D. 2004. Predator-induced behaviour shifts and natural selection in field-experimental lizard populations. *Nature* 432(7016):505–508 DOI 10.1038/nature03039.
- Loyau A, Jalme MS, Sorci G. 2005. Intra- and intersexual selection for multiple traits in the peacock (*Pavo cristatus*). *Ethology* 111(9):810–820 DOI 10.1111/j.1439-0310.2005.01091.x.
- Macedonia JM, Clark DL, Riley RG, Kemp DJ. 2013. Species recognition of color and motion signals in *Anolis grahami*: evidence from responses to lizard robots. *Behavioral Ecology* 24(4):846–852 DOI 10.1093/beheco/art027.
- Macedonia JM, Stamps JA. 1994. Species recognition in *Anolis grahami* (Sauria, Iguanidae): evidence from responses to video playbacks of conspecific and heterospecific displays. *Ethology* 98(3–4):246–264 DOI 10.1111/j.1439-0310.1994.tb01074.x.
- Maia R, Eliason CM, Bitton PP, Doucet SM, Shawkey MD. 2013. Pavo: an R package for the analysis, visualization and organization of spectral data. *Methods in Ecology and Evolution* 4(10):906–913 DOI 10.1111/2041-210x.12069.

- Møller AP, Pomiankowski A. 1993. Why have birds got multiple sexual ornaments? *Behavioral Ecology and Sociobiology* 32(3):167–176 DOI 10.1007/BF00173774.
- Montgomerie R. 2006. Analyzing colors. In: Hill GE, McGraw KJ, eds. *Bird Coloration: Mechanics and Measurements*. Vol. 1. Cambridge: Harvard University Press, 90–147.
- Moore RP, Robinson WD. 2004. Artificial bird nests, external validity, and bias in ecological field studies. *Ecology* 85(6):1562–1567 DOI 10.1890/03-0088.
- Ng J, Landeen EL, Logsdon RM, Glor RE. 2013. Correlation between *Anolis* lizard dewlap phenotype and environmental variation indicates adaptive divergence of a signal important to sexual selection and species recognition. *Evolution* 67(2):573–582 DOI 10.1111/j.1558-5646.2012.01795.x.
- Ng J, Geneva AJ, Noll S, Glor RE. 2017. Signals and speciation: *Anolis* dewlap color as a reproductive barrier. *Journal of Herpetology* 51(3):437–447 DOI 10.1670/16-033.
- Nicholson KE, Harmon LJ, Losos JB. 2007. Evolution of *Anolis* lizard dewlap diversity. *PLOS ONE* 2(3):e274 DOI 10.1371/journal.pone.0000274.
- Ord TJ, Blumstein DT, Evans CS. 2001. Intrasexual selection predicts the evolution of signal complexity in lizards. *Proceedings of the Royal Society B: Biological Sciences* 268(1468):737–744 DOI 10.1098/rspb.2000.1417.
- Ord TJ, Martins EP. 2006. Tracing the origins of signal diversity in anole lizards: phylogenetic approaches to inferring the evolution of complex behaviour. *Animal Behaviour* 71(6):1411–1429 DOI 10.1016/j.anbehav.2005.12.003.
- Ord TJ, Peters RA, Clucas B, Stamps JA. 2007. Lizards speed up visual displays in noisy habitats. *Proceedings of the Royal Society B: Biological Sciences* 274(1613):1057–1062 DOI 10.1098/rspb.2006.0263.
- Ord TJ, Stamps JA, Losos JB. 2010. Adaptation and plasticity of animal communication in fluctuating environments. *Evolution* 64(11):3134–3148 DOI 10.1111/j.1558-5646.2010.01056.x.
- Orme D, Freckleton R, Thomas G, Petzoldt T, Fritz S, Isaac N, Pearse W. 2013. Caper: comparative analyses of phylogenetics and evolution in R. R package version 0.5.2. Available at <https://cran.r-project.org/web/packages/caper/> (accessed April 2016).
- Paluh DJ, Hantak MM, Saporito RA. 2014. A test of aposematism in the dendrobatid poison frog *Oophaga pumilio*: the importance of movement in clay model experiments. *Journal of Herpetology* 48(2):249–254 DOI 10.1670/13-027.
- R Development Core Team. 2016. *R: A Language and Environment for Statistical Computing*. Vienna: R Foundation for Statistical Computing. Available at <https://www.r-project.org/>.
- Rand AS, Williams EE. 1970. An estimation of redundancy and information content of anole dewlaps. *American Naturalist* 104(935):99–103 DOI 10.1086/282643.
- Rangen SA, Clark RG, Hobson KA. 2000. Visual and olfactory attributes of artificial nests. *Auk* 117(1):136–146 DOI 10.1642/0004-8038(2000)117[0136:vaoaoa]2.0.co;2.
- Revell LJ. 2010. Phylogenetic signal and linear regression on species data. *Methods in Ecology and Evolution* 1(4):319–329 DOI 10.1111/j.2041-210x.2010.00044.x.
- Revell LJ. 2012. Phytools: an R package for phylogenetic comparative biology (and other things). *Methods in Ecology and Evolution* 3(2):217–223 DOI 10.1111/j.2041-210x.2011.00169.x.
- Richards-Zawacki CL, Yeager J, Bart HPS. 2013. No evidence for differential survival or predation between sympatric color morphs of an aposematic poison frog. *Evolutionary Ecology* 27(4):783–795 DOI 10.1007/s10682-013-9636-0.
- Ricklefs RE, Bermingham E. 2004. History and the species-area relationship in Lesser Antillean birds. *American Naturalist* 163(2):227–239 DOI 10.1086/381002.

- Ricklefs RE, Lovette IJ. 1999. The roles of island area per se and habitat diversity in the species-area relationships of four Lesser Antillean faunal groups. *Journal of Animal Ecology* 68(6):1142–1160 DOI 10.1046/j.1365-2656.1999.00358.x.
- Roberts SC, Gosling LM, Thornton EA, McClung J. 2001. Scent-marking by male mice under the risk of predation. *Behavioral Ecology* 12(6):698–705 DOI 10.1093/beheco/12.6.698.
- Rodríguez-Cabrera TM, Torres J, Marrero R, Podio-Martínez JA. 2016. Predation attempt by the Cuban Racer, *Cubophis cantherigerus* (Squamata: Dipsadidae) on the Cuban Giant Anole, *Anolis equestris buidei* (Squamata: Dactyloidae), a threatened endemic subspecies. *IRCF Reptiles and Amphibians* 23(1):46–50.
- Rodríguez-Robles JA. 1992. Notes on the feeding behavior of the Puerto Rican Racer, *Alsophis portoricensis* (Serpentes: Colubridae). *Journal of Herpetology* 26(1):100–102 DOI 10.2307/1565036.
- Saks L, McGraw K, Horak P. 2003. How feather color reflects its carotenoid content. *Functional Ecology* 17(4):555–561 DOI 10.1046/j.1365-2435.2003.00765.x.
- Santos JC, Cannatella DC. 2011. Phenotypic integration emerges from aposematism and scale in poison frogs. *Proceedings of the National Academy of Sciences of the United States of America* 108(15):6175–6180 DOI 10.1073/pnas.1010952108.
- Schoener TW, Schoener A. 1980. Ecological and demographic correlates of injury rates in some Bahamian *Anolis* lizards. *Copeia* 1980(4):839–850 DOI 10.2307/1444463.
- Schoener TW, Schoener A. 1982. The ecological correlates of survival in some Bahamian *Anolis* lizards. *Oikos* 39(1):1–16 DOI 10.2307/3544525.
- Schoener TW, Slade JB, Stinson CH. 1982. Diet and sexual dimorphism in the very catholic lizard genus *Leiocephalus* of the Bahamas. *Oecologia* 53(2):160–169 DOI 10.1007/bf00545659.
- Schoener TW, Spiller DA, Losos JB. 2002. Predation on a common *Anolis* lizard: can the food-web effects of a devastating predator be reversed? *Ecological Monographs* 72(3):383–407 DOI 10.1890/0012-9615(2002)072[0383:poacal]2.0.co;2.
- Scott MP. 1984. Agonistic and courtship displays of male *Anolis sagrei*. *Breviora* 479:1–22.
- Sigmund WR. 1983. Female preferences for *Anolis carolinensis* males as a function of dewlap color and background coloration. *Journal of Herpetology* 17(2):137–143 DOI 10.2307/1563454.
- Simon V. 2007. Not all signals are equal: Male brown anole lizards (*Anolis sagrei*) electively decrease pushup frequency following a simulated predatory attack. *Ethology* 113(8):793–801 DOI 10.1111/j.1439-0310.2007.01379.x.
- Simon V. 2011. Communication signal rates predict interaction outcome in the brown anole lizard, *Anolis sagrei*. *Copeia* 2011(1):38–45 DOI 10.1643/ce-08-022.
- Smiseth PT, Örnborg J, Andersson S, Amundsen T. 2001. Is male plumage reflectance correlated with paternal care in bluethroats? *Behavioral Ecology* 12(2):164–170 DOI 10.1093/beheco/12.2.164.
- Smith JM, Harper D. 2003. *Animal Signals*. Oxford: Oxford University Press.
- Sokal RR, Rohlf FJ. 1995. *Biometry: The Principles and Practices of Statistics in Biological Research*. Third Edition. New York: Freeman WH.
- Stamps JA. 1983. Sexual selection, sexual dimorphism, and territoriality. In: Huey RB, Pianka ER, Schoener TW, eds. *Lizard Ecology: Studies of a Model Organism*. Cambridge: Harvard University Press, 169–204.
- Stamps JA. 1999. Relationships between female density and sexual size dimorphism in samples of *Anolis sagrei*. *Copeia* 1999(3):760–765 DOI 10.2307/1447610.

- Stamps JA, Losos JB, Andrews RM. 1997.** A comparative study of population density and sexual size dimorphism in lizards. *American Naturalist* **149**(1):64–90 DOI [10.1086/285979](https://doi.org/10.1086/285979).
- Steffen JE. 2009.** Perch-height specific predation on tropical lizard clay models: implications for habitat selection in mainland neotropical lizards. *Revista de Biología Tropical* **57**(3):859–864 DOI [10.15517/rbt.v57i3.5498](https://doi.org/10.15517/rbt.v57i3.5498).
- Steffen JE, McGraw KJ. 2007.** Contributions of pterin and carotenoid pigments to dewlap coloration in two anole species. *Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology* **146**(1):42–46 DOI [10.1016/j.cbpb.2006.08.017](https://doi.org/10.1016/j.cbpb.2006.08.017).
- Steinberg DS, Losos JB, Schoener TW, Spiller DA, Kolbe JJ, Leal M. 2014.** Predation-associated modulation of movement-based signals by a Bahamian lizard. *Proceedings of the National Academy of Sciences of the United States of America* **111**(25):9187–9192 DOI [10.1073/pnas.1407190111](https://doi.org/10.1073/pnas.1407190111).
- Stuart-Fox DM, Moussalli A, Whiting MJ. 2007.** Natural selection on social signals: signal efficacy and the evolution of chameleon display coloration. *American Naturalist* **170**(6):916–930 DOI [10.1086/522835](https://doi.org/10.1086/522835).
- Stuart-Fox DM, Ord TJ. 2004.** Sexual selection, natural selection and the evolution of dimorphic coloration and ornamentation in agamid lizards. *Proceedings of the Royal Society B: Biological Sciences* **271**(1554):2249–2255 DOI [10.1098/rspb.2004.2802](https://doi.org/10.1098/rspb.2004.2802).
- Telemeco RS, Baird TA, Shine R. 2011.** Tail waving in a lizard (*Bassiana duperreyi*) functions to deflect attacks rather than as a pursuit-deterrent signal. *Animal Behaviour* **82**(2):369–375 DOI [10.1016/j.anbehav.2011.05.014](https://doi.org/10.1016/j.anbehav.2011.05.014).
- Thompson FR, Burhans DE. 2004.** Differences in predators of artificial and real songbird nests: Evidence of bias in artificial nest studies. *Conservation Biology* **18**(2):373–380 DOI [10.1111/j.1523-1739.2004.00167.x](https://doi.org/10.1111/j.1523-1739.2004.00167.x).
- Trivers RL. 1976.** Sexual selection and resource-acruing abilities in *Anolis garmani*. *Evolution* **30**(2):253–269 DOI [10.1111/j.1558-5646.1976.tb00908.x](https://doi.org/10.1111/j.1558-5646.1976.tb00908.x).
- Turner FB, Medica PA, Jennrich RI, Maza BG. 1982.** Frequencies of broken tails among *Uta stansburiana* in Southern Nevada and a test of the predation hypothesis. *Copeia* **1982**(4):835–840 DOI [10.2307/1444094](https://doi.org/10.2307/1444094).
- Vanhooydonck B, Herrel A, Meyers JJ, Irschick DJ. 2009.** What determines dewlap diversity in *Anolis* lizards? An among-island comparison. *Journal of Evolutionary Biology* **22**(2):293–305 DOI [10.1111/j.1420-9101.2008.01643.x](https://doi.org/10.1111/j.1420-9101.2008.01643.x).
- Vanhooydonck B, Herrel A, Van Damme R, Irschick DJ. 2005.** Does dewlap size predict male bite performance in Jamaican *Anolis* lizards? *Functional Ecology* **19**(1):38–42 DOI [10.1111/j.0269-8463.2005.00940.x](https://doi.org/10.1111/j.0269-8463.2005.00940.x).
- Vanhooydonck B, Huyghe K, Holáňová V, van Dongen S, Herrel A. 2015.** Differential growth of naturally and sexually selected traits in an *Anolis* lizard. *Journal of Zoology* **296**(4):231–238 DOI [10.1111/jzo.12236](https://doi.org/10.1111/jzo.12236).
- Yee J, Lee J, Desowitz A, Blumstein DT. 2013.** The costs of conspecifics: Are social distractions or environmental distractions more salient? *Ethology* **119**(6):480–488 DOI [10.1111/eth.12085](https://doi.org/10.1111/eth.12085).
- York JR, Bairds TA. 2016.** Juvenile collared lizards adjust tail display frequency in response to variable predatory threat. *Ethology* **122**(1):37–44 DOI [10.1111/eth.12442](https://doi.org/10.1111/eth.12442).
- Zuk M, Tinghitella RM. 2008.** Rapid evolution and sexual signals. In: d’Ettorre P, Hughes DP, eds. *Sociobiology of Communication*. Oxford: Oxford University Press, 139–156.